

JAST (Journal of Animal Science and Technology) TITLE PAGE

Upload this completed form to website with submission

ARTICLE INFORMATION	Fill in information in each box below
Article Type	Research article
Article Title (within 20 words without abbreviations)	Enhanced γ -aminobutyric acid and sialic acid in fermented deer antler velvet and immune promoting effects
Running Title (within 10 words)	Fermented deer antler velvet extracts and immune promotion
Author	Jiseon Yoo ^{1#} , Juyeon Lee ^{1#} , Ming Zhang ² , Daye Mun ³ , Minkyong Kang ¹ , Bohyun Yun ¹ , Yong-An Kim ⁴ , Sooah Kim ² and Sangnam Oh ^{1*}
Affiliation	1 Department of Functional Food and Biotechnology, Jeonju University, Jeonju, 55069 Korea 2 Department of Environment Science & Biotechnology, Jeonju University, Jeonju 55069, Korea 3 Department of Agricultural Biotechnology and Research Institute of Agriculture and Life Science, Seoul National University, Seoul, Korea 4 Department of Animal Science, Chonbuk National University, Jeonju 54896, Korea # These authors contributed equally to this work.
ORCID (for more information, please visit https://orcid.org)	Jiseon Yoo (https://orcid.org/0000-0002-2402-5953) Juyeon Lee (https://orcid.org/0000-0002-6390-6120) Ming Zhang (https://orcid.org/0000-0001-9351-9837) Daye Mun (https://orcid.org/0000-0002-3470-7632) Minkyong Kang (https://orcid.org/0000-0002-2366-7970) Bohyun Yun (https://orcid.org/0000-0001-6723-5849) Yong-An Kim (https://orcid.org/0000-0001-5019-1536) Sooah Kim (https://orcid.org/0000-0002-9096-4879) Sangnam Oh (https://orcid.org/0000-0002-2428-412X)
Competing interests	No potential conflict of interest relevant to this article was reported.
Funding sources State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available.	This work was supported by grants from National Agricultural Cooperative Federation (NACF) and a grant from the Agricultural and livestock products safe distribution and consumption technology development program (Project No. 320111-1), Korea Institute of Planning and Evaluation for Technology in Food, Agriculture and Forestry (IPET) and the National Research Foundation of Korea Grant funded by the Korean government (NRF-2018R1D1A3B07050304).
Acknowledgements	Not applicable.
Availability of data and material	Upon reasonable request, the datasets of this study can be available from the corresponding author.
Authors' contributions Please specify the authors' role using this form.	Conceptualization: Oh S, Kim Y, Kim S Data curation: Yoo J, Lee J, Zhang M Methodology: Yoo J, Lee J, Zhang M, Oh S Software: Zhang M, Mun D, Kang M Validation: Kang M, Yun B, Kim Y, Kim S Investigation: Yoo J, Lee J, Zhang M, Mun D Writing - original draft: Yoo J, Lee J, Zhang M, Mun D, Kang M, Yun B, Kim Y, Kim S, Oh S Writing - review & editing: Yoo J, Lee J, Zhang M, Mun D, Kang M, Yun B, Kim Y, Kim S, Oh S
Ethics approval and consent to participate	The animal experiment in this research was approved by the Institutional Animal Care and Use Committee (IACUC) of the Korea Food Industry Cluster Promotion Agency based on the Animal Protection Act No. 16977.

5 **CORRESPONDING AUTHOR CONTACT INFORMATION**

For the corresponding author (responsible for correspondence, proofreading, and reprints)	Fill in information in each box below
First name, middle initial, last name	Sangnam Oh
Email address – this is where your proofs will be sent	osangnam@jj.ac.kr
Secondary Email address	osangnam@gmail.com
Address	Department of Functional Food and Biotechnology, Jeonju University, Jeonju, 55069 Korea
Cell phone number	+82-10-2535-5326
Office phone number	+82-63-220-3109
Fax number	+82-63-220-2054

10

15

20

ACCEPTED

25 ABSTRACT

Deer antler velvet is widely used in traditional medicine for its anti-aging, antioxidant, and immunity-enhancing effects. However, few studies have reported on the discovery of probiotic strains for deer antler fermentation to increase functional ingredient absorption. This study evaluated the ability of probiotic lactic acid bacteria to enhance the concentrations of bioactive molecules (e.g., sialic acid and gamma-aminobutyric acid [GABA]) in extracts of deer antler velvet. Seventeen strains of *Lactobacillus* spp. that were isolated from kimchi and infant feces, including *L. sakei*, *L. rhamnosus*, *L. brevis*, and *L. plantarum*, and those that improved the life span of *Caenorhabditis elegans* were selected for evaluation. Of the 17 strains, 2 (*L. rhamnosus* LFR20-004 and *L. sakei* LFR20-007) were selected based on data showing that these strains increased both the sialic acid and GABA contents of deer antler extract after fermentation for 2 d and significantly improved the life span of *C. elegans*. Co-fermentation with both strains further increased the concentrations of sialic acid, GABA, and metabolites such as short-chain fatty acids and amino acids. We evaluated the biological effects of the fermented antler velvet (FAV) on the antibacterial immune response in *C. elegans* by assessing worm survival after pathogen infection. The survival of the *C. elegans* conditioned with FAV for 24 h was significantly higher compared with that of the control worm group fed only normal feed (non-pathogenic *E. coli* OP50) exposed to *E. coli* O157:H7, *Salmonella typhi*, and *Listeria monocytogenes*. To evaluate the protective effects of FAV on immune response, cyclophosphamide (Cy), an immune-suppressing agent was treated to *in vitro* and *in vivo*. We found that FAV significantly restored viability of mice splenocytes and immune promoting-related cytokines (IL-6, IL-10, iNOS, IFN- γ , and TNF- α) were activated compared to non-fermented deer antlers. This finding indicated the protective effect of FAV against Cy-induced cell death and immunosuppressed mice. Taken together, our study suggests that immune-promoting antler velvet can be produced through fermentation using *L. rhamnosus* LFR20-004 and *L. sakei* LFR20-007.

Keywords: Deer antler velvet, Probiotics, Gamma-aminobutyric acid, Sialic acid, Immune promotion, *Caenorhabditis elegans*

INTRODUCTION

Deer antler velvet is a non-osteolytic agent that is widely used in traditional medicine because it contains a variety of small ingredients (e.g., monosaccharides, uronic acid, free amino acids, sialic acid, gangliosides, and small peptides) [1] that promote recovery from fatigue [2], improve liver
55 function and mitigate skeletal weakness symptoms [3, 4], strengthen immunity [5], and reduce inflammation. As the elderly population is increasing, demand for health supplements and import of antlers velvets have increased rapidly in recent years. New Zealand, Russia, Canada, and China are countries importing deer antlers, with New Zealand importing greatest quantity and Russia importing the most expensive ones [6]. In Asian countries, deer antler velvets are dried and sold as slices or usually
60 boiled with various types of herbs, botanical materials, and other ingredients in water to be consumed as medicinal soup. Despite having numerous beneficial ingredients, there are limited ways to consume deer antlers. Therefore, there have been various attempts to apply fermentation along with herb materials in order to increase the physiological efficacy of antler velvet and herbal medicine [7].

Microbial fermentation is used in the food industry for a variety of purposes, including the
65 enhancement of flavor, preservation, and health benefits [8]. It is also used to generate useful substances through the enzymatic decomposition of organic matter. During fermentation, high-molecular weight substances are converted into low-molecular weight substances, such as organic acids and amino acids, and glycosides are broken down so that nutrients can be more easily absorbed and utilized by the body [9]. Previous studies have reported that *Bacillus subtilis*-fermented antler extract promotes
70 hematopoiesis [10], inhibits osteoclast differentiation [11], and improves exercise endurance [12].

In this study, we focused on the optimal fermentation of deer antlers using probiotic strains, mainly the lactic acid bacteria (LAB). Fermentation with *Lactobacillus* sp. has been reported to enhance absorption of bioactive molecules, and aid in immune modulation, cholesterol homeostasis, exercise performance [13-16]. The purpose of the present study was to enhance the concentration of bioactive
75 ingredients (e.g., sialic acid and gamma-aminobutyric acid [GABA]) in fermented antler velvet (FAV) and evaluate its effects on immune response in *Caenorhabditis elegans* and mouse models.

MATERIALS AND METHODS

Caenorhabditis elegans longevity assay

80 The *fer-15;ferm-1* mutants were synchronized using an alkaline hypochlorite solution. *Lactobacillus rhamnosus* GG (LGG) and *Escherichia coli* OP50 (OP50) were used as controls. Concentrated suspensions of LAB or *E. coli* OP50 (100 μ L) were plated on 35-mm diameter nematode growth medium (NGM) agar plates. Day 0 was defined as the day of L4-to-adult switch. Worms were transferred, using a platinum wire, onto fresh plates daily and maintained at 25°C. The experiment was
85 repeated three times, and survival rates were determined.

Antler velvet extract fermentation

Water extracts of deer antler, *Cervus canadensis* Erxleben, were procured from Korea RND Nonghyup (Yeonghwasaro 24 Gwangjin-gu, Seoul, Korea) and stored at 4°C. To ferment with LAB
90 and antler extract, minimal prebiotic medium with 2% antler extract (1 L standard: 20 g antler water extract, 5 g peptone, 2.5 g sodium acetate, 0.5 mL of 1 M magnesium sulfate heptahydrate, 0.5 mL of 1 M manganese sulfate tetrahydrate, 5 mL Tween 80, 1 g diammonium citrate, 1 g dipotassium phosphate, 20 g glucose, and 1 g L-glutamic acid) was prepared and autoclaved at 121°C for 15 min. The working antler medium was aliquoted into 50-mL conical tubes, and 1% of each of the 17 LAB
95 strains was then inoculated ($>10^4$ CFU/mL). After incubation at 37°C for 24, 48, 72, 96, and 120 h, the cultures were centrifuged at 4000 \times g for 10 min at 25°C, and the resulting supernatants were filtered using 0.22- μ m syringe filters (Minisart, Sartorius Stedim Biotech GmbH, Goettingen, Germany).

Total acidity, pH, and bacterial growth

100 The acidity of each culture (1 mL) was measured by adding 10 μ L of 1% phenolphthalein and then adding 0.1 N NaOH until reaching an endpoint. The amount (mL) of 0.1 N NaOH consumed was then converted and expressed as lactic acid content (%). The culture pH was measured during

fermentation using a pH meter (S20 SevenEasy pH meter; Mettler-Toledo, Greifensee, Switzerland). The pH and acidity of the cultures were measured before and after each fermentation period (24, 48, 72, 105 96, and 120 h). Bacterial concentration was assessed using the colony-forming unit (CFU) counting method. Briefly, serial diluents (100 μ L) were spread on MRS agar plates, and CFUs (\log^{10} /mL) were counted after 2 d.

Total nitrogen, amino acid, and protein quantification

110 Total nitrogen content was analyzed photometrically as described previously [17], with minor modifications. Briefly, samples of each culture were diluted 25-fold, mixed with 5 mL aqueous HCl (0.2%), 20 mL ddH₂O, and 10 mL potassium persulfate (3% alkalinity, pH 9.7); autoclaved at 120°C for 30 min; cooled to room temperature; and then combined with 5 mL HCl (0.06%). Finally, the absorbance of each mixture was measured at 220 nm, and nitrogen content was determined using a 115 calibration curve generated using potassium nitrate. To measure total amino acid content, amino acids were extracted as described by Sun [3], with slight modifications. Briefly, 50 μ L of each culture was mixed with 2 mL ddH₂O, 1 mL acetic acid-sodium acetate buffer solution (pH 6.0), and 1 mL aqueous ninhydrin (2%); incubated at 100°C for 15 min; cooled to room temperature; and then mixed with 10 mL 50% ethanol. Finally, the absorbance of each mixture was measured at 220 nm [3], and amino acid 120 concentration was determined using a calibration curve generated using an arginine standard. The protein content of each culture was quantified using Bradford protein analysis, following the manufacturers' instructions. The concentration of each mixture was determined by measuring absorbance at 595 nm.

125 GABA analysis

GABA analysis was performed according to the method described by Lü et al. [18], with slight modifications. Briefly, 3 mL of each sample was mixed with 1.5 mL NaHCO₃ (0.5 M) and 0.5 mL 1-fluoro-2,4-dinitrobenzene (0.715 mg/mL), incubated at 60°C for 1 h, cooled to room temperature, and

130 then passed through a 0.22- μ m syringe filter (PTFE; Advantec, Dublin, CA, USA). The GABA content
of each sample derivative (10 μ L) was then analyzed using a Shimadzu 20A series HPLC system
(Shimadzu, Kyoto, Japan) equipped with a Symmetry C18 column (5 μ m, 3.9 \times 150 mm; Waters,
Milford, MA, USA). Isocratic elution was performed with 0.5% ammonium acetate aqueous solution
and acetonitrile (85:15, v/v) for 20 min at a flow rate of 1 mL/min. GABA was detected based on
absorption at a wavelength of 360 nm.

135

Sialic acid analysis

Each sample (0.1 mL) was mixed with 0.1 mL H₂SO₄ (0.2 N), hydrolyzed at 85°C for 1 h, and
then reacted with 20 μ L periodate solution (0.25 M NaIO₄ in 9 M H₃PO₄) at room temperature for 20
min. Next, 0.1 mL arsenite reagent (10% NaAsO₂ in 0.1 N H₂SO₄ with 0.5 M Na₂SO₄) was added to
140 reduce the excess periodate. Thereafter, 0.25 mL TBA solution (0.6% TBA in 0.5 M Na₂SO₄) was added
and color was developed at 85 °C for 15 min, followed by cooling at room temperature. Finally, each
sample was mixed with 1 mL cyclohexanone and centrifuged at 900 \times g for 7 min. The absorbance of
each supernatant was measured at 549 nm (BioTek Synergy HTX multi-mode plate reader, USA) [19],
and sialic acid content was calculated by comparison to a standard curve for N-acetyl neuraminic acid.

145

Metabolite analysis

The supernatants of the fermented samples were collected after centrifugation at 7000 rpm for
5 min and filtered using 0.2- μ m cellulose acetate syringe filters. Each sample was then mixed with ice-
cold methanol, vigorously vortexed on ice for 1 min, and centrifuged at 10,000 rpm for 10 min at 4°C.
150 The resulting supernatants were filtered using 0.2- μ m polyvinylidene fluoride (PVDF) syringe filters
(Whatman, Maidstone, England) and completely vacuum-dried.

For GC-MS analysis, each sample was dissolved in 30 μ L methoxyamine hydrochloride
(Sigma-Aldrich, St. Louis, MO, USA) in pyridine (20 mg/mL), derivatized at 30°C for 90 min. Samples
were then trimethylsilylated by adding 50 μ L *N,O*-bis(trimethylsilyl)trifluoroacetamide (Sigma-Aldrich,

155 St. Louis, MO, USA) and incubated at 60°C for 30 min. GC-MS analysis was performed using a
TRACE™ 1310 Gas Chromatograph (Thermo Fisher Scientific, Waltham, MA, USA) equipped with
an ISQ LT single quadrupole mass spectrometer (Thermo Fisher Scientific, Waltham, MA, USA), and
separation was performed using a DB-5MS column (60 m × 0.25 mm, 0.25-um film thickness; Agilent,
USA). Oven temperature was maintained at 50°C for 2 min, gradually increased at 5°C/min, maintained
160 at 180°C for 8 min, gradually increased at 2.5°C/min to 325°C, and then maintained for 10 min. The
sample was injected at 300°C, and helium was used as the carrier gas, with a flow rate of 1.5 mL/min
and split ratio of 1:60. An electron ionization system with an ionization voltage of 70 eV was used for
GC-MS detection, and the ion source temperature was 270 °C. The mass scan range was set at 30-450
(m/z), and the acquisition rate was 5 spectra/s. Detected metabolites were identified using the NIST
165 Mass Spectral Search Program (version 2.0; NIST, Gaithersburg, MD, USA).

***Caenorhabditis elegans* killing assay**

The *C. elegans* killing assay was performed as previously described [20]. L4/young adult
worms were placed on conditioning NGM plates with non-fermented antler velvet (AV) medium (10
170 mg/mL) or FAV (10 mg/mL), and non-pathogenic *E. coli* OP50 at 25°C for 24 h. Worms were plated
at 30 per plate. Subsequently, for viability analysis, nematodes preconditioned with AVM, FAV, and
OP50 were transferred onto *Salmonella typhi* SL 1344, *Listeria monocytogenes* EDG-e, and *E. coli*
O157:H7 EDL933 culture plates; incubated at 25°C; and examined using a microscope at 24-h intervals
until all worms died. Whether worms were alive was assessed by gentle touching with a platinum wire.

175

Isolation of mouse splenocytes

Spleen tissues were extracted from C57BL/6 mice (24-wk-old female, 20-22 g). The tissues
were disaggregated via passage through a 40 µm cell strainer in RPMI-1640 medium (SH30027,
Hyclone, USA) containing 10% FBS, 1% antibiotics and the cells were purified by centrifugation at
180 2000 rpm for 10 min at 4°C. Red blood cells were removed by incubating purified cells in Red blood

cell lysis buffer (A10492-0, GIBCO, USA) for 3 min. Splenocytes were then washed with RPMI containing 10% FBS and 1% antibiotics and centrifuged at 2000 rpm for 10 min at 4°C for 10 min.

Measurement of cell cytotoxicity and proliferation

185 The cell viability assay was performed using an EZ-Cytox cell viability assay kit (EZ-3000, DoGen, Seoul, Korea). Splenocytes were seeded into 96-well plates (approximately 5×10^6 /mL) and treated with 0, 0.1, 0.5, and 1 mg/mL of the AV or FAV samples for 48 h. Cells were co-treated with 1.6 mg/mL cyclophosphamide (Cy; C0768-1G; Sigma-Aldrich, St Louis, MO) and the antler velvet sample. Subsequently, 10 μ L of EZ-Cytox reagent was added to each well and incubated for 1 h.
190 Absorbance of the samples was then measured at 450 nm. Cell viability values were normalized to those of the cells from the untreated control groups.



Preparation of the Cy-induced immunosuppression mouse model

195 Ten-week-old female C57BL/6 mice (19-20 g) were housed in a room with controlled temperature ($22^\circ\text{C} \pm 2^\circ\text{C}$) and humidity ($50\% \pm 5\%$) on a 12/12 h light/dark cycle with free access to food and water. After 1 wk of acclimatization, the mice were randomly divided into four groups ($n = 5$): control group, Cy group (100 mg/kg), AV 2 mg/20 g group, FAV 2 mg/20 g group. The AV or FAV-treated groups were orally administered the corresponding concentration of antler velvet for about 28 d, and the control and the Cy groups received oral administration of the same amount of saline. Mice
200 were administered intraperitoneal injection of Cy (100 mg/kg) in a total volume of 200 μ L of saline on days 21 and 24. All of the mice were administered with Cy except the control group. At the end of the experiment, mice were sacrificed, and the spleens were collected.

RNA isolation and quantitative real-time reverse-transcription-polymerase chain reaction (RT-qPCR)

205

Total RNA was isolated from the spleen tissues using the total RNA Mini Kit (T2010S; New England BioLabs, Ipswich, MA), as per the manufacturer's instructions. A cDNA kit (170-8891, Bio-Rad Laboratories, Hercules, CA, USA) was used to synthesize cDNA from 1 µg of total RNA, using the primers. Target gene expression was analyzed using the gene-specific primers listed in
210 Supplementary Table 1. The amplified products were subjected to electrophoresis on a 1% agarose gel and visualized with ultraviolet illumination. For qPCR, cDNA strands synthesized from the isolated total RNA were used as templates. PCR was performed on a StepOnePlus Real-Time PCR System (Applied Biosystems), using the SYBR Green PCR Mastermix (1725121; Bio-Rad Laboratories, USA). The cycle threshold values were normalized against the β-actin gene expression levels.

215

Statistical analyses

Data were analyzed using GraphPad Prism software (Prism 9 Academic), and the results are presented as mean ± standard deviation (SD). Statistical analysis was performed using Student's *t*-test for comparisons between two groups and one-way analysis of variance for comparisons among multiple
220 groups. $p < 0.05$ was considered significant.

RESULTS AND DISCUSSION

225 Probiotic candidates for fermentation

Previously, we had isolated approximately 200 strains of LAB, including *L. rhamnosus* and *L. plantarum* strains, from fermented Kimchi and infant feces. Among the isolates, we screened the strains having basic probiotic properties for further fermentation with antler velvet extract. Of the 200 strains isolated from kimchi and infant feces, 17 *Lactobacillus* strains (*L. sakei*, *L. brevis*, *L. rhamnosus*, and
230 *L. plantarum*) were selected based on their superior acid resistance, bile resistance, and intestinal adhesion, as shown in Supplementary Table 2. All the isolates were identified by 16S rRNA sequencing.

The study of the selection of 17 strains as suitable candidates for deer antler extract fermentation is unpublished. To evaluate the activity of strains during fermentation with antler extract, 2% of velvet antler water extracts (water extract 50% w/w) containing minimal prebiotic media was used for fermentation. During the first three days of fermentation, all 17 strains maintained sufficient cell populations, lowered culture pH, and increased culture acidity (Table 1). The MR ID stains from infant feces exhibited more rapid pH lowering and higher acidity rather than LFR ID stains from Kimchi, whereas higher cell population was maintained during fermentation with Kimchi strains except for LFR20-006. There have been a few studies on the strengthening of bioactive ingredients of deer antlers via fermentation [10, 12, 21]. Kim et al. utilized 11 *Bacillus* sp., 4 *Lactobacillus* sp., and 6 mushroom stains (*Cordyceps militaris*) to evaluate the growth in antler medium. All *Lactobacillus* sp. (*L. plantarum*, *L. mesenteroides*, *L. delbrueckii*) failed to grow in both antler extract and antler power-based media [21]. Surprisingly, all the strains in the present study successfully exhibited high proliferation with 10^8 ~ 10^9 CFU/g, indicating that all strains were compatible with antler extract and proliferated and produced strain-specific metabolites.

We attempted to evaluate the longevity-promoting effects of all 17 selected strains using *C. elegans* to identify optimal strains for fermentation. Strains LFR20-003, LFR20-004, LFR20-007, LFR20-008, MR4, MR5, MR14, MR19, MR43, MR45, and MR50 were as efficacious if not more than the positive control *Lactobacillus rhamnosus* GG (Fig. 1). In our previous study, *Lactobacillus plantarum* JDFM216 was selected, and it was reported that LAB strains could mitigate immune function decline and cognitive decline due to aging [22, 23]. As an important industrial application of LAB, lactic acid fermentation continues to be a highly desirable method for processing and preserving non-dairy foods owing to its low cost, low energy requirements for both processing and preparing foods for consumption, high yield, and diversified flavors as compared to fermentation with mold or *Bacillus* sp. [24].

255

Effect of fermentation on bioactive molecule contents

On the second day after fermentation with antler velvet, the amounts of total nitrogen, amino

acid, proteins, and bioactive substances were measured. In most fermented extracts, the amounts of total nitrogen, protein, and amino acids decreased (Supplementary Table 3). In general, *Lactobacillus* has the ability to hydrolyze proteins present in the medium, and this proteolytic activity produces a variety of peptides, metabolites with biological activity, as well as free amino acids required by bacteria. [25, 26]. In this experiment, it was found that even for the same *Lactobacillus* species, the protein utilization rate was variable. This result indicates the diversity in mechanisms involved in the longevity-promoting properties in *C. elegans*.

LAB producing and releasing GABA, a neurotransmitter, have been reported, and their mechanisms were reviewed [27-29]. Although there are reports of the effect of deer antler extract [30] and bone extract [31] on memory improvement in scopolamine-induced memory impaired mice, the production of GABA in deer antler extract after fermentation has not yet been investigated. The GABA content of the antler velvet was increased six times when fermented by MR43, MR45, and MR50, which were all isolated from infant feces, and was increased three and two times when fermented by LFR20-007 and LFR20-004, respectively, which were isolated from kimchi (Fig. 2A). In addition, the amount of ganglioside, a major bioactive component of deer antlers, was measured. Sialic acid in antler velvet has been considered a biomarker among bioactive molecules. Various extraction methods utilizing enzymes and fermentation, amongst others, have been developed to improve the recovery of sialic acid [32-34]. All the selected strains increased the sialic acid content; however, fermentation with LFR20-006, LFR20-007, and LFR20-008 increased the sialic acid content by approximately eight-fold (Fig. 2B). Interestingly, after fermentation, the levels of sialic acid, rather than GABA, in most fermented extracts were increased dramatically. Park et al. (2015) reported that the sialic acid content was highest, exhibiting a four-fold increase approximately in *Bacillus subtilis*-FAV extract on the fourth day [10]. The 11 LAB strains in this study (LFR20-001, -003, -004, -005, -006, -007, -008, -009, MR4, MR5, and MR19) exhibited greater ability to produce sialic acid in fermented deer antler.

Effect of FAV extract on *C. elegans* infected with pathogens

We then compared the biological activity of FAV with AV. We utilized two strains, LFR20-004 and LFR20-007, among the strains that increased the longevity in *C. elegans* (10 strains), GABA production (6 strains), and sialic acid content (10 strains) for the fermentation (Fig. 2C). GC-MS analysis of FAV revealed that free amino acid, short-chain fatty acid, and monosaccharide contents were changed significantly (Table 2). This profile may include metabolites derived from two strains and ingredients used by LAB or not used in AV samples.

We evaluated the biological effects of FAV on the antibacterial immune response in *C. elegans* by assessing whether the FAV and AV with probiotics influenced the host's survival after infection by pathogens (*E. coli* O157:H7, *Salmonella typhi*, and *Listeria monocytogenes*). The survival of the *C. elegans* conditioned with FAV or AV for 24 h was significantly higher compared with that of the control worm group fed only normal feed (non-pathogenic *E. coli* OP50) when exposed to two pathogens (*E. coli* O157:H7 and *Listeria monocytogenes*) (Fig. 3A and 3B). The worms conditioned with FAV exhibited significant resistance to infection by *S. typhi*, while worms conditioned with AV were not different from those fed normal feed (Fig. 3C). There have been few studies on the antibacterial efficacy of deer extracts against pathogens. To our knowledge, this study is the first to report that non-fermented deer antlers increase resistance of *C. elegans* to *E. coli* O157:H7 and *Listeria monocytogenes*. Dai et al. (2011) reported that antler velvet extracts stimulated anti-inflammatory cytokines in splenocytes and phagocytes and play a protective role in *Staphylococcus aureus*-infected mice [35]. However, fermentation products are known to have excellent antibacterial efficacy owing to the production of short-chain fatty acids such as lactic acid and play an important role in the safe storage of fermented foods [36-39]. Our results showed that the FAV increased resistance of *C. elegans* to pathogenic bacteria.

Cytoprotective effect of FAV on mouse splenocytes

To evaluate the effects of FAV and AV on cytotoxicity of Cy, an immune-suppressing agent, to primary splenocytes, cells isolated from spleens of mice were incubated with various concentrations

310 of AV or FAV (0, 0.1, 0.5, or 1 mg/mL) and Cy (1.6 mg/mL) for 48 h. Surprisingly, we found that FAV completely restored cell viability at 0.1 mg/mL, which was the lowest concentration. However, the viability of splenocytes treated with AV remained reduced by Cy (Fig. 4A). This finding indicated the protective effect of FAV against Cy-induced cell death and encouraged us to induce immunosuppressed mice by administering Cy (100 mg/kg). Expression of various immune modulators such as interleukin (IL)-6, IL-10, interferon (IFN)- γ , inducible nitric oxide synthase (iNOS), and tumor necrosis factor (TNF)- α by activated macrophages has been reported to play an important role in host defense against harmful factors [40]. Cytokines such as IL-6 and TNF- α exhibit essential immune functions during early stages of infection [41, 42]. IFN- γ , which is a type-1 T-helper (Th1)-derived cytokine, also plays important roles in cell-mediated immune response [43]. IL-6 and IL-10 are type-2 T-helper (Th2)-
320 derived cytokines that promote humoral immunity [44]. NO is synthesized by iNOS, an important molecule that activates macrophages to neutralize pathogenic microorganisms and tumor cells [45]. We analyzed mRNA expression of immune modulators such as IL-6, IL-10, iNOS, IFN- γ , and TNF- α in splenocytes isolated from immunosuppressed mice. We observed that FAV promoted the production of immunomodulators in spleen tissue-derived cells while AV did not (Fig.4). In addition to primary
325 splenocytes, it is necessary to investigate the effect of FAV on the single type of isolated immune cells, such as natural killer cells, T/B cells, neutrophils, eosinophils, and dendritic cells, to describe the biological mechanisms for the immune-enhancing activity of FAV.

The health-promoting effects of antler velvet in terms of chemical composition, bioactive components, and pharmacological effects have been reported [46]; however, there is not adequate
330 evidence for its immune-enhancing efficacy. Zhou et al. have suggested that peptides (3.2 kDa) extracted from antler velvet can modulate the immune system via regulation of Th1- and Th2-related cytokines [47]. Consistently, in our study, antler velvet fermented with two bacterial strains stimulated the immune system while AV did not.

The global demand for health- and immune-boosting functional foods has increased in the
335 COVID-19 era, and studies on the development of immune-promoting substances and food processing

are also increasing [48]. In this study, we demonstrated that FAV increased resistance to pathogenic bacteria in *C. elegans* and promoted splenocyte activation in mouse models by stimulating immune modulators. As deer antler velvet is fermented with LFR20-004 and LFR20-007, the bioactive molecules (GABA, sialic acid) as well as the immune function of deer antler velvet is expected to improve. In the future, an in-depth analysis of the metabolites from the LAB, *L. rhamnosus* LFR20-004 and *L. sakei* LFR20-007 that exhibit immune-promoting functions will be required. These LAB can be used as probiotic starters for health functional foods fermented for deer antler velvet as LAB.

ACKNOWLEDGMENTS

This work was supported by grants from National Agricultural Cooperative Federation (NACF) and a grant from the Agricultural and livestock products safe distribution and consumption technology development program (Project No. 320111-1), Korea Institute of Planning and Evaluation for Technology in Food, Agriculture and Forestry (IPET) and the National Research Foundation of Korea Grant funded by the Korean government (NRF-2018R1D1A3B07050304).

AUTHOR CONTRIBUTIONS

Conceptualization: Oh S, Kim Y, Kim S

Data curation: Yoo J, Lee J, Zhang M

Methodology: Yoo J, Lee J, Zhang M

Software: Zhang M, Mun D, Kang M

Validation: Kang M, Yun B, Kim Y, Kim S

Investigation: Yoo J, Lee J, Zhang M, Mun D

Writing - original draft: Yoo J, Lee J, Zhang M, Mun D, Kang M, Yun B, Kim Y, Kim S, Oh S

Writing - review & editing: Yoo J, Lee J, Oh S

ETHICS APPROVAL

The animal experiment in this research was approved by the Institutional Animal Care and Use Committee (IACUC) of the Korea Food Industry Cluster Promotion Agency based on the Animal Protection Act No. 16977.

365

ACCEPTED

REFERENCES

1. Zhou QL, Liu YQ, Wang Y, Guo YJ, Wang BX. [A comparison of chemical composition and bioactivity of polypeptides from velvet antlers of *Cervus nippon* Temminck and *Cervus elaphus* Linnaeus]. *Zhongguo Zhong Yao Za Zhi*. 2001;26(10):699-702.
370
2. Chen J-C, Hsiang C-Y, Lin Y-C, Ho T-Y. Deer Antler Extract Improves Fatigue Effect through Altering the Expression of Genes Related to Muscle Strength in Skeletal Muscle of Mice. *Evid Based Complement Alternat Med*. 2014;2014:540580-.
3. Sun S-W, Lin Y-C, Weng Y-M, Chen M-J. Efficiency improvements on ninhydrin method for amino acid quantification. *Journal of Food Composition and Analysis*. 2006;19(2):112-7.
375
4. Kim D-K, Lee S-H, Lee E-D, Kim H-J, Lee J, Lee S-S, et al. Characterizing Sections of Elk Velvet Antler by pH and Mineral, Fatty Acid, and Amino Acid Composition. *Journal of Food and Nutrition Research*. 2021;9(3):170-6.
5. Kim Y-A, Kim S-W, Lee M-H, Lee H-K, Hwang I-H. Comparisons of Chemical Composition, Flavor and Bioactive Substances between Korean and Imported Velvet Antler Extracts. *Food Science of Animal Resources*. 2021;41(3):386-401.
380
6. Kyung-Yun Choi SYH, Su-Chan Lee and Si-Kwan Kim. Effect of Antler Velvet Ethanol Extract on Common Serum Chemistry Panels and Histopathological Change in Rats Exposed to 2,3,7,8-Tetrachlorodibenzo-p-dioxin. *Korean Soc Food Sci Nutr*. 2006;35(9):1178-84.
7. Jung Y-J, Han, Dong-Oh, Park, Chul, Lee, Hye-Jung, Kim, Sung-Hoon, Hahm, Dae-Hyun. Effect of Fermented Herbal Extracts, HP-1 on Enzyme Activities and Gene Expressions Related to Alcohol Metabolism in Ethanol-loaded Rats. *Journal of Physiology & Pathology in Korean Medicine*. 2007.
385
8. Leroy F, De Vuyst L. Lactic acid bacteria as functional starter cultures for the food fermentation industry. *Trends in Food Science & Technology*. 2004;15(2):67-78.
9. Yadav H, Jain S, Rastamanesh R, Bomba A, Catanzaro R, Marotta F. Fermentation Technology in the Development of Functional Foods for Human Health: Where We Should Head. *Ferment Technol*. 2012;1(1).
390
10. Park Y, Choi HS, Lee HS, Suh HJ. Hematopoietic effect of deer antler extract fermented by *Bacillus subtilis* on murine marrow cells. *Nutr Res Pract*. 2015;9(5):451-8.
11. Choi SW, Moon SH, Yang HJ, Kwon DY, Son YJ, Yu R, et al. Antiresorptive activity of bacillus-fermented antler extracts: inhibition of osteoclast differentiation. *Evid Based Complement Alternat Med*. 2013;2013:748687.
395
12. Jang S, Park ED, Suh HJ, Lee SH, Kim JS, Park Y. Enhancement of exercise endurance capacity by fermented deer antler in BALB/c mice. *Bioscience, Biotechnology, and Biochemistry*. 2014;78(10):1716-22.
13. Zhao Z, Wu X, Chen H, Liu Y, Xiao Y, Chen H, et al. Evaluation of a strawberry fermented beverage with potential health benefits. *PeerJ*. 2021;9:e11974.
400
14. Lim PS, Loke CF, Ho YW, Tan HY. Cholesterol homeostasis associated with probiotic supplementation in

vivo. J Appl Microbiol. 2020;129(5):1374-88.

15. Widyastuti Y, Febrisiantosa A, Tidona F. Health-Promoting Properties of Lactobacilli in Fermented Dairy Products. *Front Microbiol.* 2021;12:673890.
- 405 16. Huang CC, Chen YM, Kan NW, Chao HL, Ho CS, Hsu MC. Cornu cervi pantotrichum supplementation improves exercise performance and protects against physical fatigue in mice. *Molecules.* 2014;19(4):4669-80.
17. Muntean A, Smical I, Török Z. Validation of an Alternative Method for Total Nitrogen Analysis in Water Samples. *Studia Universitatis Babeş-Bolyai Chemia.* 2017;62:101-10.
- 410 18. Lü Y, Zhang H, Meng X, Wang L, Guo X. A Validated HPLC Method for the Determination of GABA by Pre-Column Derivatization with 2,4-Dinitrofluorodinitrobenzene and Its Application to Plant GAD Activity Study. *Analytical Letters.* 2010;43(17):2663-71.
19. Kim S-H, Ameer K, Oh J-H, Park M-K. Characterization of hot water extract from Korean deer velvet antler (*Cervus canadensis* Erxleben). *Korean Journal of Food Preservation.* 2020;27(6):725-33.
- 415 20. Kim Y, Mylonakis E. *Caenorhabditis elegans* immune conditioning with the probiotic bacterium *Lactobacillus acidophilus* strain NCFM enhances gram-positive immune responses. *Infection and immunity.* 2012;80(7):2500-8.
21. Kim M-K, Jung E-Y, Lee H-S, Shin K-S, Kim Y-K, Ra K-S, et al. Isolation of strain for the preparation of the fermented antler and its physiological activities. *Journal of the Korean Society of Food Science and Nutrition.* 2009;38(9):1237-42.
- 420 22. Park MR, Ryu S, Maburutse BE, Oh NS, Kim SH, Oh S, et al. Probiotic *Lactobacillus fermentum* strain JDFM216 stimulates the longevity and immune response of *Caenorhabditis elegans* through a nuclear hormone receptor. *Sci Rep.* 2018;8(1):7441.
23. Park MR, Shin M, Mun D, Jeong S-Y, Jeong D-Y, Song M, et al. Probiotic *Lactobacillus fermentum* strain JDFM216 improves cognitive behavior and modulates immune response with gut microbiota. *Scientific Reports.* 2020;10(1):21701.
- 425 24. Canon F, Nidelet T, Guedon E, Thierry A, Gagnaire V. Understanding the Mechanisms of Positive Microbial Interactions That Benefit Lactic Acid Bacteria Co-cultures. *Front Microbiol.* 2020;11:2088.
25. Raveschot C, Cudennec B, Coutte F, Flahaut C, Fremont M, Drider D, et al. Production of Bioactive Peptides by *Lactobacillus* Species: From Gene to Application. *Front Microbiol.* 2018;9:2354.
- 430 26. Ming T, Han J, Li Y, Lu C, Qiu D, Li Y, et al. A metabolomics and proteomics study of the *Lactobacillus plantarum* in the grass carp fermentation. *BMC Microbiol.* 2018;18(1):216.
27. Cho YR, Chang JY, Chang HC. Production of gamma-aminobutyric acid (GABA) by *Lactobacillus buchneri* isolated from kimchi and its neuroprotective effect on neuronal cells. *Journal of microbiology and biotechnology.* 2007;17(1):104-9.
- 435

28. Patterson E, Ryan PM, Wiley N, Carafa I, Sherwin E, Moloney G, et al. Gamma-aminobutyric acid-producing lactobacilli positively affect metabolism and depressive-like behaviour in a mouse model of metabolic syndrome. *Sci Rep*. 2019;9(1):16323.
- 440 29. Cui Y, Miao K, Niyaphorn S, Qu X. Production of Gamma-Aminobutyric Acid from Lactic Acid Bacteria: A Systematic Review. *Int J Mol Sci*. 2020;21(3).
30. Lee M, Sun B, Gu L, Wang C, Fang Z, Wang Z, et al. Effects of the Deer Antler Extract on Scopolamine-induced Memory Impairment and Its Related Enzyme Activities. *The Korean Society of Food Science and Nutrition*. 2009;30:409–14.
- 445 31. Du CN, Min AY, Kim HJ, Shin SK, Yu HN, Sohn EJ, et al. Deer bone extract prevents against scopolamine-induced memory impairment in mice. *J Med Food*. 2015;18(2):157-65.
32. Jang DW, Ameer K, Oh JH, Park MK. Optimization and Pretreatment for Hot Water Extraction of Korean Deer (*Cervus canadensis* Erxleben) Velvet Antlers. *Journal of microbiology and biotechnology*. 2020;30(8):1116-23.
- 450 33. Lee SH, Yang HW, Ding Y, Wang Y, Jeon YJ, Moon SH, et al. Anti-inflammatory effects of enzymatic hydrolysates of velvet antler in RAW 264.7 cells in vitro and zebrafish model. *EXCLI J*. 2015;14:1122-32.
34. Je J-Y, Park P-J, Kim E-K, Kim H-A, Lim D-H, Jeon B-T, et al. Composition of Biologically Active Substances and Antioxidant Activity of New Zealand Deer Velvet Antler Extracts. *Food Science of Animal Resources*. 2010;30(1):20-7.
- 455 35. Dai TY, Wang CH, Chen KN, Huang IN, Hong WS, Wang SY, et al. The Antiinfective Effects of Velvet Antler of Formosan Sambar Deer (*Cervus unicolor swinhoei*) on *Staphylococcus aureus*-Infected Mice. *Evid Based Complement Alternat Med*. 2011;2011:534069.
36. Aldunate M, Srbinovski D, Hearps AC, Latham CF, Ramsland PA, Gugasyan R, et al. Antimicrobial and immune modulatory effects of lactic acid and short chain fatty acids produced by vaginal microbiota associated with eubiosis and bacterial vaginosis. *Frontiers in Physiology*. 2015;6(164).
- 460 37. Gao Z, Daliri EB, Wang J, Liu D, Chen S, Ye X, et al. Inhibitory Effect of Lactic Acid Bacteria on Foodborne Pathogens: A Review. *J Food Prot*. 2019;82(3):441-53.
38. Rhee SJ, Lee JE, Lee CH. Importance of lactic acid bacteria in Asian fermented foods. *Microb Cell Fact*. 2011;10 Suppl 1:S5.
- 465 39. Daliri F, Aboagye AA, Daliri EB-M. Inactivation of Foodborne Pathogens by Lactic Acid Bacteria. *Journal of Food Hygiene and Safety*. 2020;35(5).
40. Geum NG, Eo HJ, Kim HJ, Park GH, Son HJ, Jeong JB. Immune-enhancing activity of *Hydrangea macrophylla* subsp. *serrata* leaves through TLR4/ROS-dependent activation of JNK and NF- κ B in RAW264.7 cells and immunosuppressed mice. *Journal of Functional Foods*. 2020;73:104139.
- 470 41. Lee ES, Ju HK, Moon TC, Lee E, Jahng Y, Lee SH, et al. Inhibition of nitric oxide and tumor necrosis factor-alpha (TNF-alpha) production by propenone compound through blockade of nuclear factor (NF)-kappa B activation in cultured murine macrophages. *Biol Pharm Bull*. 2004;27(5):617-20.

42. Higuchi M, Higashi N, Taki H, Osawa T. Cytolytic mechanisms of activated macrophages. Tumor necrosis factor and L-arginine-dependent mechanisms act synergistically as the major cytolytic mechanisms of activated macrophages. *J Immunol.* 1990;144(4):1425-31.
- 475 43. Mosmann T, Coffman R. TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. *Annual review of immunology.* 1989;7(1):145-73.
44. Constant SL, Bottomly K. Induction of Th1 and Th2 CD4+ T cell responses: the alternative approaches. *Annual review of immunology.* 1997;15(1):297-322.
- 480 45. Ren Z, Qin T, Qiu F, Song Y, Lin D, Ma Y, et al. Immunomodulatory effects of hydroxyethylated *Hericium erinaceus* polysaccharide on macrophages RAW264. 7. *International journal of biological macromolecules.* 2017;105:879-85.
46. Sui Z, Zhang L, Huo Y, Zhang Y. Bioactive components of velvet antlers and their pharmacological properties. *J Pharm Biomed Anal.* 2014;87:229-40.
- 485 47. Zha E, Li X, Li D, Guo X, Gao S, Yue X. Immunomodulatory effects of a 3.2kDa polypeptide from velvet antler of *Cervus nippon Temminck*. *Int Immunopharmacol.* 2013;16(2):210-3.
48. Han B, Hoang BX. Opinions on the current pandemic of COVID-19: Use functional food to boost our immune functions. *J Infect Public Health.* 2020;13(12):1811-7.

490

ACCEPTED

Table 1. Changes of pH, acidity and LAB proliferation during the fermentation of antler velvet extract

Strain ID	¹ pH			¹ Total acidity (%)			² Log CFU/mL		
	1d	2d	3d	1d	2d	3d	1d	2d	3d
LFR20-001	4.50	4.09	3.99	0.45	0.54	0.63	7.31 ± 0.05	7.30 ± 0.07	7.17 ± 0.05
LFR20-002	5.33	3.52	3.46	0.32	1.01	1.13	5.12 ± 0.05	8.91 ± 0.12	8.58 ± 0.34
LFR20-003	4.14	3.78	3.60	0.60	0.77	0.99	8.26 ± 0.24	7.92 ± 0.08	8.30 ± 0.30
LFR20-004	4.14	3.75	3.60	0.52	0.76	0.99	8.36 ± 0.32	8.43 ± 0.38	7.90 ± 0.07
LFR20-005	5.31	3.58	3.44	0.28	0.92	1.22	6.32 ± 0.28	9.12 ± 0.05	8.30 ± 0.00
LFR20-006	4.19	4.03	4.01	0.60	0.81	0.63	7.71 ± 0.04	6.96 ± 0.16	3.47 ± 0.12
LFR20-007	4.82	3.75	3.41	0.36	0.85	0.99	8.46 ± 0.41	9.08 ± 0.11	8.59 ± 0.11
LFR20-008	4.09	3.76	3.62	0.69	0.87	0.99	8.39 ± 0.36	8.42 ± 0.10	7.83 ± 0.06
LFR20-009	4.13	3.56	3.56	0.72	1.13	1.26	8.84 ± 0.18	9.08 ± 0.04	8.79 ± 0.10
MR1	3.77	3.42	3.57	0.72	1.15	1.08	8.94 ± 0.21	8.59 ± 0.11	5.63 ± 0.04
MR4	3.77	3.43	3.41	0.74	1.13	1.04	9.17 ± 0.10	8.72 ± 0.12	5.66 ± 0.07
MR5	3.67	3.51	3.48	0.82	1.10	1.01	8.93 ± 0.08	8.46 ± 0.15	5.22 ± 0.14
MR14	3.66	3.50	3.48	0.91	1.10	1.26	9.08 ± 0.07	8.66 ± 0.39	5.05 ± 0.25
MR19	3.66	3.47	3.48	0.86	1.04	1.22	9.12 ± 0.15	8.43 ± 0.23	5.48 ± 0.07
MR43	3.42	3.35	3.23	0.81	0.95	1.08	8.78 ± 0.26	8.26 ± 0.24	7.11 ± 0.03
MR45	3.42	3.30	3.27	0.86	0.81	1.06	8.76 ± 0.15	8.30 ± 0.30	8.00 ± 0.17
MR50	3.40	3.30	3.25	0.81	1.04	1.13	8.74 ± 0.24	8.36 ± 0.10	7.01 ± 0.15

¹ pH and total acidity were measured continuously over 1, 2, and 3 days for independent samples

495 fermented with each strain, and are presented as average values.

² The probiotic water was measured three times per 1 mL of fermented product over 1, 2, and 3 days and presented as a log value.


Table 2. Profiling metabolites of fermentation in antler velvet extract

	¹ PC1	<i>p</i> -value	² FC
Metabolites increased by fermentation			
D-Lactic acid	0.990	<0.001	287.20
L-Proline	0.984	0.005	+
Mevalonic lactone	0.979	0.006	+
Succinic acid	0.979	0.007	+
Glutaric acid	0.968	0.01	+
Glycine	0.955	0.009	6.091
D-Mannitol	0.945	0.013	+
Propanoic acid	0.937	0.028	+
Phenylalanine	0.920	0.040	+
Aminomalonic acid	0.907	0.053	+
Metabolites reduced by fermentation			
d-Galactose	-0.908	0.040	3.712
Mannose	-0.928	0.010	1.835
D-(-)-Tagatose	-0.971	<0.001	-
d-Galactose	-0.983	0.005	1.735

¹ It was listed in the order of the most influential metabolites used as PC1 for principal component analysis (PCA)

² The numerical value is fold change (FC), the plus means metabolites detected in FAV and the minus means metabolites not-detected in FAV, but in AV

Figure Legends



505

Figure 1. Anti-aging effects of *C. elegans* by exposure to *Lactobacillus* spp. Prolonged lifespan of *C. elegans* by pre-conditioning with each strain (red line) for 24h. (A) *Lactobacillus rhamnosus* GG (LGG), (B) *Lactobacillus sakei* (LFR20-001), (C) *Lactobacillus brevis* (LFR20-002), (D) *Lactobacillus rhamnosus* (LFR20-003), (E) *Lactobacillus rhamnosus* (LFR20-004), (F) *Lactobacillus sakei* (LFR20-005), (G) *Lactobacillus rhamnosus* (LFR20-006), (H) *Lactobacillus sakei* (LFR20-


510

007), (I) *Lactobacillus brevis* (LFR20-008), (J) *Lactobacillus brevis* (LFR20-009), (K) *Lactobacillus plantarum* (MR4), (L) *Lactobacillus plantarum* (MR5), (M) *Lactobacillus plantarum* (MR14), (N) *Lactobacillus plantarum* (MR19), (O) *Lactobacillus plantarum* (MR43), (P) *Lactobacillus plantarum* (MR45), (Q) *Lactobacillus plantarum* (MR50)

515

ACCEPTED

Figure 2




520

Figure 2. Changes for amounts of bioactive ingredients and optimal selection strains for fermented antler velvet. (A) GABA content, (B) Sialic acid content, (C) Venn Diagram produced by the online software Venny 2.1.0 (<http://bioinfo.pcnb.csic.es/tools/venny/index.html>) to select optimal strains for FAV.

525

ACCEPTED

Figure 3



530 **Figure 3.** Killing assays of *C. elegans* worms fed with the FAV. *C. elegans* preconditioned with AV (10 mg/mL), FAV (10 mg/mL), and OP50 for 24 h were transferred onto each pathogen containing solid medium (A) *E. coli* O157: H7 EDL933 (ATCC 700927), (B) *Listeria monocytogenes* EGD-e (ATCC BAA-679) and (C) *Salmonella typhi* SL1344 (ATCC 14028).

Figure 4

535




Figure 4. Effects of FAV on the viability of splenocytes and on mRNA expression of immune promoting cytokine in CY-treated mice. (A) Splenocytes were treated with AV, FAV (0, 0.1, 0.5 and 1 mg/mL) and/or Cy (1.6 mg/mL). All results were expressed compared to untreated and shown as the means ± standard deviation (SD) (n=3). (B~F) As an immune promotion indicator, the expression of mRNA in cytokines as follows was expressed as a fold change to untreated (NC). (B) IL-6, (C) iNOS, (D) IL-10, (E) IFN-γ, and (F) TNF-α. Values are presented as mean ± standard deviation (n = 4). * P ≤ 0.05 ; compared with NC group, # P ≤ 0.05 ; compared with Cy group, \$ P ≤ 0.05 ; compared with Cy + AV group.

545