

c17-C18:1	0.224	0.222	0.00	0.772
c12,15-C18:2	0.041	0.037	0.00	0.057
n6-C18:2	1.869	1.885	0.03	0.673
C20:0	0.127	0.125	0.00	0.718
n6-C18:3	0.025	0.024	0.00	0.530
t11-C20:1	0.008	0.007	0.00	0.384
c11-C20:1	0.081	0.082	0.00	0.819
n3-C18:3	0.261	0.257	0.01	0.675
c9t11-C18:2	0.576	0.566	0.02	0.774
t10c12-C18:2	0.007	0.006	0.00	0.700
C21:0	0.015	0.015	0.00	0.861
c9c11-C18:2	0.007	0.006	0.00	0.745
C20:2	0.031	0.032	0.00	0.727
C22:0	0.036	0.034	0.00	0.462
n6-C20:3	0.083	0.083	0.00	0.781
n6-C20:4	0.106	0.108	0.00	0.399
C23:0	0.015	0.016	0.00	0.958
n3-20:5	0.022	0.024	0.00	0.034
C24:0	0.024	0.023	0.00	0.557
C22:4	0.018	0.017	0.00	0.526
n3-C22:5	0.047	0.046	0.00	0.742
De novo	25.17	25.38	0.31	0.473

Control, no treatment; Vinegar; 2 of 4.5% acetic acid equivalent per day

SEM, standard error of the mean

Proportion of fatty acids from de novo synthesis was calculated as the sum of 4- to 16-carbon fatty acids, odd and branched chain fatty acids were excluded.