

JAST (Journal of Animal Science and Technology) TITLE PAGE
Upload this completed form to website with submission

ARTICLE INFORMATION	Fill in information in each box below
Article Type	Research article
Article Title (within 20 words without abbreviations)	Complete genome sequence of <i>Lactobacillus curvatus</i> CACC879 and its functional probiotic properties
Running Title (within 10 words)	Complete genome sequence of <i>Lactobacillus curvatus</i> CACC879
Author	Soyeon Park ¹ , Seoyun Son ¹ , Mi Ae Park ¹ , Dae-Hyuk Kim ^{1,2} , and Yangseon Kim ¹
Affiliation	¹ Department of Research and Development, Center for Industrialization of Agricultural and Livestock Microorganisms, Jeongeup, 56212, Korea ² Department of Molecular Biology, Department of Bioactive Material Science, Institute for Molecular Biology and Genetics, Jeonbuk National University, Jeonju, 54896, Korea
ORCID (for more information, please visit https://orcid.org)	Soyeon Park (http://orcid.org/0000-0003-3788-5415) Seoyun Son (http://orcid.org/0000-0002-4753-1955) Mi Ae Park (http://orcid.org/0000-0002-7601-5976) Dae-Hyuk Kim (http://orcid.org/0000-0002-9948-5313) Yangseon Kim (http://orcid.org/0000-0002-8285-3407)
Competing interests	No potential conflict of interest relevant to this article was reported.
Funding sources State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available.	This research was supported by INNOPOLIS FOUNDATION through Science and Technology Project Opens the Future of the Region, which is funded by Ministry of Science and ICT (2022-DD-UP-0333), Korea, and partially supported by 2022 Technology commercialization support project Ministry of Agriculture, Food and Rural Affairs, Republic of Korea (Project No. 122037-02).
Acknowledgements	Not applicable.
Availability of data and material	Upon reasonable request, the datasets of this study can be available from the corresponding author.
Authors' contributions Please specify the authors' role using this form.	Conceptualization: Kim Y. Data curation: Park S, Park MA. Formal analysis: Park S. Methodology: Park S, Son S. Software: Park S. Validation: Park S, Kim DH, Son S. Investigation: Kim Y. Writing - original draft: Park S. Writing - review & editing: Park S, Son S, Park MA, Kim DH, Kim Y.
Ethics approval and consent to participate	This article does not require IRB/IACUC approval because there are no human and animal participants.

CORRESPONDING AUTHOR CONTACT INFORMATION

For the corresponding author (responsible for correspondence, proofreading, and reprints)	Fill in information in each box below
First name, middle initial, last name	Yangseon Kim
Email address – this is where your proofs will be sent	yangseon@cials.or.kr
Secondary Email address	nimitzdr93@gmail.com
Address	241, Cheomdangwahagro, Jeongeup-si, Jeonbuk, Korea
Cell phone number	82-10-5680-4340
Office phone number	82-63-536-6712
Fax number	82-63-536-6003

ACCEPTED

Abstract

Latilactobacillus curvatus CACC879 originated from swine feces in Korea, and its probiotic properties have been analyzed. The complete genome of strain CACC879 contained one chromosome 1,398,247 bp in length and three circular plasmids, namely, pCACC879-1 (591,981 bp), pCACC879-2 (14,542 bp), and pCACC879-3 (45,393 bp). The complete genome encodes a total of 2,077 genes, including 25 rRNA genes and 90 tRNA genes. In addition, probiotic stability-genes acid/bile related to salts tolerance, the biosynthesis of cobalamin (vitamin B12), riboflavin (vitamin B2), and CRISPR/Cas9 were found in the whole genomes. Remarkably, *L. curvatus* CACC879 contained the antioxidant-related (peroxiredoxin) and bacteriocin-related genes (*lysM* and *blpA*). Overall, these results demonstrate that *L. curvatus* CACC879 is a functional probiotic candidate for animal industry applications.

Keywords: *Latilactobacillus curvatus*, swine, probiotics, PacBio, genome sequence

Announcement

17

18

19 Lactic acid bacteria, such as *Latilactobacillus*, are useful microbes that produce healthy
20 metabolites, including bacteriocins and organic acids (such as lactic acid), that can regulate the gut
21 microbiome balance [1]. Lactic acid bacteria also confer health benefits via diverse mechanisms,
22 such as acid and bile tolerance, epithelial cell adherence, intestinal barrier buildup, and immune
23 system modulation [2]. *Latilactobacillus curvatus* is a potential probiotic strain that produces
24 various bacteriocins and metabolites and exhibits immunomodulatory activity [3, 4]. In this study,
25 the genome of *L. curvatus* CACC879 was sequenced and fully assembled to elucidate the genetic
26 factors associated with its probiotic characteristics.

27 *L. curvatus* CACC879 was isolated from swine feces in Korea, and the isolate was cultured
28 in De Man, Rogosa, and Sharpe (MRS) medium for 18 h at 37 °C. The genomic DNA of *L.*
29 *curvatus* CACC879 was extracted and purified using the DNeasy UltraClean kit (Qiagen, Hilden,
30 Germany) and sequenced using the PacBio Sequel (Pacific Biosciences, Menlo Park, CA, USA)
31 sequencing platform. *De novo* assembly was performed using PacBio SMRT analysis software
32 (version 2.3.0; Pacific Biosciences) [5]. The EggNOG 5.0 database (<http://eggnog5.embl.de>) was
33 used to classify all genes into clusters of ortholog gene (COG)/non-supervised orthologous group
34 (NOG) categories. Functional annotations of the predicted coding sequences (CDSs) were
35 compared with the Swiss-Prot and Kyoto Encyclopedia of Genes and Genomes (KEGG) [6]. The
36 genome sequence of CACC879 was compared with other reference strains by Orthologous average
37 nucleotide identity (OrthoANI; [http://www.ezbiocloud.net/tools /orthoani](http://www.ezbiocloud.net/tools/orthoani)) [7].

38 The whole genome of strain CACC879 consisted of one circular chromosome 1,398,247
39 bp in length (41.9% GC) along with three plasmids designated pCACC879-1 (591,981 bp, 42.2%
40 GC), pCACC879-2 (14,542 bp, 45.2% GC), and pCACC879-3 (45,393 bp, 41.2% GC) (Table 1
41 and Fig. 1A). The genome of strain CACC879 contains 2,077 CDSs and 115 non-coding genes

42 (25 rRNA and 90 tRNA genes) (Table 1). In addition, a total of 1,874 proteins (90.2%) were
43 matched and classified into 19 COG functional categories (Fig. 1B). The most abundant COG
44 categories were associated with replication, recombination, and repair (12.7%); translation,
45 ribosomal structure, and biogenesis (7.8%); transcription (7.6%), carbohydrate transport, and
46 metabolism (7.5%); and cell wall/membrane/envelope biogenesis (5.7%), excluding those with
47 unknown function (29.7%). Compared with the genome sequence of reference strains, the genome
48 of strain CACC879 was the most similar to that of the reference strains *L. curvatus* DSM 20019
49 (99.4%) and Wikim38 (99.0%) (Fig. 1C). The CACC879 strain showed common probiotic
50 properties including the CRISPR-associated endonuclease (Cas9) for antiviral-related mechanisms
51 and the biosynthesis of vitamin B groups (*ribF* and *pduO*), bacteriocin (*lysM*), and antioxidant
52 (*tpx*), compared to the reference strains [8-10]. Additionally, we confirmed that strain CACC879
53 harbors genes associated with common probiotic properties, including acid tolerance (*clpB* and
54 *grpE*), lactate synthesis (*ldh* and L-lactate dehydrogenase), and cell adhesion (*sotA*) (Table 2).
55 Interestingly, the CACC879 genome contained the *dltB* and *dltD* genes associated with the
56 modulation of the host immune response, but the reference strains did not. These findings will
57 serve as a reference for further studies on *L. curvatus* and provide a scientific basis for functional
58 probiotic development.

59

Nucleotide sequence accession number(s)

60

61 The whole-genome sequence of *L. curvatus* strain CACC879 (KACC 92511) has been deposited
62 in GenBank under accession numbers CP117683 (chromosome) and CP117684 to CP117686
63 (plasmids). The BioProject and BioSample accession numbers are PRJNA932593 and
64 SAMN33197937.

65

ACCEPTED

66

Acknowledgement

67 This research was supported by INNOPOLIS FOUNDATION through Science and Technology
68 Project Opens the Future of the Region, which is funded by Ministry of Science and ICT (2022-
69 DD-UP-0333), Korea, and partially supported by 2022 Technology commercialization support
70 project Ministry of Agriculture, Food and Rural Affairs, Republic of Korea (Project No. 122037-
71 02).

72

73

74

ACCEPTED

References

- 75
- 76
- 77 1. Hong SW, Kim JH, Bae HJ, Ham JS, Yoo JG, Chung KS, et al. Selection and characterization
78 of broad-spectrum antibacterial substance-producing *Lactobacillus curvatus* PA40 as a
79 potential probiotic for feed additives. *Anim Sci J.* 2018;89:1459–67. [https://doi.org/
80 10.1111/asj.13047](https://doi.org/10.1111/asj.13047)
- 81 2. Zommiti M, Connil N, Hamida JB, Ferchichi M. Probiotic characteristics of *Lactobacillus*
82 *curvatus* DN317, a strain isolated from chicken ceca. *Probiotics Antimicrob Proteins.*
83 2017;9:415–24. <https://doi.org/10.1007/s12602-017-9301-y>
- 84 3. Zommiti M, Almohammed H, Ferchichi M. Purification and Characterization of a Novel
85 Anti-Campylobacter Bacteriocin Produced by *Lactobacillus curvatus* DN317. *Probiotics*
86 *Antimicrob Proteins.* 2016;8:191–201. <https://doi.org/10.1007/s12602-016-9237-7>
- 87 4. Wang X, Li D, Meng Z, Kim K, Oh S. *Latilactobacillus curvatus* BYB3 isolated from kimchi
88 alleviates dextran sulfate sodium (DSS)-induced colitis in mice by inhibiting IL-6 and TNF-
89 R1 production. *J Microbiol Biotechnol.* 2022;32:348–54. [https://doi.org/10.4014/jmb.
90 2109.09054](https://doi.org/10.4014/jmb.2109.09054)
- 91 5. Park S, Kim JA, Jang HJ, Kim DH, Kim Y. Complete genome sequence of functional
92 probiotic candidate *Lactobacillus amylovorus* CACC736. *J Anim Sci Technol.* 2023;65:473–
93 7. <https://doi.org/10.5187/jast.2022.e85>
- 94 6. Kang S, Li R, Jin H, You HJ, Ji GE. Effects of selenium- and zinc-enriched *Lactobacillus*
95 *plantarum* SeZi on antioxidant capacities and gut microbiome in an ICR mouse model.
96 *Antioxidants (Basel).* 2020;9:1028. <https://doi.org/10.3390/antiox9101028>
- 97 7. Kim TY, Eun SH, Moon JS, Kwon HS, Choi S. Complete genome sequence of probiotic
98 *Lactobacillus helveticus* IDCC3801 isolated from infant feces. *Korean J Microbiol.*
99 2020;56:83–5
- 100 8. Rath D, Amlinger L, Rath A, Lundgren M. The CRISPR-Cas immune system: biology,
101 mechanisms and applications. *Biochimie.* 2015;117:119–28. [https://doi.org/10.1016/j.biochi.
102 2015.03.025](https://doi.org/10.1016/j.biochi.2015.03.025)
- 103 9. Capozzi V, Russo P, Dueñas MT, López P, Spano G. Lactic acid bacteria producing B-group
104 vitamins: a great potential for functional cereals products. *Appl Microbiol Biotechnol.*
105 2012;96:1383–94. <https://doi.org/10.1007/s00253-012-4440-2>

106 10. Averina OV, Poluektova EU, Marsova MV, Danilenko VN. Biomarkers and utility of the
107 antioxidant potential of probiotic Lactobacilli and bifidobacteria as representatives of the
108 human gut microbiota. *Biomedicines*. 2021;9:1340.
109 <https://doi.org/10.3390/biomedicines9101340>

110

ACCEPTED

111

Tables and Figures

112

113 **Table 1. Genome features of *Latilactobacillus curvatus* CACC879**


Properties	Chromosome		Plasmids		Total
	CACC879	pCACC879-1	pCACC879-2	pCACC879-3	
Length (bp)	1,398,247	591,981	14,542	45,393	2,050,163
GC content (%)	41.9	42.2	45.2	41.2	42.0
CDSs	1,421	596	11	49	2,077
tRNA	35	27	28	—	90
rRNA	14	8	3	—	25
CRISPR regions	1	—	—	—	1

114 GC, guanine and cytosine; CDSs, coding DNA sequences; tRNA, transfer RNA; rRNA, ribosomal RNA; CRISPR,
 115 clustered regularly interspaced short palindromic repeats

116 **Table 2. Predicted CDSs involved in *Latilactobacillus curvatus* CACC879 probiotic potency**

Predicted function	Gene	Start	End	Length (bp)
CRISPR-associated endonuclease	<i>Cas9</i>	31,830	35,723	3,894
Antimicrobial activity-related	<i>lysM</i>	1,089,047	1,090,990	1,944
Bacteriocin (Class II)-related	<i>blpA</i>	c557,488	c558,636	1,149
Lactate synthesis	<i>ldh</i>	c903,407	c904,384	978
Acid tolerance				
Chaperone protein ClpB	<i>clpB</i>	706,508	707,140	633
Chaperone protein GrpE	<i>grpE</i>	1,335,554	1,336,165	612
ClC family H(+)/Cl(-) exchange transporter	<i>eriC</i>	1,148,397	1,149,971	1,575
Sodium hydrogen exchanger family protein	<i>nhaP</i>	1,219,349	1,221,493	2145
F0F1 ATP synthase subunit A	<i>atpB</i>	17,075	17,788	714
F0F1 ATP synthase subunit B	<i>atpF</i>	18,084	18,605	522
F0F1 ATP synthase subunit C	<i>atpE</i>	17,807	18,019	213
F0F1 ATP synthase subunit delta	<i>atpH</i>	18,592	19,134	543
Bile salts tolerance	<i>cbh</i>	c246,195	C246,716	522
Cell adhesion	<i>sotA</i>	73,428	74,090	663
Stress response or protection				
Chaperone protein DnaK	<i>dnaK</i>	1,336,202	1,338,037	1,836
Chaperone protein DnaJ	<i>dnaJ</i>	1,338,161	1,339,309	1,149
Triose-phosphate isomerase	<i>tpiA</i>	c532,378	c533,133	756
Biosynthesis of vitamin B groups				
Riboflavin (B2)	<i>ribF</i>	1,332,094	1,333,047	954
	<i>ribT</i>	c271,721	c272,089	369
Cobalamin (B12)	<i>pduO</i>	58,751	59,305	555
Modulation of Immune response				
D-alanyl-lipoteichoic acid biosynthesis proteins	<i>dltB</i>	c220,566	c221,774	1,209
	<i>dltD</i>	c219,030	c220,298	1,269
Antioxidant (peroxiredoxin)	<i>tpx</i>	565,480	565,974	495
	<i>tpxA</i>	780,278	780,601	324

117 CDSs, coding sequences; CRISPR, clustered regularly interspaced short palindromic repeats



119

120 Figure 1. Genome features of *Lactobacillus curvatus* CACC879. (A) Circular genome mapping
 121 of *L. curvatus* CACC879. (B) Functional classification of clusters of orthologous groups (COG).
 122 (C) Orthologous average nucleotide identity (OrthoANI) values of *L. curvatus* CACC879
 123 compared to other reference strains.

124