











































374 Table 5. Effects of phytase supplementation on the nutritional characteristics of compost from broiler feces  
 375 (n=6)

	PC	NC	NC500	NC1000	NC1500	SEM	<i>p</i> value
N (%)	2.92	3.13	2.82	2.81	2.54	0.127	0.721
P <sub>2</sub> O <sub>5</sub> (%)	1.76 <sup>a</sup>	1.51 <sup>b</sup>	1.42 <sup>bc</sup>	1.35 <sup>bc</sup>	1.21 <sup>c</sup>	0.052	0.005
K <sub>2</sub> O (%)	1.38	1.45	1.47	1.31	1.32	0.047	0.781
MS (%)	59.2	57.5	60.0	59.0	59.3	1.317	0.986
pH	7.09	7.08	6.95	6.92	6.97	0.062	0.882

376 <sup>a,b,c</sup> means within the column with no common superscripts differ significantly (*p* < 0.05).  
 377 PC: positive control, NC: negative control, NC500: NC+phytase at 500 FTU/kg, NC1000: NC+phytase at  
 378 1,000 FTU/kg, NC1500: NC+phytase at 1,500 FTU/kg.  
 379 SEM: standard error of mean, N: nitrogen, P<sub>2</sub>O<sub>5</sub>: phosphorus pentoxide, K<sub>2</sub>O: potassium oxide, MS: moisture.

380

381

ACCEPTED