

Performance, hemato-biochemical indices and oxidative stress markers of broiler chicken fed phyto-genic during heat stress condition

Olatunji Abubakar Jimoh^{1*}, Olajumoke Temidayo Daramola¹, Hafsat Ololade Okin-Aminu² and Olayinka Abosede Ojo³

¹Department of Agricultural Technology, The Federal Polytechnic Ado-Ekiti, Ado Ekiti 360102, Ekiti State, Nigeria

²Animal Science Department, Ahmadu Bello University, Zaria 810211, Kaduna State, Nigeria

³Department of Animal Production, Fisheries and Aquaculture, Kwara State University, Molete 241103, Kwara State, Nigeria



Received: Jan 6, 2022

Revised: Mar 25, 2022

Accepted: Jun 1, 2022

*Corresponding author

Olatunji Abubakar Jimoh
Department of Agricultural Technology,
The Federal Polytechnic Ado-Ekiti, Ado
Ekiti 360102, Ekiti State, Nigeria.
Tel: +234-805-978-3712
E-mail: abubakarjimoh2011@gmail.com

Copyright © 2022 Korean Society of
Animal Sciences and Technology.
This is an Open Access article
distributed under the terms of the
Creative Commons Attribution
Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted
non-commercial use, distribution, and
reproduction in any medium, provided
the original work is properly cited.

ORCID

Olatunji Abubakar Jimoh
<https://orcid.org/0000-0001-8204-5816>
Olajumoke Temidayo Daramola
<https://orcid.org/0000-0002-0035-5245>
Hafsat Ololade Okin-Aminu
<https://orcid.org/0000-0001-7738-811X>
Olayinka Abosede Ojo
<https://orcid.org/0000-0001-5868-9793>

Competing interests

No potential conflict of interest relevant
to this article was reported.

Abstract

Thermal stress is a tremendous health predicament encountered by poultry farmers with adverse effects on the performance, product stature, health condition, survival, and overall welfare of poultry birds, and so requires urgent dietary user-friendly strategy to curb. This study was conducted with 200-day old broilers for the purpose of investigating the potential of phyto-genics in refining the negative effects of heat stress on broiler chicken. Moringa, Phyllanthus and mistletoe leaves were processed as phyto-genic supplements and incorporated into standard ration for broilers as treatments B1 (control), B2, B3 and B4 diet during the peak of thermal discomfort in humid tropics. Growth and carcass indices were monitored in a 49-day trial and blood samples were harvested at the end of the ordeal period to assess haematology, serum biochemical and oxidative stress markers with the use of standard procedures. The results obtained showed that the prevailing environmental condition in the study site indicated that the birds were exposed to heat stress. Birds fed on moringa and mistletoe supplements had higher performance index than birds without supplementation during heat stress condition, while birds fed on mistletoe supplement had the highest survival rate across the treatments. The liveweight, slaughter weight, dressed weight and eviscerated weight of heat stressed birds fed on moringa, phyllanthus and mistletoe supplements were significantly higher than birds on control treatment. Heterophyll/lymphocyte ratio of heat stressed birds without supplement were higher than birds on phyto-genic supplements, with least values recorded in phyllanthus and mistletoe fed birds. Birds on phyto-genic supplement tend to have lower cholesterol profile, lipid peroxidation and better antioxidant profile than birds on control treatment during heat stress conditions. Mistletoe supplementation in broiler ration enhances the survival rate, as well as promotes growth indices better among the phyto-genic supplements. However, phyto-genic supplements did ameliorate the negative effects of ther-

Funding sources

Not applicable.

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Jimoh OA.
 Data curation: Jimoh OA, Daramola OT.
 Formal analysis: Okin-Aminu HO, Ojo OA.
 Methodology: Jimoh OA, Ojo OA.
 Software: Okin-Aminu HO.
 Validation: Jimoh OA, Daramola OT.
 Investigation: Jimoh OA, Ojo OA.
 Writing - original draft: Daramola OT.
 Writing - review & editing: Jimoh OA, Daramola OT, Okin-Aminu HO, Ojo OA.

Ethics approval and consent to participate

Approval for the conduct of this research was obtained with Institutional Ethics Committee (Federal Polytechnic, Ado-Ekiti) with IACUC (approval no: FPA/EC/20/0405). Appropriate measures were taken to minimise pain or discomfort to the animals in line with the National Institute of Health guide for the care and use of laboratory animals.

mal discomfort on performance, physiological and oxidative stress in heat-stressed broiler chicken.

Keywords: Antioxidant, Chicken, Leukocyte, Survival rate, Thermal discomfort

INTRODUCTION

The shock of heat stress on poultry birds is a significant predicament and a cause for economic dispossession in the poultry industry, hence it exerts a detrimental influence on physiological responses like immunity, oxidative stress, and intestinal and muscular functions [1]. Poultry birds enjoys to be raised under situations of minor energy requirement for thermoregulation purposes and ultimate net energy available for production goal, otherwise poultry will be under stress [2]. Reports of Abreu and Abreu [3] and Bueno et al. [4] stated that the actual temperature varies from 32°C to 35°C (1st week), 23°C–32°C (2nd and 5th week), and 20°C (6th week) and relative humidity ranging from 60%–70%, which are difficult conditions to keep in the tropics. However, heat stress is one of the most deleterious contentions in tropical and subtropical corridors of the world, which causes substantive economic forfeitures in poultry production [5]. Thermal-stressed birds elevated heterophil/lymphocyte ratios, liver enzymes and correlates with higher mortality levels [6]. The exposure of poultry to a range of stressors account for depressed productive, reproductive performance and exposed health [7,8]. This in turn necessitates the requirement for complementary reinforcement for the antioxidant system in the poultry industry [9].

Emerging reports suggest that enhancement with natural bioactive compounds potently palliate the heat stress in chicken due to their pooled biological activities [1]. Jang et al. [10] reported that the tool by which oxidative stress induced by heat stress is mutually connected with the initiation of cell-mediated immunity via antioxidant defense system. Thus, antioxidant enrichment to mitigate oxidative stress will influence humoral responses.

Phytochemicals with antioxidant activity such as polyphenols, which are vital secondary metabolite present in plants, are slated to meditate heat stress in poultry birds [11]. Complementing the poultry diet with a capable phytochemical-containing feed could definitely ameliorate the shock of heat stress in poultry production [11]. The potentials of medicinal plants such as *Phyllanthus amarus*, *Moringa oleifera* and *Viscum album* to enhance animal welfare has been investigated in animal [12]. Herbs possess bioactive constituent which confers optimal antioxidant defense to mitigate adverse effects of heat stress [13]. *Moringa oleifera* leaves have been reported to enhance the immune responses and improve the intestinal health of broilers [14,15]. Use of antioxidant products derived from plants is favoured over chemical antioxidants due to food safety reasons [16]. This study was conducted to investigate the effect of three phytochemicals on stress markers and performance of broiler chicken raised under heat stress conditions.

MATERIALS AND METHODS

Experimental site

This study was carried out at the Teaching and Research Farm at the Federal Polytechnic, Ado-Ekiti, Ekiti State in Southwest Nigeria, from February to March, 2020; which is the peak of heat stress in southern Nigeria. Severity of heat stress based on the combination of temperature and humidity (Temperature - Humidity Index) was established to peak in southern Nigeria between February and March [17–19]. Temperature and relative humidity (RH) of the poultry microclimate was recorded at 08:00 h, 12:00h and 18.00 h daily during the period covered by the study using a Thermo-Hygrometer. Daily records of the ambient temperature and RH were used to compute the

temperature humidity index (THI) as outlined in Jimoh et al. [20].

Phytonics processing and evaluation

Fresh Moringa (*Moringa oleifera*), Phyllanthus (*Phyllanthus amarus*) and Mistletoe (*Viscum album*) leaves were harvested from an established orchard within Ado-Ekiti metropolis of Ekiti State. The plants were identified and indexed with herbarium voucher numbers; *Viscum album* UILH/002/084/1210/2021; *Moringa oleifera* UILH/001/1008/2021; *Phyllanthus amarus* UILH/003/1109/2021.

Leaves were detached from twigs and shade dried until it was crispy to touch while retaining their greenish coloration. The leaves were milled and stored in air-tight containers until incorporation into the diet.

Experimental animals and management

Two hundred (200) day-old arbor acres broilers were purchased from a reputable hatchery. They were weighed and allotted to four treatment diets (B1, B2, B3 and B4) of 10 replicates and five birds per replicate in a completely randomized design.

Four experimental diets were formulated to meet the nutrient requirement for broilers at starter and finisher phases as shown in Table 1 and 2, respectively; diet 1 without leaf meal (B1), diet 2 with 5% *Moringa oleifera* (B2), diet 3 with 5% *Phyllanthus amarus* (B3) and diet 4 with 5% *Viscum album* (B4). Proximate composition of the experimental diets was carried out using AOAC standard procedures and presented in Table 3. Similarly, phytochemical screening and proximate

Table 1. Gross composition of broiler starter ration

Ingredient	Diet 1	Diet 2	Diet 3	Diet 4
Maize	58.5	58.5	58.5	58.5
Groundnut cake	9.8	8.8	8.8	8.8
Soyabean meal	25	21	21	21
Fishmeal	3.1	3.1	3.1	3.1
Methionine	0.25	0.25	0.25	0.25
Lysine	0.25	0.25	0.25	0.25
Oyster shell	1.5	1.5	1.5	1.5
Limestone	1.1	1.1	1.1	1.1
Premix	0.25	0.25	0.25	0.25
Salt	0.25	0.25	0.25	0.25
Moringa	-	5	-	-
Phyllanthus	-	-	5	-
Mistletoe	-	-	-	5
Total	100	100	100	100
Dry matter (%)	85.848	85.878	85.878	85.878
Crude protein (%)	22.992	21.407	21.407	21.407
Metabolisable energy (kcal/kg)	3031.27	3026.87	3026.87	3026.87
Ether extract (%)	3.9425	3.808	3.808	3.808
Crude fibre (%)	3.316	4.596	4.596	4.596
Lysine (%)	1.37755	1.28155	1.28155	1.28155
Methionine (%)	0.60564	0.59674	0.59674	0.59674
Calcium (%)	1.17455	1.31155	1.31155	1.31155
Phosphorus (%)	0.31525	0.30475	0.30475	0.30475

Table 2. Gross composition of broiler finisher ration

Ingredient	Diet 1	Diet 2	Diet 3	Diet 4
Maize	66.5	66.5	66.5	66.5
Groundnut cake	8.3	6.3	6.3	6.3
Soyabean meal	19.5	16.5	16.5	16.5
Fishmeal	2	2	2	2
Methionine	0.25	0.25	0.25	0.25
Lysine	0.25	0.25	0.25	0.25
Oyster shell	1.5	1.5	1.5	1.5
Limestone	1.2	1.2	1.2	1.2
Premix	0.25	0.25	0.25	0.25
Salt	0.25	0.25	0.25	0.25
Moringa	-	5	-	-
Phylanthus	-	-	5	-
Mistletoe	-	-	-	5
Total	100	100	100	100
Dry matter (%)	85.546	85.781	85.781	85.781
Crude protein (%)	18.575	19.1625	19.1625	19.1625
Metabolisable energy (kcal/kg)	3,086.43	3,088.948	3,088.948	3,088.948
Ether extract (%)	3.9305	3.81995	3.81995	3.81995
Crude fibre (%)	3.0325	4.33995	4.33995	4.33995
Lysine (%)	1.17005	2.27055	2.27055	2.27055
Methionine (%)	0.56059	0.55654	0.55654	0.55654
Calcium (%)	1.15925	0.748475	0.748475	0.748475
Phosphorus (%)	0.47845	0.252875	0.252875	0.252875

Table 3. Proximate composition of broiler experimental diets

Variable	Diet 1	Diet 2	Diet 3	Diet 4
Broiler starter ration				
Dry matter (%)	91.39	92.16	91.45	91.60
Crude protein (%)	21.00	22.85	20.95	21.51
Ether extract (%)	5.50	5.50	5.20	5.70
Crude fibre (%)	4.50	4.90	4.80	4.80
Ash	6.40	6.60	6.80	6.90
Nitrogen free extract	62.6	63.15	66.25	65.09
Broiler finisher ration				
Dry matter (%)	91.80	91.91	92.23	92.10
Crude protein (%)	20.56	20.30	20.66	20.70
Ether extract (%)	5.90	6.20	6.10	5.80
Crude fibre (%)	5.40	5.30	4.80	5.20
Ash	5.60	6.10	6.20	6.20
Nitrogen free extract	62.54	62.1	62.24	62.1

composition of the leaf meals were carried out using AOAC standard procedures and presented in Table 4.

Birds were fed *ad libitum* and fresh water was offered daily. The vaccination programme as

Table 4. Proximate and Phytochemical analysis of the leafmeals

Variable	Mistletoe leaf meal	Moringa leaf meal	Phyllanthus leaf meal
Dry matter (%)	89.82	90.76	90.50
Crude protein (%)	18.81	31.06	27.13
Ether extract (%)	2.10	2.7	2.5
Crude fibre (%)	12.10	11.30	9.90
Ash (%)	14.90	12.40	12.06
Nitrogen free extract (%)	52.09	42.54	48.41
Alkaloids (mg/100 g)	14.68	8.5	10.34
Saponins (mg/100 g)	31.01	39.81	23.96
Glycosides (mg/100 g)	75.08	95.92	81.66
Steroids (mg/100 g)	18.82	25.00	20.54
Flavonoids (mg/100 g)	62.93	25.03	44.86
Tannins (mg/100 g)	114.81	96.53	95.98

recommended by the hatchery were followed and no medication was offered throughout the study. The feed intake and weight changes were monitored throughout the study to evaluate their growth in a 49-day trial. Mortality was recorded and feed conversion were computed per replicate.

Record of feed intake, weight gain and mortality were taken weekly. Feed conversion ratio (FCR) was obtained by calculation.

$$\text{Total Feed Intake (g)} = \text{Total feed supplied (g)} - \text{Total feed left over (g)}$$

$$\text{Average feed intake (g/bird)} = \text{Total feed intake} / \text{Number of birds}$$

$$\text{Total weight gain} = \text{Final weight} - \text{Initial weight}$$

$$\text{FCR} = \text{Total feed intake (g)} / \text{Total weight gain (g)}$$

$$\% \text{Mortality} = \frac{\text{Number of dead birds}}{\text{Total number of stocked birds}} \times 100$$

Blood sample collection

At the end of the feeding trial, blood samples were collected from 3 birds per replicate into plain sample and heparinized bottles for serum and haematology respectively. For serum, blood samples in plain tubes were centrifuged and serum obtained using standard procedures and stored at -20°C until analysis. The samples in heparinized tubes were analysed for packed cell volume, hemoglobin, red blood cells, white blood cells and its differential counts using standard procedures. Serum biochemical assay; glucose, total protein, albumin, globulin, creatinine, urea, alanine amino transferase (AST), Aspartate amino transferase (AST), cholesterol, triglyceride, high density lipoprotein (HDL), low density lipoprotein (LDL) was carried out using fortress diagnostics commercial assay kits and its procedures.

Determination of serum total antioxidant capacity activities, superoxide dismutase (SOD), glutathione peroxidase, catalase activities and lipid peroxidation were assay as outlined in Jimoh et al. [21].

Data analysis

All data were subjected to one way analysis of variance and significant means were separated using Duncan multiple range test IBM SPSS.

RESULT

The proximate composition of broiler starter and finisher rations are presented in Table 3. The proximate compositions of the diets show similarity in the nutrient profile of the diets. Proximate and phytochemical analysis of the leaf meals is shown in Table 4. Moringa possess higher crude protein, saponins, glycosides, steroids among the three leaf meals. Mistletoe possesses higher crude fibre, ash, nitrogen free extract, alkaloids, flavonoids and tannins among the three leaf meals. Phyllanthus possess least crude fibre, ash, saponin and tannin among the three leaf meals.

The THI of the broiler pen is shown in Fig. 1. The range of values obtained indicate that the birds were exposed to an average THI of 24.13 (absence of heat stress) in the 8:00 hrs. At 12:00 hrs, the birds were exposed to an average THI of 30.35 (very severe of heat stress). At 18:00 hrs, the birds were exposed to an average THI of 31.95 (very severe heat stress).

The performance characteristics of broilers fed on phytogetic supplement during heat stress condition is shown in Table 5. At the starter phase, final body weight of birds fed on treatment B2 and B4 based diets were significantly ($p < 0.05$) higher than birds on treatment B3, while birds fed treatment B1 had significantly ($p < 0.05$) least. Weight gain of birds on phytogetic supplements were statistically similar, but birds on treatments B2 and B4 had significantly ($p < 0.05$) higher weight gain than birds on B1. The feed intake of birds fed on phytogetic supplements were statistically ($p < 0.05$) lower than birds on control. FCR of birds on treatments B1, B3 and B4 were significantly ($p < 0.05$) higher than those of birds on B2

In the finisher phase, final body weight and weight gain of birds fed on treatments B2 and B4 were significantly ($p < 0.05$) higher than those on treatment B1. The feed intake of birds on B3 were significantly ($p < 0.05$) lower than birds on treatment B1, while birds on treatments B2 and B4 were statistically similar to all the treatments. The FCR of birds on treatments B2, B3 and B4 were statistically ($p > 0.05$) similar, but significantly ($p < 0.05$) lower than birds on treatment B1. The percentage mortality of heat stressed birds on treatments B2 and B3 were significantly ($p < 0.05$) lower than birds on treatment B1, with the statistically ($p < 0.05$) least value obtained in birds on B4.

The overall performance of the birds showed that birds fed on B2 and B4 had statistically ($p > 0.05$) similar body weight, weight gain and feed intake and were significantly ($p < 0.05$) higher than those of birds on treatment B1. While birds fed on treatment B3 had significantly ($p > 0.05$) similar




Fig. 1. Temperature humidity index of broiler pen.

Table 5. Performance characteristics of broiler chicken fed phytochemical feed supplement in a hot dry climate

Variable	B1	B2	B3	B4	SEM	p-value
Starter phase (0–21 d)						
Initial body weight (g/bird)	61.20	64.00	63.60	63.05	1.06	0.00
Body weight at 21 d (g/bird)	894.07 ^c	963.89 ^a	909.28 ^b	941.23 ^a	12.41	0.00
Average weight gain (g/bird)	832.86 ^b	899.89 ^a	845.67 ^{ab}	878.18 ^a	12.09	0.00
Average feed intake (g/bird/day)	97.21 ^c	101.49 ^b	101.87 ^{ab}	104.33 ^a	0.68	0.02
Feed conversion ratio	2.53 ^a	2.37 ^b	2.54 ^a	2.50 ^a	0.04	0.00
Mortality	0.00	0.00	0.00	0.00	0.00	0.00
Finisher phase (21–49 d)						
Body weight at 21 d (g/bird)	894.07 ^c	963.89 ^a	909.28 ^b	941.23 ^a	12.41	0.01
Body weight at 49 d (g/bird)	2,428.94 ^b	2,689.03 ^a	2,528.01 ^{ab}	2,650.72 ^a	34.53	0.00
Average weight gain (g/bird)	1,534.87 ^b	1,789.14 ^a	1,682.34 ^{ab}	1,772.55 ^a	36.06	0.01
Average feed intake (g/bird/day)	143.15 ^a	132.71 ^{ab}	123.54 ^b	138.53 ^{ab}	3.15	0.00
Feed conversion ratio	2.63 ^a	2.08 ^b	2.06 ^b	2.19 ^b	0.07	0.03
Mortality	11.67 ^a	7.50 ^b	7.50 ^b	2.78 ^c	2.20	0.02
Overall (49 d)						
Initial body weight (g/bird)	61.20	64.00	63.60	63.05	1.06	0.00
Body weight at 49 d (g/bird)	2,428.94 ^b	2,689.03 ^a	2,528.01 ^{ab}	2,650.72 ^a	34.53	0.02
Average weight gain (g/bird)	2,367.74 ^b	2,625.03 ^a	2,464.41 ^{ab}	2,587.68 ^a	34.09	0.03
Average feed intake (g/bird/day)	113.48 ^b	123.46 ^a	119.23 ^a	123.87 ^a	1.81	0.00
Feed conversion ratio	2.58 ^a	2.23 ^b	2.27 ^b	2.34 ^b	0.03	0.01
Mortality	11.67 ^a	7.50 ^b	7.50 ^b	2.78 ^c	2.20	0.02

B1, diet without leaf meal; B2, diet with *Moringa oleifera*; B3, diet with *Phyllanthus amarus*; B4, diet with *Viscum album*.

^{a-c}Means with different superscripts in the same row significantly differ at $p < 0.05$.

body weight, weight gain and feed intake as birds on treatment B2 and B4. The FCR of the birds on phytochemical supplements were significantly ($p < 0.05$) lower than the control. The percentage mortality of heat stressed birds on treatments B2 and B3 were significantly ($p < 0.05$) lower than birds on treatment B1, with the statistically ($p < 0.05$) least value obtained in birds on treatment B4.

Carcass traits of broilers fed on phytochemical feed supplement is shown in Table 6. All carcass indices assessed were not significantly ($p > 0.05$) affected by the treatments except live weight, slaughter weight, dressed weight, eviscerated weight, pancreas, head and gizzard weight. The live weight, slaughter weight, dressed weight, eviscerated weight of heat stressed birds fed on treatments B2, B3 and B4 were significantly higher than birds fed on B1. The pancreas weight of birds fed on treatments B2 and B4 were statistically ($p > 0.05$) similar to those of other treatments, but pancreas weight of birds fed on treatment B1 was significantly ($p < 0.05$) higher than that of birds on B3. Head of birds fed on treatments B2, B3 and B4 were statistically ($p > 0.05$) similar, but significantly ($p < 0.05$) higher than those of birds on treatment B1.

The haematological indices of broilers fed on phytochemical supplements is shown in Table 7. The packed cell volume, erythrocyte, hemoglobin, mean cell hemoglobin concentration (MCHC), mean cell volume (MCV) and mean cell hemoglobin (MCH) of birds were not significantly ($p > 0.05$) affected by the phytochemical supplement. Leukocytes of birds fed on treatment B2 was significantly ($p < 0.05$) highest, and the significantly ($p < 0.05$) least value was obtained in B3. Birds fed on treatment B1 had significantly ($p < 0.05$) higher leukocyte than birds fed on B4. The heterophils of birds fed on B1 was significantly ($p < 0.05$) higher than birds fed on treatment B2, while birds fed on treatments B3 and B4 had statistically least values. Lymphocytes of birds fed on B2 were

Table 6. Carcass traits of broilers fed phytogetic feed supplement in a hot dry climate

Variable	B1	B2	B3	B4	SEM	p-value
Live weight (g)	2,478.68 ^b	2,692.98 ^a	2,518.56 ^b	2,631.68 ^a	54.79	0.02
Slaughter weight (g)	2,412.08 ^b	2,600.58 ^a	2,496.08 ^b	2,595.82 ^a	51.85	0.01
Eviscerated weight (g)	2,378.46 ^b	2,533.26 ^a	2,411.38 ^{ab}	2,477.30 ^a	52.00	0.03
Dressed weight (g)	1,860.26 ^b	2,197.12 ^a	1,965.72 ^a	2,095.24 ^a	42.18	0.00
Heart (g)	10.34	8.86	9.42	11.20	0.51	0.08
Lungs (g)	13.32	14.58	13.80	12.16	0.72	0.11
Shank (g)	94.92	83.98	95.22	80.42	4.71	0.15
Kidney (g)	13.82	13.62	11.22	13.64	0.69	0.20
Liver (g)	49.94	54.10	45.12	43.78	2.21	0.17
Pancreas (g)	5.40 ^a	4.20 ^{ab}	3.92 ^b	4.76 ^{ab}	0.23	0.00
Head (g)	63.06 ^b	70.24 ^a	74.72 ^a	79.44 ^a	2.18	0.00
Spleen (g)	2.40	1.86	2.58	2.00	0.19	0.32
Gizzard (g)	49.70 ^a	40.80 ^b	43.68 ^b	50.18 ^a	1.88	0.02
Proventriculus (g)	9.60	7.14	8.24	8.64	0.49	0.22
Gall Bladder (g)	1.64	2.08	1.76	2.42	0.16	0.42
Intestine (g)	131.20	122.56	113.80	123.24	3.51	0.64

B1, diet without leaf meal; B2, diet with *Moringa oleifera*; B3, diet with *Phyllanthus amarus*; B4, diet with *Viscum album*.

^{a,b}Means with different superscripts in the same row significantly differ at $p < 0.05$.

Table 7. Haematology of broilers fed phytogetic feed supplement in a hot dry climate

Variable	B1	B2	B3	B4	SEM	p-value
Packed cell volume (%)	26.67	26.67	25.33	23.67	0.66	0.21
Erythrocyte ($\times 10^6/L$)	2.87	2.00	2.77	2.40	0.21	0.17
Haemoglobin (Hbg/dL)	8.89	8.89	8.44	7.89	0.22	0.08
Mean cell hemoglobin concentration (g/dL)	33.13	32.81	33.13	33.10	0.09	0.23
Mean cell volume (fL)	101.28	139.11	95.89	101.20	9.25	0.26
Mean cell hemoglobin (pg/cell)	33.76	46.37	31.96	33.73	3.08	0.16
Leukocytes ($\times 10^9/L$)	4.27 ^b	5.20 ^a	2.80 ^d	3.80 ^c	0.49	0.02
Heterophils ($\times 10^9/L$)	2.28 ^a	1.35 ^b	0.50 ^c	0.66 ^c	0.30	0.01
Lymphocytes ($\times 10^9/L$)	1.93 ^b	2.71 ^a	2.26 ^{ab}	2.14 ^{ab}	0.27	0.04
Monocytes ($\times 10^9/L$)	0.05 ^b	0.12 ^a	0.02 ^b	0.01 ^b	0.02	0.03
Heterophil/Lymphocyte ratio	1.18 ^a	0.85 ^b	0.22 ^c	0.31 ^c	0.16	0.00

B1, diet without leaf meal; B2, diet with *Moringa oleifera*; B3, diet with *Phyllanthus amarus*; B4, diet with *Viscum album*.

^{a-d}Means with different superscripts in the same row significantly differ at $p < 0.05$.

significantly ($p < 0.05$) higher than that of birds fed on treatment B1. Monocytes of birds fed on treatment B2 was significantly ($p < 0.05$) higher than birds fed on B1, B3 and B4. Heterophils/lymphocyte ratio of birds fed on treatments B2 B3, and B4 were statistically ($p > 0.05$) similar, while birds fed on treatment B1 had significantly ($p < 0.05$) higher values than those of birds fed on treatments B3 and B4.

The serum biochemistry of birds fed on phytogetic feed supplements is shown in Table 8. Blood glucose and albumin of birds fed on treatment B3 were significantly ($p < 0.05$) higher than birds fed on treatments B1, B2 and B4. Blood total protein of birds fed on treatment B2 were significantly ($p < 0.05$) higher than birds fed treatments B3 and B4 and the significantly ($p < 0.05$) least blood total protein was obtained in birds fed on treatment B1. Blood globulin of birds fed on

Table 8. Serum Biochemistry of broilers fed phytogetic feed supplement in a hot dry climate

Variable	B1	B2	B3	B4	SEM	p-value
Glucose (mmol/L)	4.53 ^b	4.45 ^b	7.06 ^a	4.87 ^b	0.47	0.03
Total protein (g/L)	43.59 ^c	65.16 ^a	54.61 ^b	57.41 ^b	5.35	0.00
Albumin (g/L)	16.93 ^b	19.10 ^b	22.11 ^a	18.94 ^b	1.93	0.04
Globulin (g/L)	27.44 ^b	46.06 ^a	41.44 ^a	38.48 ^a	5.26	0.02
Triglyceride (mmol/L)	2.35 ^a	1.79 ^b	1.92 ^b	2.30 ^a	0.21	0.00
Cholesterol (mmol/L)	2.89 ^a	1.72 ^c	2.73 ^b	2.46 ^b	0.21	0.00
High density lipoprotein (mmol/L)	0.29 ^b	0.24 ^b	0.41 ^a	0.34 ^a	0.04	0.00
Low density lipoprotein (mmol/L)	1.67 ^a	1.09 ^b	1.44 ^b	1.14 ^b	0.20	0.02
Creatinine (mg/dL)	4.67 ^a	1.58 ^b	3.75 ^a	2.00 ^b	0.90	0.01
Aspartate amino transferase (U/l)	29.96	31.60	26.67	26.28	2.69	0.10
Alanine amino transferase (U/l)	3.77	3.10	4.01	3.36	0.54	0.08
Uric acid (mg/dL)	5.83 ^b	5.38 ^b	7.21 ^a	8.40 ^a	0.65	0.04

B1, diet without leaf meal; B2, diet with *Moringa oleifera*; B3, diet with *Phyllanthus amarus*; B4, diet with *Viscum album*.

^{a-c}Means with different superscripts in the same row significantly differ at $p < 0.05$.

phytogetic supplemented diets (B2, B3 and B4) were significantly ($p < 0.05$) higher than those fed on treatment B1. Blood triglyceride of birds fed on treatments B1 and B4 were significantly ($p < 0.05$) higher than that of birds fed on treatments B2 and B3. Blood cholesterol of birds fed on treatment B1 were significantly ($p < 0.05$) higher than birds fed on treatments B3 and B4, which share statistically ($p > 0.05$) similar values, while the significantly ($p < 0.05$) least cholesterol were obtained in birds fed on treatment B2. Blood HDL of birds fed on treatments B3 and B4 were significantly ($p < 0.05$) higher than those of birds fed on treatments B1 and B2. LDL of birds fed phytogetic supplemented diets (B2, B3 and B4) were significantly ($p < 0.05$) lower than those fed on B1. Blood creatinine of birds fed on treatments B1 and B3 were significantly ($p < 0.05$) higher than birds fed on treatments B2 and B4. AST and ALT of birds were not significantly ($p > 0.05$) influenced by the treatments. Uric acid of birds fed on treatments B3 and B4 were statistically ($p > 0.05$) similar, however, significantly ($p < 0.05$) higher than birds fed on treatments B1 and B2 which had statistically ($p > 0.05$) similar values.

Serum oxidative stress markers of broilers fed on phytogetic feed supplements is shown in Table 9. Serum total antioxidant activity of birds fed on phytogetic supplemented groups were significantly ($p < 0.05$) higher than birds on treatment B1. Serum lipid peroxidation of birds on treatments B2, B3 and B4 were statistically ($p > 0.05$) similar and significantly ($p < 0.05$) lower than birds on treatment B1. Blood SOD of birds fed on treatments B1, B3 and B4 were statistically ($p > 0.05$) similar and significantly ($p < 0.05$) lower than birds fed on treatment B2. Blood GPx of birds

Table 9. Oxidative stress markers of broilers fed phytogetic supplement in a hot dry climate

Variable	B1	B2	B3	B4	SEM	p-value
Total antioxidant activity (mmol/L)	0.62 ^b	2.23 ^a	2.85 ^a	2.98 ^a	0.26	0.01
Lipid peroxidation ($\times 10^{-5}$ MDA/mg protein)	3.26 ^a	1.66 ^b	0.75 ^b	0.95 ^b	0.59	0.02
Superoxide dismutase (U/min/mg protein)	1.78 ^b	10.61 ^a	1.33 ^b	3.06 ^b	2.05	0.02
Glutathione peroxidase (mg GSH/min/mg protein)	19.74 ^b	20.34 ^{ab}	23.68 ^a	24.15 ^a	3.93	0.00
Catalase (nm H ₂ O ₂ /min/mg protein)	0.36 ^b	0.41 ^{ab}	0.57 ^a	0.71 ^a	0.16	0.00

B1, diet without leaf meal; B2, diet with *Moringa oleifera*; B3, diet with *Phyllanthus amarus*; B4, diet with *Viscum album*.

^{a,b}Means with different superscripts in the same row significantly differ at $p < 0.05$.

fed on treatments B3 and B4 were significantly ($p < 0.05$) higher than that of birds fed on treatment B1. Blood catalase activity of birds fed on treatments B3 and B4 were significantly ($p < 0.05$) higher than those of birds fed on treatment B1.

DISCUSSION

The rich phytochemical constituents of the phytogetic supplements suggest the enrichment they confer on the diets fed to the birds. The phytogetic supplements possess flavonoids (kaempferol, quercetin and rutin), glycosides, steroids and phenolic acids (gallic, chlorogenic, ellagic and ferulic acid) [22], glucosinolates and are valuable source of β -carotene (precursor of vit. A) and vitamins (B-complex, C, D and K) [23], which confers high antioxidant activities.

The trend of results shows that the phyto-genics supplements enhanced feed intake, and body weight gain and resulted in better feed conversion efficiency of birds during heat stress condition. This suggests that antimicrobial combination in the phyto-genics moderate the microflora in the gastrointestinal tract, which accordingly enhances the digestibility of feed and growth performance in broiler chickens [24]. The inclusion of the phyto-genic supplements also reduced mortality associated with thermal discomfort, especially during the finisher phase. Birds fed on mistletoe supplement during heat stress condition had the highest survival rate across the treatment. Birds fed on moringa and mistletoe supplement had higher performance index than birds without supplementation during heat stress condition. This is in concurrence with Selim et al. [25] that moringa had a favourable impact on the rabbit performance as it plays a key role in boosting nutrient utilization, owing to its bioactive compounds. Similarly, Saleh et al. [24] suggested that medical plants enhance appetite and digestive enzymes or have coccidiostat and improved digestibility, and accounts for promoting performance and digestibility due to the antimicrobial and antifungal effects.

Mortality recorded during finisher phase would be attributed to higher heat load due to metabolic size compared to starter phase, and lesser heat dissipation mechanisms associated with fast growing birds. Similar claims of supplementing *Moringa oleifera* leaves affirms that it ameliorates the negative effect of thermal discomfort on productive performances and physiological variables of heat stressed broiler in tropical regions [26]. Incongruent with our result is the claim of higher FCR on herbs fed/treated animals. Attributed to flavonoids, which impede the *in vitro* digestibility of proteins and starch in bread through the accumulation of indigestible complexes with the nutrients, and by impeding the activity of several digestive enzymes [27]. The supplements fed to these birds during heat stress condition, enhances appetite, weight gain and survival rate of the birds due to the myriad of phyto-constituents in the supplements.

Reports that dietary moringa essential oil improves growth performance in stressed poultry [2,28,29] supports the claims of this study. Similarly, Daramola et al. [30] has reported that moringa leaf meal can be used as feed additive to enhance performance and antioxidant status of laying birds.

The trend of result shows that carcass merit of heat stressed birds fed on phyto-genic supplement were better than birds without supplement. However, the organ weights were not affected by the supplements except the gizzard. Similar claims of Hassan et al. [31] that heat stressed broiler chicks fed on *Moringa oleifera* improves body weight and FCR, with no effect on carcass indices.

The phyto-genic supplements had no effect on erythrocytic indices of heat stressed broilers. White blood cell and its differential counts were influenced by heat stress and phyto-genic supplement in broiler. Birds fed on moringa supplement and non-supplement group had higher leukocyte and its differential counts than birds fed phyllanthus and mistletoe supplements. Heterophyl/lymphocyte ratio of heat stressed birds without supplement were higher than birds on phyto-genic supplements,

with least values recorded in phyllanthus and mistletoe fed birds. This study supports Jang et al. [10] that antioxidant activated immune function via the accrual of lymphocytes and macrophages in heat stressed chickens [32]. Similarly, Jang et al. [10] reported that heat stressed birds fed on diet supplemented with vitamin E have improved immunity under stressed conditions.

Birds fed on phyllanthus supplements during heat stress condition tends to have higher glucose values. Adequate glucose concentration is crucial for energy accessibility, liver glycogen synthesis, and normal gene expression of liver phosphoenolpyruvatecarboxy-kinase, which might bestow to energy production by animals and might provide energy for cellular metabolism without additional heat production [33]. Heat stress reduces protein deposition, elevates protein catabolism (pronounced by an elevated plasma uric acid level), and reduces protein synthesis and N retention [34]. Total protein of birds on phytochemical supplements were better than those on control, largely due to the higher serum globulin in birds fed on phytochemical supplement. This could indicate higher immunoglobulin activity for cellular immunological response to combat the adverse effects of heat stress in the birds. Similarly, the addition of moringa leaves in broiler diet increase plasma total proteins associated with protein synthesis which mirrored positively on the productive performance of broilers (increased live body weight and body weight gain) [31]. This could be linked with better performance associated with phytochemical supplemented birds during heat stress conditions in this study. Similar to higher albumin in birds fed on phyllanthus supplements is the report that moringa improve serum albumin concentration [35] due to its responsibility for 75% to 80% of the vascular colloidal osmotic pressure and a vital agent in maintaining equilibrium with tissue fluids. Incongruence with Habibi et al. [36] who reported non-beneficial effect of diet supplementation on total protein level. Similarly, total protein and albumin concentrations increased with dietary vitamin C and folic acid supplementation to heat stressed Japanese quails' concentration [37].

The trend of result shows that birds on phytochemical supplement tend to have lower cholesterol profile, with higher HDL and lower LDL than birds on control. It could be due to pancreatic cholesterol esterase hydrolyses dietary cholesterol esters which releases free cholesterol in the lumen of the small intestine, the constraint of cholesterol esterase would limit the absorption of dietary cholesterol and thereby reduce cholesterol concentration [37].

Similarly, Tayer et al. [38] related that the enhance effects of flavonoids in medicinal plants relates to the hypoglycemic and hepatic glucokinase activity of the liver. Ferit Gursu et al. [39] reported that decline in corticosterone via inhibitory effect of vitamin C on glucocorticoid synthesis a response to heat stress, led to decrease in protein derived gluconeogenesis led to decreased serum cholesterol, triglyceride, HDL cholesterol, and glucose concentrations. Similar to the trend of results obtained in this study, is the report that *Moringa oleifera* reduce lipids and lipid peroxidation levels in rats [40]. There is no indication of organ toxicity due to similar ALT and AST across the treatments, creatinine was lower in birds fed on moringa and mistletoe supplements, uric acid was higher in birds fed on phyllanthus and mistletoe. Contrariwise, levels of aspartate amino transferase, ALT and alkaline phosphatase were reduced by *Moringa oleifera* leaves in rats [41]. Also, decreased Heterophil/Lymphocyte ratio, increased Plasma total protein, globulin and thyroid hormones were the effects of *Moringa oleifera* on heat stressed broiler chicks [23].

Environmental heat stress associated with increased lipid peroxidation by raising both of thiobarbituric acid reactive substance (TBARS) and malondialdehyde (MDA) levels in broilers [42] were mitigated by the phytochemical supplements administered to the birds during heat stress conditions. Birds fed on phytochemical supplements had better oxidative stability due to superior antioxidant activity and reduced lipid peroxidation compared to birds without supplement during heat stress condition. The antioxidants potency of the phytochemicals can be linked to this result as claimed by Jang et al. [10] that antioxidant vitamins employed a beneficial influence on sustaining

immunity and antioxidant status within summer conditions. The addition of *Moringa oleifera* leaves meal enhanced broiler performance, physiology and improved resistance to heat stress conditions [23]. As Helal et al. [43] reported beneficial role of moringa growth indices, improve immune responses and reduced lipid peroxidation owing to its antioxidant potency. Mistletoe has been reported to boost antioxidant profile and inhibits oxidative stress in laying pullets [44]. The antioxidant improvement of herb on birds was reported with upregulation of antioxidant activity by ginger supplementation in heat stressed birds to lower malondialdehyde [45]. Similarly, Akbarian et al. [46] found that enhancement with dietary essential oils promoted antioxidant defense against heat stress-induced pathophysiology of heat stressed broilers.

The promoted antioxidant profile of heat stressed chickens fed on phytogetic supplements was achieved using phenolic compounds, like flavonoids [47]. Lowered cholesterol and lipid peroxidation by phytogetics could be associated with apparatus of soursop juice decrease of cholesterol by contributing to reducing lipid peroxidation, shows hypocholesterolemia is related to low lipid peroxidation [19]. Similarly, Selim et al. [25] opined that moringa inclusion in rabbits diet lowered blood lipids; Triglycerides, cholesterol, and LDL, elevates blood antioxidant status. Ameliorating strategies to improve the negative impact of heat stress via enhancing antioxidant cover through exogenous charge of antioxidant substrate and/or its precursors [19], is supported by this study.

CONCLUSION

The findings of this study shows that feeding broiler chicken with phytogetic supplements did ameliorate the negative effects of thermal discomfort on productive performance, haematology and physiological indices of heat stressed broilers. Productive index of heat stressed broilers fed on *Viscum album* supplement were better than other supplements. Thermal stress-induced oxidative stress in broilers were combated by the phytogetic supplements via reduced lipid peroxidation and enhanced antioxidant activity.

REFERENCES

1. Shehata AM, Saadeldin IM, Tukur HA, Habashy WS. Modulation of heat-shock proteins mediates chicken cell survival against thermal stress. *Animals*. 2020;10:2407. <https://doi.org/10.3390/ani10122407>
2. Tekce E, Bayraktar B, Aksakal V, Dertli E, Kamiloğlu A, Topcu KÇ, et al. Response of Japanese quails (*Coturnix coturnix japonica*) to dietary inclusion of *Moringa oleifera* essential oil under heat stress condition. *Ital J Anim Sci*. 2020;19:514-23. <https://doi.org/10.1080/1828051X.2020.1760740>
3. Abreu VMN, Abreu PG. Os desafios da ambiência sobre os sistemas de aves no Brasil. *Rev Bras Zootec*. 2011;40:1-14.
4. Bueno JPR, Gotardo LRM, dos Santos AM, Litz FH, Olivieri OCL, Alves RLOR, et al. Effect of cyclic heat stress on thyroidal hormones, thyroid histology, and performance of two broiler strains. *Int J Biometeorol*. 2020;64:1125-32. <https://doi.org/10.1007/s00484-020-01886-0>
5. Mehaisen GMK, Ibrahim RM, Desoky AA, Safaa HM, El-Sayed OA, Abass AO. The importance of propolis in alleviating the negative physiological effects of heat stress in quail chicks. *PLOS ONE* 2017;12:e0186907. <https://doi.org/10.1371/journal.pone.0186907>
6. Saeed M, Abbas G, Alagawany M, Kamboh AA, El-Hack MEA, Khafaga AF, et al. Heat

- stress management in poultry farms: a comprehensive overview. *J Therm Biol.* 2019;84:414-25. <https://doi.org/10.1016/j.jtherbio.2019.07.025>
7. Surai PF, Fisinin VI. Vitagenes in poultry production: part 1. technological and environmental stresses. *Worlds Poult Sci J.* 2016;72:721-33. <https://doi.org/10.1017/S0043933916000714>
 8. Surai PF, Fisinin VI. Vitagenes in poultry production: part 2. nutritional and internal stresses. *Worlds Poult Sci J.* 2016;72:761-72. <https://doi.org/10.1017/S0043933916000726>
 9. Surai PF, Kochish II, Fisinin VI, Kidd MT. Antioxidant defence systems and oxidative stress in poultry biology: an update. *Antioxidants.* 2019;8:235. <https://doi.org/10.3390/antiox8070235>
 10. Jang IS, Ko YH, Moon YS, Sohn SH. Effects of vitamin C or E on the pro-inflammatory cytokines, heat shock protein 70 and antioxidant status in broiler chicks under summer conditions. *Asian-Australas J Anim Sci.* 2014;27:749-56. <https://doi.org/10.5713/ajas.2013.13852>
 11. Vandana GD, Sejian V, Lees AM, Pragna P, Silpa MV, Maloney SK. Heat stress and poultry production: impact and amelioration. *Int J Biometeorol.* 2021;65:163-79. <https://doi.org/10.1007/s00484-020-02023-7>
 12. Jimoh OA, Oyeyemi WA, Okin-Aminu HO, Oyeyemi BF. Reproductive characteristics, semen quality, seminal oxidative status, steroid hormones, sperm production efficiency of rabbits fed herbal supplements. *Theriogenology.* 2021;168:41-9. <https://doi.org/10.1016/j.theriogenology.2021.03.020>
 13. Jimoh OA, Oyeyemi BF, Oyeyemi WA. Soursop juice enhanced seminal antioxidant defence and semen quality of rabbit bucks in extremely dry climatic condition of Southwestern Nigeria. *J Therm Biol.* 2021;100:103034. <https://doi.org/10.1016/j.jtherbio.2021.103034>
 14. Olugbemi TS, Mutayoba SK, Lekule FP. Effect of moringa (*Moringa oleifera*) inclusion in cassava based diets fed to broiler chickens. *Int J Poult Sci.* 2010;9:363-7. <https://doi.org/10.3923/ijps.2010.363.367>
 15. Melesse A, Getye Y, Berihun K, Banerjee S. Effect of feeding graded levels of *Moringa stenopetala* leaf meal on growth performance, carcass traits and some serum biochemical parameters of Koekoek chickens. *Livest Sci.* 2013;157:498-505. <https://doi.org/10.1016/j.livsci.2013.08.012>
 16. Stoyanova S, Geuns J, Hideg É, Van den Ende W. The food additives inulin and stevioside counteract oxidative stress. *Int J Food Sci Nutr.* 2011;62:207-14. <https://doi.org/10.3109/09637486.2010.523416>
 17. Jimoh OA, Ewuola EO. Thermoregulatory response of exotic rabbit breeds during peak temperature - humidity index of Ibadan. *Trop Anim Prod Invest.* 2016;19:41-7.
 18. Jimoh OA, Ewuola EO, Balogun AS. Oxidative stress markers in exotic breeds of rabbit during peak of heat stress in Ibadan, Nigeria. *J Adv Biol Biotechnol.* 2017;12:1-9. <https://doi.org/10.9734/JABB/2017/30437>
 19. Jimoh OA, Ayedun ES, Oyelade WA, Oloruntola OD, Daramola OT, Ayodele SO, et al. Protective effect of soursop (*Annona muricata* Linn.) juice on oxidative stress in heat stressed rabbits. *J Anim Sci Technol.* 2018;60:28. <https://doi.org/10.1186/s40781-018-0186-4>
 20. Jimoh OA, Ewuola EO. Semen characteristics, seminal biochemical and oxidative stress markers in rabbits during heat stress. *J Vet Androl.* 2018;3:35-44.
 21. Jimoh AO, Ojo OA, Ihejirika UD. Metabolic and oxidative status of West African dwarf does at different reproductive stages in southwest Nigeria. *Bull Natl Res Cent.* 2019;43:190. <https://doi.org/10.1186/s42269-019-0223-6>
 22. Mbikay M. Therapeutic potential of *Moringa oleifera* leaves in chronic hyperglycemia and dyslipidemia: a review. *Front Pharmacol.* 2012;3:24. <https://doi.org/10.3389/fphar.2012.00024>

23. Abd El-Hack ME, Abdelnour SA, Taha AE, Khafaga AF, Arif M, Ayasan T, et al. Herbs as thermoregulatory agents in poultry: an overview. *Sci Total Environ.* 2020;703:134399. <https://doi.org/10.1016/j.scitotenv.2019.134399>
24. Saleh AA, Ebeid TA, Abudabos AM. Effect of dietary phytochemicals (herbal mixture) supplementation on growth performance, nutrient utilization, antioxidative properties, and immune response in broilers. *Environ Sci Pollut Res.* 2018;25:14606-13. <https://doi.org/10.1007/s11356-018-1685-z>
25. Selim S, Seleiman MF, Hassan MM, Saleh AA, Mousa MA. Impact of dietary supplementation with *Moringa oleifera* leaves on performance, meat characteristics, oxidative stability, and fatty acid profile in growing rabbits. *Animals.* 2021;11:248. <https://doi.org/10.3390/ani11020248>
26. Gouda A, El-Moniary MM, Youssef AW, Hamouda Y, Hassan HMA, El-Daly EF. Response of broiler chicks to diets supplemented with *Moringa oleifera* dry leaves and some antioxidants under tropical summer conditions. *Biosci Res.* 2018;15:637-44.
27. North MK, Dalle Zotte A, Hoffman LC. Effect of quercetin supplementation on the growth, feed efficiency and serum hormone levels of New Zealand White rabbits. *S Afr J Anim Sci.* 2018;48:1128-39. <https://doi.org/10.4314/sajas.v48i6.17>
28. Cui YM, Wang J, Lu W, Zhang HJ, Wu SG, Qi GH. Effect of dietary supplementation with *Moringa oleifera* leaf on performance, meat quality, and oxidative stability of meat in broilers. *Poult Sci.* 2018;97:2836-44. <https://doi.org/10.3382/ps/pey122>
29. Vedendar G. Effect of supplementation of *Moringa oleifera* leaf powder on performance and antioxidant activity to alleviate heat stress in broilers. *PVNarsimha Rao Telangana Veterinary University.* 2018;116-30.
30. Daramola OT, Jimoh AO, Oloruntola OD, Ayodele SO. Effect of herbal supplement on the laying performance and antioxidant status of serum and egg yolk of laying birds. *Biotechnol J Int.* 2018;21:1-9. <https://doi.org/10.9734/BJI/2018/41401>
31. Hassan HMA, El-Moniary MM, Hamouda Y, El-Daly EF, Youssef AW, Abd El-Azeem NA. Effect of different levels of *Moringa oleifera* leaves meal on productive performance, carcass characteristics and some blood parameters of broiler chicks reared under heat stress conditions. *Asian J Anim Vet Adv.* 2016;11:60-6. <https://doi.org/10.3923/ajava.2016.60.66>
32. Niu ZY, Liu FZ, Yan QL, Li WC. Effects of different levels of vitamin E on growth performance and immune responses of broilers under heat stress. *Poult Sci.* 2009;88:2101-7. <https://doi.org/10.3382/ps.2009-00220>
33. Hosny NS, Hashem NM, Morsy AS, Abo-elezz ZR. Effects of organic selenium on the physiological response, blood metabolites, redox status, semen quality, and fertility of rabbit bucks kept under natural heat stress conditions. *Front Vet Sci.* 2020;7:290. <https://doi.org/10.3389/fvets.2020.00290>
34. Belhadj Slimen I, Najjar T, Ghram A, Abdrrabba M. Heat stress effects on livestock: molecular, cellular and metabolic aspects, a review. *J Anim Physiol Anim Nutr.* 2015;100:401-12. <https://doi.org/10.1111/jpn.12379>
35. El-Desoky NI, Hashem NM, Elkomy A, Abo-elezz ZR. Physiological response and semen quality of rabbit bucks supplemented with *Moringa* leaves ethanolic extract during summer season. *Animal.* 2017;11:1549-57. <https://doi.org/10.1017/S1751731117000088>
36. Habibi R, Sadeghi G, Karimi A. Effect of different concentrations of ginger root powder and its essential oil on growth performance, serum metabolites and antioxidant status in broiler chicks under heat stress. *Br Poult Sci.* 2014;55:228-37. <https://doi.org/10.1080/00071668.2014.887830>

37. Hajati H, Hassanabadi A, Golian A, Nassiri-Moghaddam H, Nassiri MR. The effect of grape seed extract and vitamin C feed supplementation on some blood parameters and HSP70 gene expression of broiler chickens suffering from chronic heat stress. *Ital J Anim Sci.* 2015;14:3273. <http://doi.org/104081/ijas2014.3273>
38. Tayer M, Nobakht A, Pishjangh J. Effects of different levels of green grape (*Vitis vinifera*) leaves on performance, carcass traits, blood biochemical and immunity parameters of broilers. *Int Res J Appl Basic Sci.* 2012;3:103-11.
39. Ferit Gursu M, Onderci M, Gulcu F, Sahin K. Effects of vitamin C and folic acid supplementation on serum paraoxonase activity and metabolites induced by heat stress in vivo. *Nutr Res.* 2004;24:157-64. <https://doi.org/10.1016/j.nutres.2003.11.008>
40. Vergara-Jimenez M, Almatrafi MM, Fernandez ML. Bioactive components in *Moringa oleifera* leaves protect against chronic disease. *Antioxidants.* 2017;6:91. <https://doi.org/10.3390/antiox6040091>
41. Halaby MS, Elmetwaly EM, Omar AAA. Effect of *Moringa oleifera* on serum lipids and kidney function of hyperlipidemic rats. *J Appl Sci Res.* 2013;9:5189-98.
42. Mujahid A, Pumford NR, Bottje W, Nakagawa K, Miyazawa T, Akiba Y, et al. Mitochondrial oxidative damage in chicken skeletal muscle induced by acute heat stress. *J Poult Sci.* 2007;44:439-45. <https://doi.org/10.2141/jpsa.44.439>
43. Helal FIS, El-Badawi AY, El-Wardany I, Ali NGM, Aboelazab OM. Effect of dietary moringa (*Moringa oleifera*) and rosemary (*Rosmarinus officinalis*) leaves or their mixture on productive performance, carcass characteristics and antioxidant enzymes of rabbits reared under heat stress conditions. *Agric Eng Int CIGR J.* 2017;184-92.
44. Jimoh OA, Ihejirika UG, Balogun AS, Uwaeziozi UC. Antioxidative effect of mistletoe leaf meal supplemented diets in laying pullets. *Arch Zootec.* 2018;67:526-30. <https://doi.org/10.21071/az.v0i0.3883>
45. Ibtisham F, Nawab A, Niu Y, Wang Z, Wu J, Xiao M. The effect of ginger powder and Chinese herbal medicine on production performance, serum metabolites and antioxidant status of laying hens under heat-stress condition. *J Therm Biol.* 2019;81:20-4. <https://doi.org/10.1016/j.jtherbio.2019.02.002>
46. Akbarian A, Michiels J, Golian A, Buyse J, Wang Y, De Smet S. Gene expression of heat shock protein 70 and antioxidant enzymes, oxidative status, and meat oxidative stability of cyclically heat-challenged finishing broilers fed *Origanum compactum* and *Curcuma xanthorrhiza* essential oils. *Poult Sci.* 2014;93:1930-41. <https://doi.org/10.3382/ps.2014-03896>
47. Kamboh AA, Hang SQ, Bakhsetgul M, Zhu WY. Effects of genistein and hesperidin on biomarkers of heat stress in broilers under persistent summer stress. *Poult Sci.* 2013;92:2411-8. <https://doi.org/10.3382/ps.2012-02960>