

ARTICLE INFORMATION	Fill in information in each box below
Article Type	Research article
Article Title (within 20 words without abbreviations)	Evaluation of <i>In vitro</i> Method for Estimating <i>In vivo</i> Digestibility in Mixed-Breed Dogs across Each Life Stage
Running Title (within 10 words)	Validating <i>In vitro</i> Digestibility Method for Dogs
Author	Kyeongho Jeon ^{1, #} , Minho Song ^{2, #} , Jihwan Lee ^{3, #} , Dongcheol Song ¹ , Hyuck Kim ¹ , Jinmo Yang ¹ , Hyohyeon Yu ¹ , Hyeunbum Kim ^{4, *} , Jinho Cho ^{1, *}
Affiliation	¹ Department of animal science, Chungbuk National University, Cheongju 28644, Korea ² Division of Animal and Dairy Science, Chungnam National University, Daejeon 34134, Korea ³ Department of Animal Science, Jeonbuk National University, Jeonju 54896, Korea ⁴ Department of Animal Resources Science, Dankook University, Cheonan 31116, Korea
ORCID (for more information, please visit https://orcid.org)	Kyeongho Jeon / jeonkh1222@gmail.com (https://orcid.org/0000-0003-2321-3319) Minho Song / mhsong@cnu.ac.kr (https://orcid.org/0000-0002-4515-5212) Jihwan Lee/ jl26112@jbnu.ac.kr (https://orcid.org/0000-0001-8161-4853) Dongcheol Song / paul741@daum.net (https://orcid.org/0000-0002-5704-603X) Hyuck Kim / harrck85@naver.com (https://orcid.org/0000-0002-5280-0734) Jinmo Yang / mike000315@gmail.com (https://orcid.org/0009-0007-4272-3441)

	<p>Hyohyeon Yu / dbgygus123@naver.com (https://orcid.org/0009-0004-7633-8013)</p> <p>Hyeunbum Kim / hbkim@dankook.ac.kr (https://orcid.org/0000-0003-1366-6090)</p> <p>Jinho Cho / jinhcho@cbnu.ac.kr (http://orcid.org/0000-0001-7151-0778)</p>
Competing interests	No potential conflict of interest relevant to this article was reported.
Funding sources State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available.	This work was carried out with the support of ‘Cooperative Research Program for Agriculture Science and Technology Development (Project No. RS-2023-00230754)’ Rural Development Administration, Republic of Korea.
Acknowledgements	Not applicable.
Availability of data and material	All data generated or analyzed during this study are included in this published article.
Authors' contributions Please specify the authors’ role using this form.	<p>Conceptualization: KH Jeon, JH Lee, MH Song, JH Cho</p> <p>Data curation: H Kim, JM Yang</p> <p>Formal analysis: HH Yu, DC Song</p> <p>Methodology: DC Song, HH Yu</p> <p>Software: JH Lee, JM Yang</p> <p>Validation: H Kim, KH Jeon</p> <p>Investigation: HB Kim, JH Cho</p> <p>Writing - original draft: KH Jeon, JH Lee, HB Kim</p> <p>Writing - review & editing: KH Jeon, MH Song, JH Lee, DC Song, H Kim, JM Yang, HH Yu, HB Kim, JH Cho</p>
Ethics approval and consent to participate	This experiment received approval (approval # 202412-CNU-409) from the Institutional Animal Care and Use Committee of Chungnam National University

For the corresponding author (responsible for correspondence, proofreading, and reprints)	Fill in information in each box below
First name, middle initial, last name	Jinho Cho Hyeunbum Kim
Email address – this is where your proofs will be sent	jinhcho@chungbuk.ac.kr hbkim@dankook.ac.kr
Secondary Email address	
Address	Department of animal science, Chungbuk National University, Cheongju 28644, Korea Department of Animal Resources Science, Dankook University, Cheonan 31116, Korea
Cell phone number	+82-10-5488-8580 (Jinho Cho)
Office phone number	+82-43-261-2544 (Jinho Cho) +85-41-550-3632 (Hyeunbum Kim)
Fax number	+82-43-273-2240 (Jinho Cho) +85-41-550-3632 (Hyeunbum Kim)

ACCEPTED

4 **Abstract**

5 This study aimed to evaluate the agreement between a two-step *in vitro* digestion procedure and *in vivo*
6 apparent total tract digestibility (ATTD) across three life stages in mixed-breed dogs. Eighteen dogs were
7 divided into three groups: puppies (n = 6; < 1 year; initial body weight (iBW): 10.68 ± 1.55 kg), adults (n = 6;
8 2–7 years; iBW: 8.34 ± 0.38 kg), and seniors (n = 6; > 8 years; iBW: 8.87 ± 0.98 kg). An extruded diet based on
9 oat, turkey, and chicken breast meal was used for both *in vitro* and *in vivo* digestibility determinations of dry
10 matter (DM), organic matter (OM), crude protein (CP), gross energy (GE), crude fiber (CF), and ether extract
11 (EE). *In vitro* digestibility values were consistently higher than *in vivo* values across all nutrients and life stages
12 ($p < 0.05$). Linear regression revealed strong *in vitro*–*in vivo* agreement for gross energy across all life stages (r^2
13 = 0.87–0.98), for organic matter in adult dogs ($r^2 = 0.96$), and for ether extract and dry matter in senior dogs (r^2
14 = 0.91–0.93), as well as for crude fiber in puppies and adults ($r^2 = 0.85$ –0.87) and for ether extract in puppies (r^2
15 = 0.91), whereas agreement was weaker for crude protein across life stages ($r^2 = 0.55$ –0.66) and for organic
16 matter in puppies ($r^2 = 0.40$). These findings indicate that the two-step *in vitro* digestion method can serve as a
17 practical screening tool for energy and macronutrient digestibility in mixed-breed dogs of the body-size range
18 studied here, with predictive accuracy that depends on both the nutrient and the life stage.

19

20 Keywords: *in vitro* digestibility, *in vivo* digestibility, nutrient digestibility, mixed-breed dog, life stage

21

Introduction

Nutrient digestibility is a key factor in assessing the nutritional quality of companion animal diets, as it determines the proportion of ingested nutrients available for metabolic use [1]. Apparent total tract digestibility (ATTD) significantly impacts feed efficiency, fecal characteristics, and overall nutritional status. Meanwhile, undigested residues that reach the hindgut undergo microbial fermentation, producing metabolites that influence gut health and fecal quality [2, 3]. Highly digestible diets are therefore particularly important for companion dogs, including the heterogeneous mixed-breed populations that account for a substantial fraction of the global pet-dog population [4].

Body size and breed are well-known modulators of canine digestive physiology: small-breed dogs exhibit higher mass-specific basal metabolic rates and shorter gastrointestinal transit times than large-breed dogs, while large-breed dogs have more developed cecal and colonic compartments that prolong transit time and increase fermentative activity [4, 5, 6]. These anatomical and functional differences suggest that digestive efficiency may vary across breeds; however, systematic *in vivo* digestibility data for the heterogeneous mixed-breed populations that dominate the global pet-dog population remain scarce.

Among breeds, copy-number variation in the amylase gene has been associated with breed-level differences in carbohydrate digestive capacity [8], implying that breed-specific adaptations may extend beyond carbohydrate metabolism to overall nutrient utilization. Validation of *in vitro* digestibility methods against *in vivo* data in genetically diverse populations is therefore needed if the resulting equations are to be applied outside of single-breed reference colonies.

In vitro digestibility methodologies have become accepted alternatives to *in vivo* trials due to their standardized conditions, rapid throughput, and reduced ethical concerns [9]. Previous studies have demonstrated a strong correlation between *in vitro* and *in vivo* digestion rates, supporting the validity of *in vitro* methods as a reliable approach for evaluating comparative digestibility [10, 11].

Most existing canine digestibility research has used Beagles as a standardized model or studied populations without explicit breed specification [1, 13], leaving the *in vivo* digestibility profile of mixed-breed dogs—and its agreement with *in vitro* estimates—largely uncharacterized across life stages. We hypothesized that the agreement between two-step *in vitro* digestion and *in vivo* ATTD would vary with both the nutrient and the life stage in mixed-breed dogs. Therefore, the present study was conducted to evaluate, across three life stages (puppies, adults, and seniors), the level of agreement between a two-step enzymatic *in vitro* digestion method and *in vivo* apparent

51 total tract digestibility for dry matter, organic matter, crude protein, gross energy, crude fiber, and ether extract in
52 mixed-breed dogs, and to derive life-stage-specific regression equations linking the two approaches.
53

ACCEPTED

54 **Materials and Methods**

55

56 **Experimental diet**

57 The experimental diet, comprising oat, turkey, and chicken breast meal as primary ingredients, was
58 formulated in our laboratory specifically for this study. The diet was formulated to meet the AAFCO [14]
59 nutrient profile for all life stages including growth, ensuring that nutrient supply was adequate for puppies,
60 adult, and senior dogs without re-formulation (Table 1). Manufacturing was carried out under a contract
61 arrangement at a pilot-scale extrusion facility, following a standard sequence of dry-ingredient grinding through
62 a 1.0-mm screen, mixing in a horizontal ribbon mixer, steam conditioning, single-screw extrusion, drying to a
63 shelf-stable moisture, perilla-oil coating, and ambient cooling.

64

65 ***In vitro* method**

66 A two-step *in vitro* digestion procedure adapted from Hervera et al. [15] was performed in six independent
67 replicates of the experimental diet. Each replicate consisted of a separately weighed subsample of the same bulk
68 experimental diet, processed through the full two-step procedure described below.

69 Preparation phase: Prior to digestion, all samples were oven-dried at 65°C to a constant weight and subsequently
70 ground into a fine powder with a particle size of less than 1.0 mm.

71 Gastric phase: Each container received 25 mL of phosphate buffer (0.1 M, pH 6.0) together with 10 mL of HCl
72 solution (0.2 M, pH 0.7). The pH was brought to 2.0 by titrating with 1 M HCl and 1 M NaOH. Gastric digestion
73 was initiated by the addition of 1 mL of pepsin solution (10 mg/mL; ≥ 250 units/mg solid, P7000, pepsin from
74 porcine gastric mucosa; Sigma-Aldrich). To inhibit microbial contamination, 1 mL of chloramphenicol solution
75 (C0378, chloramphenicol; Sigma-Aldrich; 5 g/L in ethanol) was also introduced. Flasks were then covered with
76 Parafilm M® film and incubated at 39°C for 2 h in a shaking incubator (SWB-35; Hanyang Science Lab Co.).

77 Small intestinal phase: Once cooled to room temperature, each flask received 5 mL of 0.6 M NaOH and 10 mL
78 of phosphate buffer (0.2 M, pH 6.8), followed by pH adjustment to 6.8 with 1 M HCl and NaOH. Intestinal
79 digestion was then simulated by adding 1 mL of pancreatin solution (100 mg/mL; 4 × USP, P1750, pancreatin
80 from the porcine pancreas; Sigma-Aldrich). The flasks were sealed with Parafilm M® film and incubated at 39°C
81 for 4 h with continuous agitation.

82 Sample collection and filtration phase: Following digestion, undigested residues were recovered by vacuum
83 filtration through pre-weighed, pre-dried glass filter crucibles (Gooch Type Filter Crucibles, PYREX®). Each

84 flask was rinsed three times with distilled water during the filtration step. Residues in the crucibles were
85 subsequently washed with two sequential additions of 10 mL of 95% ethanol and 10 mL of 99.5% acetone.

86

87 **Chemical analyses and calculation**

88 To determine DM content following *in vitro* digestion, filter crucibles containing undigested residues were
89 oven-dried at 70°C for 24 h. Subsequent ashing at 550°C for 4 h was conducted to obtain Organic matter (OM)
90 values. Crucibles were cooled to room temperature before weighing. All proximate analyses followed AOAC
91 methods [16]: Dry matter (DM; method 930.15), OM (method 942.05), Crude Fiber (CF; method 978.10), and
92 Ether extract (EE; method 920.39). CP was quantified by the Dumas combustion method (Rapid MAX N-Exceed,
93 Elementar), and GE was measured using a bomb calorimeter (Parr 6400 Bomb Calorimeter, Parr Instrument Co.).

94 *In vitro* DM digestibility was calculated as follows:

$$95 \quad \text{“Digestibility (\%)} = 100 - \{(\text{residue weight} / \text{sample weight}) \times 100\}”$$

96 *In vitro* digestibility of OM, CP, GE, CF, and EE was derived using the equation below:

$$97 \quad \text{“Digestibility (\%)} = 100 - \{Nr \times (100 - \text{IDDM}) / Nd\}”$$

98 where Nr = nutrient concentration in residues (DM %), Nd = nutrient concentration in diet (DM %), and IDDM
99 = *in vitro* DM digestibility (%).

100

101 ***In vivo* method**

102 **Animal ethics**

103 This experiment received approval (approval # 202412-CNU-409) from the Institutional Animal Care and
104 Use Committee of Chungnam National University, Daejeon, Korea. Dogs were housed and handled in
105 accordance with the approved procedures throughout the study.

106

107

108 **Animals and experiment design**

109 Eighteen mixed-breed dogs of comparable medium body size (mature weight approximately 8–11 kg) were
110 used in this study. The dogs were derived from medium-sized parent stock, were of unidentified pedigree, and
111 showed no dominant morphological traits of any single recognized breed. Animals were allocated to three life-
112 stage groups: six puppies (< 1 year old; initial body weight, BW: 10.68 ± 1.55 kg), six adults (2–7 years old;
113 BW: 8.34 ± 0.38 kg), and six seniors (> 8 years old; BW: 8.87 ± 0.98 kg). Within each life stage, sex was
114 balanced (3 males and 3 females). Each dog was individually housed in a kennel maintained at 23°C. The total
115 study duration was 17 days, with the first 7 days designated as an adaptation period. Maintenance energy
116 requirements (MER) were estimated on the basis of metabolic BW (mBW) using the following equations:

117 Calculating the MER used the following formula:

118 “Puppies = $132 \times \text{mBW} (\text{BW}^{0.75}) \times 1.5$; Adult dogs = $132 \times \text{mBW} (\text{BW}^{0.75})$; Senior dogs = $105 \times \text{mBW}$
119 $(\text{BW}^{0.75})$ ”.

120 Daily feed requirements of each dog were determined using MER, and the dogs were fed twice a day at 9:00
121 and 17:00.

123 **Nutrient digestibility**

124 The ATTD of DM, OM, CP, GE, CF, and EE were determined using 0.5% Cr₂O₃ incorporated into the diet as
125 an indigestible external marker. Fresh fecal samples were collected on days 3 through 6 of the collection period.
126 Both freshly collected fecal samples and the corresponding diet samples were immediately frozen at –20°C for
127 subsequent analysis. Upon completion of the experiment, fecal samples were oven-dried at 70°C for 72 h and
128 ground through a 1 mm screen. Nutrient digestibility of DM, OM, CP, GE, CF, and EE was analyzed using the
129 dried fecal samples. Analytical methods for DM (method 930.15), OM (method 942.05), CF (method 978.10),
130 and EE (method 920.39) followed AOAC methods [16]. CP and GE were determined by Dumas combustion
131 (Rapid MAX N-Exceed, Elementar) and bomb calorimetry (Parr 6400 Bomb Calorimeter, Parr Instrument Co.),
132 respectively.

133 Calculating the ATTD digestibility of nutrients used the following formula:

134 “Digestibility = $1 - [(\text{Nf} \times \text{Cd}) / (\text{Nd} \times \text{Cf})] \times 100$ ”

135 where Nf = concentration of nutrient in fecal, Nd = concentration of nutrient in the diet, Cd = concentration of
136 Cr₂O₃ in the diet, and Cf = concentration of Cr₂O₃ in the fecal.

137

138 **Statistical analysis**

139 For the *in vivo* experiment, the individual dog was considered the experimental unit; for the *in vitro*
140 experiment, each independent digestion replicate was treated as the experimental unit. Differences in
141 digestibility among life-stage groups were tested by one-way analysis of variance (ANOVA) followed by
142 Tukey's honestly significant difference (HSD) test for pairwise comparisons. The assumptions of normality of
143 residuals (Shapiro–Wilk test) and homogeneity of variance (Levene's test) were verified prior to ANOVA;
144 where assumptions were violated, the non-parametric Kruskal–Wallis test was applied as a sensitivity check,
145 and the conclusions were unchanged. Orthogonal contrasts were used to compare the *in vitro* method against the
146 *in vivo* measurements within each life stage. The relationship between the two methods was examined by linear
147 regression analysis within a general linear model (GLM) framework. All statistical computations were carried
148 out using JMP® Pro version 16.0.0 (SAS Institute Inc., Cary). Differences were regarded as statistically
149 significant at $p < 0.05$.

150

ACCEPTED

151 **Results**

152 ***In vitro* and *in vivo* Digestibility**

153 The *in vitro* and *in vivo* digestibility of DM, OM, CP, GE, CF and EE in puppies, adult dogs, and senior dogs
154 are presented in Table 2. The *in vitro* digestibility of DM, OM, CP, CF, and EE was significantly higher ($p <$
155 0.05) than *in vivo* digestibility in all life stages. The *in vitro* digestibility of GE was also significantly higher (p
156 < 0.05) than *in vivo* digestibility in all life stages except puppies.

157 Among *in vivo* measurements, a significant life-stage effect was detected only for gross energy digestibility (p
158 < 0.013), with senior dogs showing a lower value than adult dogs ($p < 0.05$). For the remaining nutrients (DM,
159 OM, CP, CF, and EE), no significant life-stage difference was observed ($p > 0.05$; Table 3).

160

161 **The relationships between *in vitro* and *in vivo* digestibility**

162 The statistical relationships between *in vitro* and *in vivo* digestibility as linear regression equations are shown
163 in Table 4. In puppies, strong correlations were observed for GE, CF, and EE ($r^2 = 0.93, 0.87, \text{ and } 0.91,$
164 respectively), whereas DM, OM, and CP showed moderate correlations ($r^2 = 0.60, 0.40, \text{ and } 0.55, \text{ respectively}$).
165 In adult dogs, OM and GE exhibited the strongest correlations ($r^2 = 0.96 \text{ and } 0.98, \text{ respectively}$), followed by CF
166 ($r^2 = 0.85$), while DM, CP, and EE showed moderate correlations ($r^2 = 0.67, 0.56, \text{ and } 0.51, \text{ respectively}$). In
167 senior dogs, strong correlations were observed for most nutrients, with EE, DM, and GE showing the highest
168 values ($r^2 = 0.93, 0.91, \text{ and } 0.87, \text{ respectively}$), whereas CP exhibited a moderate correlation ($r^2 = 0.66$).

169

170 **Discussion**

171 The present study assessed and compared nutrient digestibility in mixed-breed dogs at three life stages: puppies,
172 adults, and seniors, both *in vitro* and *in vivo*. Consistently, the *in vitro* digestibility values were higher than the
173 corresponding *in vivo* values across all measured nutrients, aligning with findings from previous studies [15, 17,
174 18]. This systematic overestimation in *in vitro* methods can be attributed to the lack of consideration for
175 endogenous nitrogen and fat losses that occur *in vivo*. In living animals, fecal output includes not only undigested
176 dietary residues but also epithelial cell debris, mucus secretions, and bacterial biomass [19, 20]. Furthermore, the
177 two-step enzymatic procedure does not account for hindgut microbial activity, where resident bacteria modify
178 undigested substrates through fermentation [21, 22]. Additionally, the static nature of *in vitro* systems fails to
179 replicate the dynamic variations in enzyme secretion, gastric emptying rates, and intestinal motility that occur in
180 response to meal composition and individual physiological status [23, 24].

181 Contrast analysis showed a progressive decline in *in vivo* digestibility values from puppies to seniors for most
182 nutrients. This age-related decrease in digestive efficiency can be attributed to several physiological changes
183 associated with aging, including reduced pancreatic enzyme secretion, decreased brush border enzyme activity,
184 and a diminished intestinal absorptive surface area due to villous atrophy [25, 26]. Prior studies have also shown
185 that senior dogs experience alterations in gut microbiota composition, characterized by reduced bacterial diversity
186 and lower populations of beneficial fermentative bacteria [3, 27], which may further hinder nutrient utilization
187 efficiency. The smallest difference between *in vitro* and *in vivo* values observed in puppies suggests that their
188 heightened digestive capacity during growth, marked by increased enzyme secretion and enhanced absorptive
189 function, more closely resembles the optimal conditions simulated *in vitro* [28]. The rapid tissue accretion and
190 organ development during the growth phase necessitate maximal nutrient extraction efficiency, and the developing
191 gastrointestinal system appears optimized for high digestive performance [29, 30].

192 The strong *in vitro*–*in vivo* agreement observed in senior dogs (DM, GE, EE $r^2 = 0.87$ – 0.93), despite their
193 lower mean digestibility, can be reconciled by noting that the *in vitro* method captures relative differences in
194 substrate susceptibility to enzymatic hydrolysis, whereas the absolute reduction in *in vivo* digestibility in senior
195 dogs reflects age-related changes in pancreatic enzyme secretion, mucosal surface area, and microbial activity
196 that act as a roughly uniform downward shift across samples. Because such a uniform shift preserves the rank
197 order of samples, the slope of the regression line is reduced but its r^2 remains high. The relatively narrow within-
198 group variability among senior dogs further contributed to the high coefficients of determination.

199 The linear regression analysis revealed that gross energy and organic matter digestibility had the strongest
200 correlations between *in vitro* and *in vivo* methods. This suggests that the two-step enzymatic procedure most
201 effectively simulates the digestion of energy-yielding macronutrients. The sequential hydrolysis using pepsin and
202 pancreatin accurately mimics the gastric and small intestinal phases, where most lipid and carbohydrate digestion
203 occurs. The chemical bonds cleaved during *in vitro* incubation closely resemble the enzymatic reactions occurring
204 *in vivo* [9, 15]. CP digestibility showed moderate correlations, likely due to the complexity of protein digestion,
205 which involves the sequential action of multiple proteolytic enzymes with specific substrate affinities.
206 Additionally, the resistance of different protein sources to enzymatic hydrolysis varies based on their amino acid
207 composition, tertiary structure, and thermal processing history [13, 31]. Prior studies have indicated that protein
208 digestibility is particularly sensitive to ingredient quality and processing conditions. For instance, Maillard
209 reaction products formed during extrusion can reduce protein bioavailability in ways not fully captured by *in vitro*
210 methods [19, 32]. Adult dogs exhibited the most consistent correlations across all nutrient parameters, reflecting
211 the physiological stability characteristic of the maintenance life stage. At this stage, digestive enzyme profiles,
212 gut microbiota composition, and intestinal morphology reach a mature equilibrium [33, 34].

213 Because all observations originated from a single extruded oat–poultry-based diet, the regression coefficients
214 reported here should be interpreted as life-stage–specific calibrations valid for diets of comparable composition,
215 rather than as universal predictive equations. Within the present scope, the regression equations describe the
216 distributional agreement between *in vitro* and *in vivo* digestibility values and may help reduce the need for repeated
217 *in vivo* trials when evaluating closely related formulations [1]. The equations derived from adult dogs showed the
218 highest agreement, making them the most informative reference among the three life-stage models for quality
219 control of pet food formulations of similar matrix [24, 32].

220 It should be noted that the sequential ethanol and acetone washes used in the filtration step of the present *in*
221 *vitro* protocol (adapted from Hervera et al. [15]) may solubilize part of the residual lipid fraction, potentially
222 shifting the absolute *in vitro* EE digestibility upward. Because all *in vitro* samples were processed identically,
223 this offset is expected to be systematic across life stages and should not affect the relative comparison or the
224 slope of the *in vivo*–*in vitro* regression. Consistent with this interpretation, the *in vitro*–*in vivo* EE agreement
225 was among the strongest observed in the present dataset ($r^2 = 0.91$ in puppies, $r^2 = 0.93$ in senior dogs),
226 indicating that the *in vitro* EE measurement tracks the corresponding *in vivo* digestibility with high fidelity
227 despite the wash step. The absolute *in vitro* EE values reported here may nonetheless carry a small upward

228 offset relative to a lipid-conserving wash protocol, and a comparative evaluation of the two wash schemes could
229 further refine these absolute estimates in future studies.

230 Additionally, the *in vitro* approach supports systematic comparisons of novel ingredients and processing
231 techniques, aiding in the development of optimized formulations tailored to specific physiological needs [21, 35].
232 Moreover, the use of validated *in vitro* methods can reduce the number of animals needed for nutrient digestibility
233 assessments while maintaining scientific rigor [36].

234

235 **Conclusion**

236 The two-step *in vitro* digestion method showed strong agreement with *in vivo* ATTD for energy and organic
237 matter in adult dogs ($r^2 = 0.96\text{--}0.98$), and for ether extract, dry matter, and energy in senior dogs ($r^2 = 0.87\text{--}$
238 0.93). Agreement was weaker for crude protein in all life stages ($r^2 = 0.55\text{--}0.66$) and for organic matter in
239 puppies ($r^2 = 0.40$), indicating that the predictive value of the *in vitro* method is nutrient- and life-stage-
240 dependent and should not be generalized uniformly. Within the scope of the single experimental diet and the
241 mixed-breed population studied here, the regression equations derived from adult dogs may serve as a practical
242 screening reference for diet evaluation, while broader application to chemically distinct diets will require
243 additional multi-diet validation.

244

245 **Disclosure statement**

246

247 **Acknowledgments**

248 This work was carried out with the support of 'Cooperative Research Program for Agriculture Science and
249 Technology Development (Project No. RS-2023-00230754)' Rural Development Administration, Republic of
250 Korea.

251

252 **References**

- 253 1. Carciofi AC, Takakura F, De-Oliveira L, Teshima E, Jeremias J, Brunetto MA, et al. Effects of six
254 carbohydrate sources on dog diet digestibility and post-prandial glucose and insulin response. *J Anim Physiol*
255 *Anim Nutr.* 2008;92:326-36. <https://doi.org/10.1111/j.1439-0396.2007.00794.x>
- 256 2. Nery J, Goudez R, Biourge V, Tournier C, Leray V, Martin L, et al. Influence of dietary protein content and
257 source on colonic fermentative activity in dogs differing in body size and digestive tolerance. *J Ani Sci.*
258 2012;90:2570-80. <https://doi.org/10.2527/jas.2011-4112>
- 259 3. Middelbos IS, Vester Boler BM, Qu A, White BA, Swanson KS, Fahey Jr GC. Phylogenetic characterization
260 of fecal microbial communities of dogs fed diets with or without supplemental dietary fiber using 454
261 pyrosequencing. *PloS one.* 2010;5:e9768. <https://doi.org/10.1371/journal.pone.0009768>
- 262 4. Weber M, Biourge V, Nguyen P. Digestive sensitivity varies according to size of dogs: a review. *J Anim*
263 *Physiol Anim Nutr.* 2017;101:1-9. <https://doi.org/10.1111/jpn.12507>
- 264 5. Rainbird A, Kienzle E. Studies on the energy requirement of dogs depending on breed and age. 1990.
- 265 6. Deschamps C, Humbert D, Zentek J, Denis S, Priymenko N, Apper E, et al. From Chihuahua to Saint-Bernard:
266 how did digestion and microbiota evolve with dog sizes. *Int J Biol Sci.* 2022;18:5086.
267 <http://doi.org/10.7150/ijbs.72770>
- 268 7. Parker HG, Harris A, Dreger DL, Davis BW, Ostrander EA. The bald and the beautiful: hairlessness in
269 domestic dog breeds. *Philosophical Transactions of the Royal Society B: Biol Sci.* 2017;372.
270 <https://doi.org/10.1098/rstb.2015.0488>
- 271 8. Axelsson E, Ratnakumar A, Arendt M-L, Maqbool K, Webster MT, Perloski M, et al. The genomic signature
272 of dog domestication reveals adaptation to a starch-rich diet. *Nature.* 2013;495:360-4.
273 <https://doi.org/10.1038/nature11837>
- 274 9. Boisen S, Fernández JA. Prediction of the total tract digestibility of energy in feedstuffs and pig diets by *in*
275 *vitro* analyses. *Anim Feed Sci Technol.* 1997;68:277-86. [https://doi.org/10.1016/S0377-8401\(97\)00058-8](https://doi.org/10.1016/S0377-8401(97)00058-8)
- 276 10. Biagi G, Cipollini I, Grandi M, Pinna C, Vecchiato CG, Zaghini G. A new *in vitro* method to evaluate
277 digestibility of commercial diets for dogs. *Ital J Anim Sci.* 2016;15:617-25.
278 <https://doi.org/10.1080/1828051X.2016.1222242>
- 279 11. Jeon KH, Song MH, Lee JH, Chang SY, Song DC, An JW, et al. New *In vitro* Method Complement Low-
280 value Existing *In vitro* Method in Commercial Dog Diets in the Republic of Korea. *J Agric Life Sci.*
281 2024;58:101-10. <https://doi.org/10.14397/jals.2024.58.2.101>

- 282 12. Bellumori TP, Famula TR, Bannasch DL, Belanger JM, Oberbauer AM. Prevalence of inherited disorders
283 among mixed-breed and purebred dogs: 27,254 cases (1995–2010). *J Am Vet Med Assoc.* 2013;242:1549-55.
- 284 13. Clapper G, Grieshop C, Merchen N, Russett J, Brent Jr J, Fahey Jr G. Ileal and total tract nutrient digestibilities
285 and fecal characteristics of dogs as affected by soybean protein inclusion in dry, extruded diets. *J Anim Sci.*
286 2001;79:1523-32. <https://doi.org/10.2527/2001.7961523x>
- 287 14. AAFCO (Association of American Feed Control Officials). Proceedings of the AAFCO. 2024: Champaign,
288 IL: AAFCO.
- 289 15. Hervera M, Baucells M, Blanch F, Castrillo C. Prediction of digestible energy content of extruded dog food
290 by *in vitro* analyses. *J Anim Physiol Anim Nutr.* 2007;91:205-9. [https://doi.org/10.1111/j.1439-](https://doi.org/10.1111/j.1439-0396.2007.00693.x)
291 [0396.2007.00693.x](https://doi.org/10.1111/j.1439-0396.2007.00693.x)
- 292 16. AOAC (Association of Official Analytical Chemists) International. Official methods of analysis of the AOAC
293 International. 18th ed AOAC International. 2006
- 294 17. De-Oliveira L, Takakura F, Kienzle E, Brunetto M, Teshima E, Pereira G, et al. Fibre analysis and fibre
295 digestibility in pet foods—a comparison of total dietary fibre, neutral and acid detergent fibre and crude fibre.
296 *J Anim Physiol Anim Nutr.* 2012;96:895-906. <https://doi.org/10.1111/j.1439-0396.2011.01203.x>
- 297 18. Castrillo C, Baucells M, Vicente F, Muñoz F, Andueza D. Energy evaluation of extruded compound foods for
298 dogs by near-infrared spectroscopy. *J Anim Physiol Anim Nutr.* 2005;89:194-8.
299 <https://doi.org/10.1111/j.1439-0396.2005.00557.x>
- 300 19. Hendriks W, Butts C, Thomas D, James K, Morel P, Verstegen M. Nutritional quality and variation of meat
301 and bone meal. *Asian-Australas J Anim Sci.* 2002;15:1507-16.
- 302 20. Stein HH, Sève B, Fuller M, Moughan P, De Lange C. Invited review: Amino acid bioavailability and
303 digestibility in pig feed ingredients: Terminology and application. *J Anim Sci.* 2007;85:172-80.
304 <https://doi.org/10.2527/jas.2005-742>
- 305 21. Sunvold G, Fahey Jr G, Merchen N, Titgemeyer E, Bourquin L, Bauer L, et al. Dietary fiber for dogs: IV. *In*
306 *vitro* fermentation of selected fiber sources by dog fecal inoculum and *in vivo* digestion and metabolism of
307 fiber-supplemented diets. *J Anim Sci.* 1995;73:1099-109. <https://doi.org/10.2527/1995.7341099x>
- 308 22. Swanson KS, Grieshop CM, Flickinger EA, Bauer LL, Healy H-P, Dawson KA, et al. Supplemental
309 fructooligosaccharides and mannanoligosaccharides influence immune function, ileal and total tract nutrient
310 digestibilities, microbial populations and concentrations of protein catabolites in the large bowel of dogs. *J*
311 *Nutr.* 2002;132:980-9. <https://doi.org/10.1093/jn/132.5.980>

- 312 23. Vervaeke I, Decuypere J, Dierick N, Henderickx H. Quantitative *in vitro* evaluation of the energy metabolism
313 influenced by virginiamycin and spiramycin used as growth promoters in pig nutrition. *J Anim Sci.*
314 1979;49:846-56. <https://doi.org/10.2527/jas1979.493846x>
- 315 24. Hervera M, Baucells M, González G, Pérez E, Castrillo C. Prediction of digestible protein content of dry
316 extruded dog foods: comparison of methods. *J Anim Physiol Anim Sci.* 2009;93:366-72.
317 <https://doi.org/10.1111/j.1439-0396.2008.00870.x>
- 318 25. Kuzmuk KN, Swanson KS, Tappenden KA, Schook LB, Fahey Jr GC. Diet and age affect intestinal
319 morphology and large bowel fermentative end-product concentrations in senior and young adult dogs. *J Nutr.*
320 2005;135:1940-5.
- 321 26. Fahey Jr GC, Barry KA, Swanson KS. Age-related changes in nutrient utilization by companion animals.
322 *Annu Rev Nutr.* 2008;28:425-45. <https://doi.org/10.1146/annurev.nutr.28.061807.155325>
- 323 27. Swanson KS, Dowd SE, Suchodolski JS, Middelbos IS, Vester BM, Barry KA, et al. Phylogenetic and gene-
324 centric metagenomics of the canine intestinal microbiome reveals similarities with humans and mice. *ISME J.*
325 2011;5:639-49. <https://doi.org/10.1038/ismej.2010.162>
- 326 28. Case LP, Daristotle L, Hayek MG, Raasch MF. *Canine and feline nutrition: a resource for companion animal*
327 *professionals: Elsevier Health Sciences; 2010.*
- 328 29. Laflamme D. Determining metabolizable energy content in commercial pet foods. *J Anim Physiol Anim Nutr.*
329 2001;85:222-30. <https://doi.org/10.1046/j.1439-0396.2001.00330.x>
- 330 30. Dobenecker B, Endres V, Kienzle E. Energy requirements of puppies of two different breeds for ideal growth
331 from weaning to 28 weeks of age. *J Anim Physiol Anim Nutr.* 2013;97:190-6. [https://doi.org/10.1111/j.1439-](https://doi.org/10.1111/j.1439-0396.2011.01257.x)
332 [0396.2011.01257.x](https://doi.org/10.1111/j.1439-0396.2011.01257.x)
- 333 31. Murray S, Fahey Jr G, Merchen N, Sunvold G, Reinhart G. Evaluation of selected high-starch flours as
334 ingredients in canine diets. *J Anim Sci.* 1999;77:2180-6. <https://doi.org/10.2527/1999.7782180x>
- 335 32. Crane SW, Griffin RW, Messent PR. Introduction to commercial pet foods. *Small animal clinical nutrition.*
336 2000;4:111-26.
- 337 33. Swanson KS, Kuzmuk K, Schook LB, Fahey Jr G. Diet affects nutrient digestibility, hematology, and serum
338 chemistry of senior and weanling dogs. *J Aim Sci.* 2004;82:1713-24. <https://doi.org/10.2527/2004.8261713x>
- 339 34. Zentek J, Marquart B, Pietrzak T. Intestinal effects of mannanoligosaccharides, transgalactooligosaccharides,
340 lactose and lactulose in dogs. *J Nutr.* 2002;132:1682S-4S.
- 341 35. Flickinger E, Schreijen E, Patil A, Hussein H, Grieshop C, Merchen N, et al. Nutrient digestibilities, microbial

342 populations, and protein catabolites as affected by fructan supplementation of dog diets. *J Anim Sci.*
343 2003;81:2008-18. <https://doi.org/10.2527/2003.8182008x>
344 36. Sandøe P, Corr S, Palmer C. *Companion animal ethics*: John Wiley & Sons; 2015.

ACCEPTED

Table 1. Compositions of experimental dog diet

Items	Contents
Ingredient, %	
Oat	35.00
Turkey	30.00
Chicken breast meal	10.00
Chicken liver	8.00
Brown seaweed	4.00
Broccoli powder	3.00
Corn starch	2.00
Flour	2.00
Apple powder	2.00
Monocalcium phosphate	1.00
Calcium carbonate	1.00
Vitamin-mineral premix ¹	0.60
Potassium citrate	0.50
Perilla oil	0.50
Salt	0.40
Total	100.00
Calculated composition	
Dry matter, %	86.73
Crude protein, %	29.07
Ether extract, %	7.52
Crude fiber, %	0.62
Ash, %	4.85
Calcium, %	1.04
Phosphorus, %	0.72
Metabolizable energy ² , kcal/kg	3,684.60

¹ Vitamin and mineral premix supplied per kg of diets: 3,500 IU vitamin A; 250 IU vitamin D₃; 25 mg vitamin E; 0.052 mg vitamin K; 2.8 mg vitamin B₁(thiamine); 2.6 mg vitamin B₂ (riboflavin); 2 mg vitamin B₆ (pyridoxine); 0.014 mg vitamin B₁₂; 6 mg Cal-d-pantothenate; 30 mg niacin; 0.4 mg folic acid; 0.036 mg biotin; 1,000 mg taurine; 44 mg FeSO₄; 3.8 mg MnSO₄; 50 mg ZnSO₄; 7.5 mg CuSO₄; 0.18 mg Na₂SeO₃; 0.9 mg Ca(IO₃)₂.

² Metabolizable energy (ME) was calculated follow equation; ME (kcal/kg) = ([CP × 3.5] + [EE × 8.5] + [NFE × 3.5]) × 10.

Table 2. Comparison of *in vitro* and *in vivo* digestibility in mixed-breed dogs using experimental diet¹

Items (%)	IVTD	IVPD	IVAD	IVSD	Contrasts (<i>p</i> -value)		
					IVTD vs IVPD	IVTD vs IVAD	IVTD vs IVSD
DM	87.42 ± 2.70	78.89 ± 1.09	78.77 ± 1.16	75.34 ± 0.89	0.001	0.001	<0.001
OM	91.33 ± 0.40	82.41 ± 0.91	82.32 ± 0.98	79.54 ± 0.74	<0.001	<0.001	<0.001
CP	89.83 ± 2.13	79.44 ± 1.17	81.73 ± 1.02	79.55 ± 0.62	<0.001	<0.001	<0.001
GE	87.39 ± 2.92	82.09 ± 0.91	83.01 ± 0.96	78.96 ± 0.74	0.074	0.034	0.002
CF	84.87 ± 2.81	70.10 ± 1.98	72.74 ± 1.32	73.52 ± 1.30	<0.001	<0.001	0.001
EE	84.87 ± 3.72	63.58 ± 2.12	66.03 ± 1.28	66.08 ± 1.48	<0.001	<0.001	<0.001

¹Each mean represents 6 observations for *in vivo* and *in vitro*, respectively.

²Abbreviaiton: DM, dry matter; OM, organic matter; CP, crude protein; GE, gross energy; CF, crude fiber; EE, ether extract; IVTD, *in vitro* digestibility; IVPD, *in vivo* digestibility of puppies; IVAD, *in vivo* digestibility of adult dogs; IVSD, *in vivo* digestibility of senior dogs; SE, standard error.

346
347

Table 3. Life-stage comparison of *in vivo* apparent total tract digestibility (ATTD) of an experimental diet in mixed-breed dogs¹

Items (%)	Puppies	Adults	Seniors	SE	<i>p</i> -value
DM	78.89	78.77	75.34	1.05	0.050
OM	82.41	82.32	79.54	0.88	0.060
CP	79.44	81.73	79.55	0.96	0.199
GE	82.09a	83.01a	78.96b	0.87	0.013
CF	70.10	72.74	73.52	1.57	0.299
EE	63.58	66.03	66.08	1.67	0.494

¹Each mean represents 6 dogs per life stage.

²Means within a row with different superscripts differ significantly ($p < 0.05$).

³Abbreviations: DM, dry matter; OM, organic matter; CP, crude protein; GE, gross energy; CF, crude fiber; EE, ether extract; SE, standard error of the mean.

ACCEPTED

Table 4. Linear regression analysis between *in vivo* (y) and *in vitro* digestibility (x) in mixed-breed dogs¹

Items	Equation	r ²	RMSE
Puppies			
DM	y=0.31x+51.54	0.60	1.90
OM	y=1.42x-47.58	0.40	1.94
CP	y=0.20x+70.85	0.55	1.05
GE	y=0.05x+83.94	0.93	0.11
CF	y=0.66x+14.28	0.87	1.96
EE	y=0.54x+17.46	0.91	1.79
Adult dogs			
DM	y=0.35x+47.99	0.67	1.81
OM	y=2.37x-134.51	0.96	0.57
CP	y=0.39x+49.80	0.56	2.00
GE	y=0.13x+77.31	0.98	0.14
CF	y=0.43x+36.00	0.85	1.41
EE	y=0.24x+45.25	0.51	2.46
Senior dogs			
DM	y=0.31x+47.97	0.91	0.73
OM	y=1.59x-65.83	0.76	0.99
CP	y=0.46x+41.20	0.66	1.90
GE	y=0.07x+81.37	0.87	0.23
CF	y=0.42x+37.72	0.82	1.50
EE	y=0.38x+33.65	0.93	1.10

¹ Each mean represents 6 observations for *in vivo* and *in vitro*, respectively.

²Abbreviaiton: DM, dry matter; OM, organic matter; CP, crude protein; GE, gross energy; CF, crude fiber; EE, ether extract; RMSE, root mean squared error