# JAST (Journal of Animal Science and Technology) TITLE PAGE

	Fill in information in each box below
Article Type	Research article
Article Title (within 20 words without abbreviations)	Evaluating the effects of finishing diet and feeding location on
	sheep performance, carcass characteristics, and internal parasites
Running Title (within 10 words)	The effects of finishing diet and feeding location on sheep
Author	Ragen Devon L1, Butler Molly R1, Boles Jane A1, Layton William
	A2, Craig Thomas M3, and Hatfield Patrick G1
Amiliation	I Department of Animal and Range Sciences, Montana State
	2Montana Department of Livestock, Bozeman, MT 59718
	3Department of Veterinary Pathobiology, Texas A&M University,
	College Station, TX, 77843
ORCID (for more information, please visit	Ragen DL (https://orcid.org/0000-0002-0140-1581)
https://oreid.org/	Boles JA (https://orcid.org/0000-0001-8381-3434) Butler MB (https://orcid.org/0000_0000_0000_0000)
	Lavton WA (https://orcid.org/0000-0000-0000-0000)
	Craig TM (https://orcid.org/0000-0000-0000-0000)
	Hatfield PG (https://orcid.org/0000-0000-0000-0000)
Competing interests	No potential conflict of interest relevant to this article was reported.
Funding sources	We would like to thank WSADE for the funding of this project:
State funding sources (grants, funding sources,	"Integrating Livestock into Cropping Systems and the Impacts on
equipment, and supplies). Include name and number of	Soil Health" SW17-080.
grant if available.	
Acknowledgements	The authors would like to acknowledge Ryan Knuth for his
	assistance with the SAS software. We also appreciate the help of
	Chad Page, Olivia Fernandez, and Miriah Butler for field work.
Availability of data and material	Upon reasonable request, the datasets of this study can be available
	from the corresponding author.
Authors' contributions	Conceptualization: Hatfield PG Boles IA
Please specify the authors' role using this form.	Data curation: Ragen DL, Butler MR.
	Formal analysis: Ragen DL, Butler MR.
	Methodology: Ragen DL, Butler MR.
	Investigation: Ragen DL, Butler MR.
	Writing - original draft: Kagen DL, Butler MK.
	Lavton WL, Thomas CM, Ragen DL,
Ethics approval and consent to participate	All animal procedures were approved by the Montana State
	University Agricultural Animal Care and Use Committee (Protocol
	#2013-AA07 approved October 29, 2013)
4	

 $\begin{array}{c}
 1 \\
 2 \\
 3
 \end{array}$ 

### 5 CORRESPONDING AUTHOR CONTACT INFORMATION

For the corresponding author (responsible for correspondence, proofreading, and reprints)	Fill in information in each box below
First name, middle initial, last name	Devon, L, Ragen
Email address – this is where your proofs will be sent	Devon.ragen@montana.edu
Secondary Email address	none
Address	103 Animal Bioscience Bldg, Montana State University, Bozeman, MT 59717
Cell phone number	406-579-1539
Office phone number	406-994-7749
Fax number	406-994-5589

## 8 Abstract

9	A 3 yr experiment was conducted to evaluate the influence of diet and feeding location on animal
10	performance, carcass characteristics, whole blood counts, and internal parasite burden of lambs assigned to
11	1 of 4 treatments: 1) confinement fed 71% alfalfa, 18% barley pellet, 5% molasses, 0.013% Bovatec, 6.1%
12	vitamin/mineral package diet (CALF), 2) confinement fed 60% barley, 26% alfalfa pellet, 4% molasses,
13	2.5% soybean-hi pro, 0.016% Bovatec, 7.4% vitamin/mineral package diet (CBAR), 3) field fed 71%
14	alfalfa, 18% barley pellet, 5% molasses, 0.013% Bovatec, 6.1% vitamin/mineral package diet (FALF), and
15	4) field fed 60% barley, 26% alfalfa pellet, 4% molasses, 2.5% soybean-hi pro, 0.016% Bovatec, 7.4%
16	vitamin/mineral package diet (FBAR). Lambs had <i>ad libitum</i> access to feed and water during the 60-65 d
17	experimental period each year. A year $\times$ location interaction was detected for ending BW, ADG, and DMI;
18	therefore results are presented by year. In all years, cost of gain and DMI were greater for CALF and FALF
19	than for CBAR and FBAR feed treatments ( $p \le 0.03$ ). In yr 2 and 3 field treatments had greater ending BW
20	and ADG than confinement treatments. For all years, diet did not affect ending BW or ADG. In yr 1
21	dressing percent and rib eye area were greater for field finished lambs than confinement finished ( $p \le 0.02$ )
22	and Warner-Bratzler shear force was greater for CALF and FALF ( $p = 0.03$ ). In yr 2 lambs in FALF and
23	FBAR treatments had greater leg scores and conformation than CALF and CBAR ( $p = 0.09$ ). In yr 1, FALF
24	had a greater small intestine total worm count than all other treatments. In yr 1, ending Trichostrongyle
25	type egg counts were greater for FALF (p=0.05). In yr 2, ending Nematodirus spp. egg counts were greater
26	for FALF and lowest for CBAR ( $p < 0.01$ ). Abomasum <i>Teladorsagia circumcinta</i> worm burden was greater
27	in CALF than all other treatments ( $p=0.07$ ) in yr 2. While field finishing lambs with a grain- or forage-
28	based diet we conclude that it is possible to produce a quality lamb product without adverse effects to
29	animal performance, carcass quality or increasing parasite burdens.
30	Keywords: carcass, confinement, field, finishing, parasites, sheep.

33	Introduction
34	Integrated crop and livestock systems as an alternative to confinement feedlot operations have
35	gained popularity in recent years. The excretion and concentration of manure at concentrated animal
36	feeding operations has resulted in environmental concerns associated with nuisance issues (odors, disease
37	vectors), water quality, and air quality [1]. Conversely, the application of manure to soil provides potential
38	benefits including improving the fertility, structure, water holding capacity of soil, increasing soil organic
39	matter and reducing the amount of synthetic fertilizer needed for crop production [2-4]. Incorporating
40	livestock into cropping systems may offer alternative uses of crops and provide new grazing opportunities
41	for livestock producers [5]. It also offers an alternative to traditional stubble management, and enables
42	improved pest management [6-8]. In Montana, sheep producers commonly manage their flock on
43	rangeland or pasture during the grazing months of spring and summer and feed harvested forages during
44	the fall and winter. Lambs raised for meat production are typically finished in a confinement feedlot
45	operation [9].
46	Internal parasitism of grazing livestock is a significant world-wide disease problem [10, 11].
47	Gastrointestinal parasites (GIP) in the abomasum and small intestine cause extensive protein loss in the
48	digestive tracts of sheep [12] and failure to control GIP results in poor animal growth rates and thrift, and
49	can result in animal mortality [13]. Previous research has reported that gastrointestinal nematode eggs per
50	gram (EPG) were lowest in lambs fed in confinement, moderate in semi-confinement, and highest in
51	grazing lambs [14]. Research by Ebrahim [15] revealed that blood samples taken from sheep infested with
52	gastrointestinal parasites had a significant decrease in hemoglobin, packed cell volume and red blood cell
53	count values, while white blood cell count was significantly increased, compared to blood samples from a
54	group of sheep that were parasite-free.
55	Previous research has only focused on grazed or harvested forage as the diet for finishing livestock
56	and there is an increasing demand for non-confinement, forage-fed animals which are stimulating market
57	interest in alternative production systems [16, 17]. Providing a finishing ration to livestock on cropland is
58	an unconventional way to finish livestock while adding manure to the soil. The objectives of this 3 yr

59	study were to compare the effects of finishing diet and location on the performance, carcass characteristics,
60	whole blood counts, and parasite loads of weaned, crossbred lambs.
61	
62	<b>Materials and Methods</b>
63	Ethics Statement
64	All animal procedures were approved by the Montana State University Agricultural Animal Care
65	and Use Committee (Protocol #2013-AA07 approved October 29, 2013).
66	Sampling
67	Ninety crossbred lambs (ewes and wethers; Blackface $\times$ Western whiteface; 6-mo-old; BW =
68	$35.6\pm5.7$ kg) and 18 Targhee lambs (ewes; 6-mo-old; BW = $35.8\pm5.2$ kg) were used in yr 1 (September 25
69	to November 25) of the study. Targhee ewe lambs were used because of an insufficient number of
70	crossbred lambs in yr 1. The Targhee ewes were evenly distributed through all treatments. One hundred
71	and eight crossbred lambs (ewes and wethers; Blackface $\times$ Western whiteface; 5-mo-old; BW = 31.8 $\pm$ 4.7
72	kg) were used in yr 2 (September 3 to November 3) of the study. One hundred and eight crossbred lambs
73	(ewes and wethers; Blackface $\times$ Western whiteface; 5-mo-old; BW = 36.8 $\pm$ 5.7 kg) were used in yr 3
74	(September 1 to November 5) of the study. For yr 1, yr 2 and yr 3, lambs were transported to the Fort Ellis
75	Research Facility (45°40'N, 111° 2'W, altitude 1468 m) in Bozeman, Montana, USA on September 19th,
76	September 1 <sup>st</sup> , and August 30 <sup>th</sup> , respectively. Mean monthly air temperature at the site ranges from -5.6 °C
77	in January to 19 °C in July and the total annual precipitation (113-yr average) is 465 mm. On d 0 all lambs
78	were placed in a dry lot pen and held off of feed and water overnight. On d 1 lambs were weighed, and
79	paint-branded for identification purposes. Lambs were then stratified by BW and sex and allocated to
80	treatments. In all years, the four treatments consisted of: 1) confinement fed 71% alfalfa, 18% barley pellet
81	diet (CALF), 2) confinement fed 60% barley, 26% alfalfa pellet diet (CBAR), 3) field fed 71% alfalfa,
82	18% barley pellet diet (FALF), and 4) field fed 60% barley, 26% alfalfa pellet diet (FBAR)(Table 1).
83	For all three years of the study, pen was the experimental unit with nine lambs per pen in
84	confinement (4 pens total). Field was the experimental unit with six fields per treatment (12 fields total),

with six lambs per field. Harvest occurred at the end of the 60 d finishing period for yr 1 and yr 2; carcass
data was not collected in yr 3 as lambs were not harvested the third year of the study.

87 All lambs on CBAR and FBAR treatment diets had a step-up period of 2 weeks, during which a 88 combination of the barley and alfalfa pellets were fed to lambs in 2 to 3 day increments (25-75, 35-65, 45-89 55, 55-45, 65-35, 75-25, 85-15, and 100-0 percent respectively). Lambs being field finished, both FALF 90 and FBAR, were on organic wheat stubble fields measuring 15.2 m by 44.2 m and fed from self-feeders. 91 Self-feeders were moved to a different area of the field at ~30 d to help distribute manure and urine. The 92 winter wheat stubble fields used for this experiment were part of a common 5-yr crop rotation planted at 93 the site: 1) safflower under-sown to biennial sweet clover, 2) sweet clover cover crop/green manure, 3) 94 winter wheat, 4) lentils, and 5) winter wheat. Further details of the farm management practices used at the 95 research site can be found in [18-21]. 96 Confinement finishing took place in dry lot pens and used a GrowSafe feed intake system 97 (GrowSafe Systems, LDT., Airdrie, AB, Canada) to measure intake. GrowSafe is a feed intake acquisition 98 technology using an electronic radio frequency identification (RFID) system that has enabled researchers 99 and producers to monitor individual animal intake and behavior to more precisely evaluate feed efficiency 100 and health status [22]. GrowSafe records feeding behavior traits such as total intake, frequency and 101 duration of feeding, and eating rate for each individual animal through the use of RFID tags that provide a 102 continuous transmission of data to a computer located at the facility. The lambs in each confinement 103 treatment, CALF and CBAR, were housed together in an 11.3 by 7.1 m pen with one GrowSafe bunk; 104 modifications were made to the beef cattle stanchions through elevated platforms for sheep. 105 Prior to all weigh days, animals were kept in a separate dry lot pen where feed and water were 106 withheld overnight for approximately 16 h. Each lamb's BW was recorded at start and at the end of the trial. 107 Average daily gain (ADG) was calculated for the entire ~2 mo trial period for each year. Daily dry matter 108 intake (DMI) was determined between days that BW was recorded for all lambs and the average ratio of 109 gain to feed (G:F) was calculated between weigh days by dividing ADG by average daily DMI. Cost of 110 gain was calculated by multiplying G:F by cost per kg of feed. Feed cost was determined by the purchase 111 price of the treatment diets which was based on current market value (Table 1). The higher price of feed in

112 yr 1 was due to a persistent drought that increased the demand for hay as well as lingering winter

113 temperatures that wilted spring crops.

#### 114 Carcass data

Carcass data was only collected for yr 1 and yr 2. At the end of yr 1 and yr 2, 32 wether lambs (8 from each treatment) were transported to Pioneer Meats in Big Timber, MT where harvest occurred the following day, using standard industry practices for a small packing plant. After harvest, carcasses were hung by the Achilles tendon, a hot carcass weight was recorded, and dressing percentage was measured. Carcasses were chilled for 24 h at 4°C and then transported to the Montana State University Meat Lab for further processing and data collection. An experienced evaluator obtained individual-level carcass data. These measurements included

122 back fat depth, rib-eye area, leg score, conformation, flank streaking, and quality grade. Additionally, a 123 sample of four rib chops were obtained from the posterior portion of the left longissimus thoracis, vacuum-124 packaged and frozen at -20°C for later tenderness analysis. Warner-Bratzler shear force (WBSF) was 125 determined on cooked and chilled rib chops. Samples were defrosted for 24 h at 4°C, dried, weighed, and 126 broiled in an electric oven until an internal temperature of 35 °C was reached. Temperature was monitored 127 using a Digi Sense Scanning Thermometer from Cole Palmer (625 East Bunker Court Vernon Hills, IL 128 60061 USA) fitted with copper constantan needle thermocouples (Omega Engineering) placed in the 129 geometric center of the chop. All chops were turned over when the temperature reached 35 °C and 130 continued cooking until a final internal temperature of  $71\pm1$  °C was reached at which point chops were 131 removed from the heat source. Samples were bagged and allowed to chill at 4°C for at least 1 h. Samples 132 were then reweighed to determine cook loss, and two circular cores measuring 1.27cm in diameter were 133 obtained parallel to the fiber direction using a hand-held coring device (cork borer) with a total of 6 cores 134 collected from 3 chops. Cores were allowed to reach room temperature prior to being sheared perpendicular 135 to the fiber direction using a TMS-90 Texture System (Food Technology Corp., Rockville, MD) fitted with 136 a Warner-Bratzler shear attachment (crosshead speed 200mm//min). The average of the maximum force 137 necessary to shear all cores per carcass was used for statistical analysis.

138 Blood Collection

139 Blood samples were collected into serum and EDTA vacutainers from the jugular vein of lambs on

140 d 0 and d 60 for yr 1 and yr 2 of the study. Immediately after collection, blood was put on ice and

141 transported to the Montana Veterinary Diagnostic Lab for a complete blood count (CBC). The blood count

142 was performed on a CELL-DYN 3700 System (Abbott Laboratories, Abbott Park, IL).

- 143 Internal Parasites
- 144 To evaluate which species, and the amounts of parasites that occurred in the lambs, a fecal egg

count (FEC) was performed on rectal grab samples using the Modified McMaster's technique [23].

146 Samples were collected on days 0 and 60 in both yr 1 and yr 2. Trichostrongyle type or *Nematodirus* were

147 identified and counted individually. One egg inside the grid of the slide represented 50 eggs per gram of

148 fresh feces.

149 During harvest, at the end of yr 1 and yr 2, the abomasum and first meter of the small intestine 150 were collected at random from 4 lambs from each of the four treatments to be used for a total worm count 151 analysis using the modified methods of Wood et al. [24]. The two organs and their contents were separated 152 and washed thoroughly with tap water in a 20 liter bucket. The water level was brought to 10 liters and both 153 the water and organs were left to settle for  $\geq 15$  minutes. Next, the top one half to two thirds of the water 154 was decanted and the process was repeated twice, returning the water level to 10 liters each time. After the 155 third wash, the organ was rinsed again and disposed of and the water level was then returned to 10 liters for 156 a final settling. After all washings had occurred and the fourth wash was decanted, the water level was 157 returned to 10 liters, then mixed with a stir rod and 4-100 mL aliquots were taken and stored in jars with 158 100 mL of 10% formalin for later identification of adult worms in the gastrointestinal tract. The worms 159 were plated onto slides with lactophenol and placed under a compound microscope for identification of 160 species, via anatomical structure, and total number of each parasite was recorded.

161 Feed Analysis

162 The feed was sampled weekly by grab sampling, composited for the entire experiment, and stored 163 in sealed sample bags for analysis at a later date. All feed samples for all years, including pelleted rations 164 and wheat stubble samples, were sent to Midwest Laboratories for nutrient analysis. Moisture, crude 165 protein, acid detergent fiber (ADF), neutral detergent fiber (NDF), total digestible nutrients (TDN), net 166 energy (Gain, Lactation, Maintenance), and relative feed value (RVF) were estimated using the F10:

- 167 Relative Feed Value package (Table 1).
- 168 In addition to the analysis by Midwest Laboratories, all samples were analyzed for organic matter. 169 Two 2-g samples of the 1-mm ground forage were weighed and placed in a muffle oven at 550°C for 15 h 170 to determine ash weights for calculation of OM content [25].

171 **Data Analyses** 

- 172 The study design was a two by two factorial. The model included the effect of year, feeding 173 location, and type of feed. Response variables for performance were: final lamb BW, ADG, feed efficiency, 174 DMI, and cost per kg of gain. Response variables for carcass quality characteristics were: dressing 175 percentage, hot carcass weight, back fat, rib eye area, leg score, conformation, flank streaking, quality 176 grades, and WBSF. A year  $\times$  feed  $\times$  location interaction was detected for dressing percentage, abomasum T. 177 circumcinta counts, small intestine Nematodirus counts, and total small intestine counts. Year × feed 178 interactions were detected for ending white blood cell counts and abomasumal T. circumcinta counts. Year  $\times$  location interactions were detected for DMI, ending BW, ADG, dressing percentage, and abomasum H. 179 180 contortus counts. The data was then analyzed within year using the GLM procedure of SAS (SAS Inst., Inc., 181 Cary, NC). Beginning lamb BW was used as a covariate in the analysis of final BW and ADG. Treatment 182 means were compared using the LSMeans procedure when a significant p value was found ( $p \le 0.10$ ). 183 Fecal egg counts were transformed to the logarithmic base 10 scale prior to analysis. The GLM 184 procedure was run with beginning count as a covariate but it added nothing to the model and was removed. 185 Total worm counts were also logarithmic base 10 transformed and Proc GLM was used for analysis. 186
- 187

### **Results and Discussion**

- 188 **Performance**
- 189 In North America, confinement-finished lamb-meat production promotes rapid growth and is 190 based on diets containing high levels of grain concentrates. The majority of research has reported that 191 lambs grow faster on concentrate-based diets than on forage-based diets [26-35] and ad libitum

192 consumption of concentrates results in fatter lambs compared to those fed forage diets when the lambs are

- 193 slaughtered at a constant final weight [27, 36, 37]. However, there is a rapidly increasing demand for grass-
- 194 fed or organically produced livestock [38] and in the U.S. retail sales of pasture-finished beef have risen
- 195 from \$17 million in 2012 to \$272 million in 2016 [39].
- 196 A year  $\times$  location interaction was detected for ending BW, ADG, and DMI; therefore results are
- presented by year. In yr 1, there were no interactions between location and feed (p > 0.29) for all response
- 198 variables (Table 2). There was also no effect of location (p > 0.42) on ending BW, ADG, DMI, gain to feed,
- and cost of gain. Cost of gain and DMI were greater for CALF and FALF than for CBAR and FBAR feed
- 200 treatments (p < 0.01) across all years. Gain to feed ratio was greater for CBAR and FBAR than CALF and
- 201 FALF (0.15, 0.14, 0.12, and 0.13 G:F respectively; *p* = 0.01).
- In yr 2, there were no interactions between location and feed (p > 0.32) for all response variables (Table 3). Location had an effect on ending BW and ADG, and both FALF and FBAR were greater than CALF and CBAR (p < 0.01). Dry matter intake, gain to feed ratio, and cost of gain had both a location and feed effect and differed among all treatments (p < 0.01; Table 3).
- 206 In yr 3, there was a location  $\times$  feed interaction for ending BW and ADG (p=0.09 and p=0.08, 207 respectively) (Table 4). Ending BW and ADG were greater for both FALF and FBAR than CALF and 208 CBAR (p < 0.01). Feed had an effect on DMI and was greater for FALF and CALF which were similar (p =209 0.03). This difference in DMI was expected as it has been reported that the decrease in DMI as a result of 210 feeding higher proportion of concentrate can be attributed to the regulatory effect of dietary energy intake. 211 Generally, animals eat food mainly to satisfy their desire for energy [40]. Gain to feed did not differ among 212 treatments (p=0.53). Cost of gain had both a location and feed effect and was highest for CALF and FALF 213 (\$4.15/kg and \$3.72/kg, respectively; p < 0.01) than CBAR and FBAR with FBAR having the lowest cost 214 of gain ( $\frac{2.75}{\text{kg}}$ ; p < 0.01). Even though the cost of gain was greater for alfalfa fed lambs, a survey 215 conducted by Ripoll et al. [41] reported that 70.4% of consumers surveyed believe that grass-fed lamb is 216 better and may be willing to pay a premium price for what they perceive as a "higher-quality product". 217 Also, Jacques et al. [42] concluded that using forage finishing systems may improve processing efficiency

by reducing the amount of external fat to be removed by preventing excessively fat carcasses from lambs slaughtered. However, in our study there was no difference in back fat thickness among treatments ( $p \le 0.33$ ; Tables 5 and 6).

In yr 2 and 3, both field treatments had greater ending BW and ADG than the confinement treatments. In yr 1, extreme weather conditions (temperatures down to -29°C for ~ 1 week) may have affected animals in the field and had an impact on animal performance. Our results are in agreement with those of Phillips et al. [43] who reported that lambs can be adequately finished on a forage-based diet (alfalfa or kenaf) and doing so does not adversely affect performance or feed intake. McClure et al. [32] and Aurosseau et al. [44] also determined that finishing lambs on high-quality forages can yield similar ADG to those achieved in confinement feeding a concentrate diet while producing comparable carcasses.

228 Carcass

229 A year  $\times$  location and a year  $\times$  location  $\times$  feed interaction was detected for dressing percentage; 230 therefore, results are presented by year. In yr 1, there was no interaction between location and feed (p>231 0.09) for all response variables (Table 5). Dressing percent differed by location but not feed; dressing 232 percent was greater for FALF and FBAR than for CALF and CBAR (51.92, 52.85, 49.18, and 46.45% 233 respectively; p < 0.02). Rib eye area differed by location but not feed; rib eye area was greater for FALF 234 and FBAR than for CALF and CBAR (6.03, 6.32, 5.30, and 5.33 cm<sup>2</sup> respectively; p < 0.02; Table 5). 235 WBSF was greater for CALF and FALF than CBAR and FBAR (3.8, 3.8, 2.7, and 3.1kg respectively; p =236 0.01). All other carcass measurements did not differ among treatments. Nichols et al. [45] reported that 237 lambs overwintered on stubble fields graded choice after confinement feeding; however, they did not 238 investigate alternatives to confinement feeding. There was no difference in quality grade amongst 239 treatments or years; all treatments graded in the good-plus to choice-minus range ( $p \le 0.77$ ; Table 5). 240 In vr 2, there were no interactions between location and feed (p > 0.23) for all variables. Lambs in 241 FALF and FBAR treatments tended to have greater leg scores and conformation than CALF and CBAR (p=242 0.09). All other carcass measurements did not differ among treatments (Table 6). In our study we 243 observed lambs in the field treatments exercising more than lambs in confinement pens; further research

should be conducted to corroborate these observations and determine if exercise has an effect on leg scores

and conformation. Our results are in conflict with the results of Jones et al. [46], where confinement fed

246 lambs produced heavier carcasses and larger rib eye areas than pasture fed lambs. Our results are also in

- 247 disagreement with Purchas et al. [47] who reported that WBSF values were significantly lower for *M*.
- 248 seminembranosus in the pasture vs. grain treatments of 50 kg harvest weight lambs (4.04, 4.67 kg,
- 249 respectively). In our study, location did not have an effect on tenderness but WBSF was greater for CALF
- and FALF than CBAR and FBAR (3.8, 3.8, 2.7, and 3.1 respectively; p = 0.01). Both Duckett et al. [48] and
- 251 Realini et al. [49] reported that WBSF values were similar between forage and concentrate-fed animals. In
- 252 our trial the forage-based treatment diet and grain-based treatment diet were both in concentrate-form with
- the same particle size. Past studies have only investigated forage-fed (in pasture or hay form) vs.
- 254 concentrate-fed diets and therefore may not be comparable to our study. Forage particle size influences
- 255 feed intake, saliva production, rumination, and the passage rate of feed in the rumen, as well as bio-
- 256 hydrogenation pathways and fatty acid composition in lamb meat [50].
- 257 Blood

258 A year  $\times$  feed interaction was detected for ending white blood cell counts; therefore, results are 259 presented by year. In yr 1, there were no interactions between location and feed (p > 0.12) for all variables 260 (Table 7). Both hematocrit and mean cell hemoglobin concentration were affected by feed and were greater 261 for CALF and FALF than CBAR and FBAR (p < 0.08). This agrees with the results of Gawel and Grzelak 262 [51] who reported that alfalfa concentrate may be important as a dietary supplement for animals and may 263 improve the hematological indices of blood. Mean cell hemoglobin and red cell distribution width differed 264 by location; FALF and FBAR were both greater than CALF and CBAR (p < 0.06). All other blood counts 265 did not differ between treatments. The principal clinical sign of Haemonchus contortus (H. contortus) 266 infections is anemia, due to the blood-letting activities of the parasite [52]. The cool season parasite T. 267 *circumcinta* (formerly *Ostertagia circumcinta*) interferes with absorption of nutrients and may cause 268 weight loss and possibly diarrhea [53].

269

In yr 2, there were no interactions between location and feed (p > 0.38) for all variables (Table 8).

270 Red blood cell counts were affected by feed and were greater for CALF and FALF than CBAR and FBAR 271 (p<0.07). The tendency for CALF to have greater white blood cell counts in yr 2, compared to other 272 treatments, may be influenced by its abomasum T. circumcinta worm burden which was greater in CALF 273 than all other treatments (p=0.07). This would agree with the results of Ebrahim [15] who reported that 274 blood samples taken from sheep infested with gastrointestinal parasites had greater white blood cell counts 275 than sheep that were parasite-free. Kowalczuk-Vasilev et al. [54] reported that the use of iron-rich alfalfa 276 concentrate in feeding lambs significantly improved hematological blood indices: hematocrit, hemoglobin 277 and erythrocytes (RBC), and may be due to more efficient iron absorption. Hematocrit and mean cell 278 hemoglobin differed by location and FALF and FBAR were greater than CALF and CBAR (p < 0.08). 279 Variation in factors that affect the rumen bacterial community (diet composition, feed types, feeding 280 strategy) can have a robust effect on rumen metabolism, which can impact both productivity and health 281 of ruminants [55]. Hemoglobin and mean cell hemoglobin concentration had an effect of both location and 282 feed therefore they differed between all treatments (p < 0.06). All other blood counts did not differ between 283 treatments (Table 8).

#### 284 Parasites

285 A year  $\times$  location and year  $\times$  feed interaction was detected for *H. contortus* worm counts along 286 with a year  $\times$  feed  $\times$  location interaction for *T. circumcinta* worm counts, *Nematodirus* worm counts and 287 small intestine total worm counts, therefore, results are presented by year (p < 0.01). In yr 1, there were no 288 interactions between location and feed (p > 0.72) for all variables. Ending *Trichostrongyle* egg counts 289 differed at p < 0.05. Ending *Nematodirus* spp. egg counts did not differ between treatments, although there 290 was a tendency (p = 0.11) for CALF and FALF to be greater than CBAR and FBAR (Table 9). 291 In yr 1, there was an interaction between location and feed for small intestine total worm count 292 (p < 0.01). FALF had a greater small intestine total worm count than all other treatments, CBAR was 293 intermediate, and CALF and FBAR had the lowest counts (176, 18, 2, 0 small intestine worm count;

- 294 p=0.01). An interaction between location and feed was also present for *Nematodirus* worm counts (p < 0.01).
- FALF was greater (p < 0.03) than all other treatments, CBAR was intermediate and differed from all other

treatments (p=0.03) and FBAR and CALF were the lowest (p=1.00). All other worm burdens in the

abomasum and first meter of the small intestine did not differ among treatments (Table 10).

- 298 In vr 2, there were no interactions between location and feed (p > 0.24) for all parasite variables. 299 Ending *Trichostrongyle* type egg counts did not differ between all treatments (Table 11). Ending 300 Nematodirus spp. egg counts were affected by location and feed and were greater for FALF and CALF 301 (20.62 and 9.99 EPG respectively; p > 0.30), however CALF did not differ from FBAR (5.60 EPG; p > 0.44), 302 and CBAR was lowest and differed from all other treatments (0.52 EPG; p < 0.03). It is unknown why 303 CBAR had the lowest EPG in yr 2 but there was a tendency for both barley treatment groups to have lower 304 EPG than the field treatments. It is widely accepted that a high grain diet causes a drop in ruminal pH and 305 may cause drastic shifts in the rumen microbial community [56-59]. Ruminal bacterial and protozoal 306 populations increase or decrease in response to pH changes [57], however, the effect of pH on internal 307 parasites has not been studied. It is possible that a lamb's rumen environment is less favorable to internal 308 parasites while consuming a high grain (barley) diet and thus we see lower EPG in these lambs; more 309 research is needed to investigate this relationship. 310 In yr 2, there were no interactions (p > 0.11) between location and feed for all parasite variables.
- Abomasum *H. contortus* worm burden was greater in CALF than all other treatments (p= 0.07). All other worm burdens in the abomasum and first meter of the small intestine did not differ among treatments (Table 12). Marley et al. [60] determined that legume forages have the potential to contribute to the control of abomasal but not small intestine nematode parasites in finishing lamb systems. This is in contrast to our results where alfalfa-fed lambs in confinement had greater abomasum *T. circumcinta* worm counts than other treatments.
- 317 Parasite results are in conflict with those of Cai and Bai [14] who reported that gastrointestinal 318 nematode eggs per gram (EPG) were lowest in lambs fed in confinement and highest in grazing lambs. 319 Fecal egg counts (FEC) in our study appeared to trend with barley fed animals having lower counts than 320 those of alfalfa fed animals. Studies have shown that the degree of parasite infestation in sheep may be 321 reduced by some plant species [61-63]. Research has focused on the effects of secondary plant compounds

	322	(e.g. condensed tannins) [64] on the reduction of parasites in the gut but the underlying mechanisms for
	323	such effects have not been determined. Overall in our study FEC worm counts taken from all slaughtered
	324	lambs were low and likely did not adversely influence the weight gains and hematocrit levels of the lambs.
	325	Integrated crop and livestock systems as an alternative to confinement feedlot operations may
	326	increase marketing opportunities for sheep producers. While field finishing lambs with a grain- or forage-
	327	based diet, we conclude that it is possible to produce a quality lamb product without adverse effects to
	328	animal performance, carcass quality or increasing parasite burdens.
329		
330		<b>Competing Interests</b>
331		No potential conflict of interest relevant to this article was reported.
332		Acknowledgements
333		The authors would like to acknowledge Ryan Knuth for his assistance with the SAS software. We also
334	appreci	ate the help of Chad Page, Olivia Fernandez, and Miriah Butler with field work.
335		
336		References
337		
338 339	1. Anim S	Meyer D, Mullinax DD. Livestock nutrient management concerns: Regulatory and legislative overview. J Sci. 1999;77:51-62. 10.2527/1999.77suppl_251x
340 341	2. the yiel	Blay ET, Danquah EY, Ofosu-Anim J, Ntumy JK. Effect of poultry manure and/or inorganic fertilizer on d of shallot [allium cepa var. Aggregatum (g. Don)]. Adv Hort Sci. 2002;16:13-6.
342 343	3. produc	Ofosu-Anim J, Leitch M. Relative efficacy of organic manures in spring barley (hordeum vulgare l.) tion. Aust J Crop Sci. 2009;3.
344 345	4. phosph	Phan TC RM, Cong SS and Nguyen Q, editor Beneficial effects of organic amendment on improving orus availability and decreasing aluminum toxicity in two upland soils. World Cong Soil Sci 2002; Thailand.
346 347	5. impact	Snyder EE, Goosey HB, Hatfield PG, Lenssen AW, editors. Sheep grazing wheat summer fallow and the on soil nitrogen, moisture, and crop yield. Proc West Sec Amer Soc Anim Sci; 2007.
348 349 350	6. densitie 610.	Goosey HB, Hatfield PG, Blodgett SL, Cash SD. Evaluation of alfalfa weevil (coleoptera: Curculionidae) es and regrowth characteristics of alfalfa grazed by sheep in winter and spring. J Entomol Sci. 2004;39:598-

351 7. Goosey HB, Hatfield PG, Lenssen AW, Blodgett SL, Kott RW. The potential role of sheep in dryland
 352 grain production systems. Ag Eco Environ. 2005;111:349-53. <u>http://dx.doi.org/10.1016/j.agee.2005.06.003</u>

8. Hatfield PG, Blodgett SL, Spezzano TM, Goosey HB, Lenssen AW, Kott RW, et al. Incorporating sheep
into dryland grain production systems: I. Impact on over-wintering larva populations of wheat stem sawfly, cephus
cinctus norton (hymenoptera: Cephidae). Small Rumin Res. 2007;67:209-15.

356 <u>http://dx.doi.org/10.1016/j.smallrumres.2005.10.002</u>

357 9. USDA. Lamb from farm to table 2013 [2021 Jan 22]. Available from:

358 <u>https://www.fsis.usda.gov/wps/portal/fsis/topics/food-safety-education/get-answers/food-safety-fact-sheets/meat-</u>
 359 <u>preparation/focus-on-lambfrom-farm-to-table/ct\_index.</u>

360 10. Min BR, Hart SP. Tannins for suppression of internal parasites. J Anim Sci. 2003;81:E102-E9.

Waller PJ. Sustainable nematode parasite control strategies for ruminant livestock by grazing
 management and biological control. Anim Feed Sci Tech. 2006;126:277-89. 10.1016/j.anifeedsci.2005.08.007

Kimambo AE, MacRae JC, Walker A, Watt CF, Coop RL. Effect of prolonged subclinical infection with
 trichostrongylus colubriformis on the performance and nitrogen metabolism of growing lambs. Vet Parasitol.
 1988;28:191-203. <u>http://dx.doi.org/10.1016/0304-4017(88)90107-0</u>

Min BR, Pomroy WE, Hart SP, Sahlu T. The effect of short-term consumption of a forage containing
 condensed tannins on gastro-intestinal nematode parasite infections in grazing wether goats. Small Rumin Res.
 2004;51:279-83. <u>http://dx.doi.org/10.1016/S0921-4488(03)00204-9</u>

Cai KZ, Bai JL. Infection intensity of gastrointestinal nematodosis and coccidiosis of sheep raised under
 three types of feeding and management regims in ningxia hui autonomous region, china. Small Rumin Res.
 2009;85:111-5. <u>https://doi.org/10.1016/j.smallrumres.2009.07.013</u>

Ebrahim ZK. Effect of gastrointestinal parasites infestation on some hematological and biochemical
 parameters in sheep. Alex J Vet Sci. 2018;59:44-7.

Blackburn HD, Snowder GD, Glimp H. Simulation of lean lamb production systems. J Anim Sci.
 1991;69:115-24. /1991.691115x

Font i Furnols M, Realini C, Montossi F, Sañudo C, Campo MM, Oliver MA, et al. Consumer's
purchasing intention for lamb meat affected by country of origin, feeding system and meat price: A conjoint study in
spain, france and united kingdom. Food Qual Pref. 2011;22:443-51.
http://dx.doi.org/10.1016/j.foodcual.2011.02.007

- 379 <u>http://dx.doi.org/10.1016/j.foodqual.2011.02.007</u>
- Barroso J, Miller ZJ, Lehnhoff EA, Hatfield PG, Menalled FD. Impacts of cropping system and
   management practices on the assembly of weed communities. Weed Res. 2015;55:426-35.
   https://doi.org/10.1111/wre.12155

Johnson S, editor Effects of organic and conventional cropping systems on plant diversity and plant soil
 feedbacks 2015.

Lehnhoff E, Miller Z, Miller P, Johnson S, Scott T, Hatfield P, et al. Organic agriculture and the quest for
 the holy grail in water-limited ecosystems: Managing weeds and reducing tillage intensity. Agric. 2017;7:33.

Ragen DL BM, Miller PR, Yeoman CJ, Meccage EC, Weeding JL, and Hatfield Patrick G. Soil
 microbial, chemical and physical properties in tilled organic, conventional no-till, and organic integrated crop–
 livestock systems. In: University MS, editor. 2020.

Mendes ED, Carstens GE, Tedeschi LO, Pinchak WE, Friend TH. Validation of a system for monitoring
 feeding behavior in beef cattle. J Anim Sci. 2011;89:2904-10. 10.2527/jas.2010-3489

- Whitlock HV. Some modifications of the mcmaster helminth egg-counting technique and apparatus. J
   Counc Sci Indust Res. 1948;21:177-80.
- Wood IB, Amaral NK, Bairden K, Duncan JL, Kassai T, Malone JB, Jr., et al. World association for the
  advancement of veterinary parasitology (w.A.A.V.P.) second edition of guidelines for evaluating the efficacy of
  anthelmintics in ruminants (bovine, ovine, caprine). Vet Parasitol. 1995;58:181-213. 10.1016/0304-4017(95)008062
- 398 25. Latimer G. Official methods of analysis of aoac. 19th ed. Arlington, VA: AOAC Intl; 2012.
- 399 26. Taylor MA. Parasite control in sheep: A risky business. Small Rumin Res. 2013;110:88-92.
  400 10.1016/j.smallrumres.2012.11.010
- 401 27. Archimède H, Pellonde P, Despois P, Etienne T, Alexandre G. Growth performances and carcass traits of
   402 ovin martinik lambs fed various ratios of tropical forage to concentrate under intensive conditions. Small Rumin Res.
   403 2008;75:162-70. <u>http://dx.doi.org/10.1016/j.smallrumres.2007.10.001</u>
- 404 28. Borton RJ, Loerch SC, McClure KE, Wulf DM. Characteristics of lambs fed concentrates or grazed on
  405 ryegrass to traditional or heavy slaughter weights. Ii. Wholesale cuts and tissue accretion1. J Anim Sci.
  406 2005;83:1345-52. /2005.8361345x
- 40729.Demirel G, Ozpinar H, Nazli B, Keser O. Fatty acids of lamb meat from two breeds fed different forage:408Concentrate ratio. Meat Sci. 2006;72:229-35. <a href="http://dx.doi.org/10.1016/j.meatsci.2005.07.006">http://dx.doi.org/10.1016/j.meatsci.2005.07.006</a>
- 409 30. Fimbres H, Hernández-Vidal G, Picón-Rubio JF, Kawas JR, Lu CD. Productive performance and carcass
  410 characteristics of lambs fed finishing ration containing various forage levels. Small Rum Res. 2002;43:283-8.
  411 <u>http://dx.doi.org/10.1016/S0921-4488(02)00014-7</u>
- 412 31. McClure KE, Solomont MB, Loerch SC. Body weight and tissue gain in lambs fed an all-concentrate diet
  413 and implanted with trenbolone acetate or grazed on alfalfa. J Anim Sci. 2000;78:1117-24. /2000.7851117x
- 414 32. McClure KE, Van Keuren RW, Althouse PG. Performance and carcass characteristics of weaned lambs
  415 either grazed on orchardgrass, ryegrass, or alfalfa or fed all-concentrate diets in drylot. J Anim Sci. 1994;72:3230-7.
  416 /1994.72123230x
- 41733.Murphy TA, Loerch SC, McClure KE, Solomon MB. Effects of grain or pasture finishing systems on<br/>carcass composition and tissue accretion rates of lambs. J Anim Sci. 1994;72:3138-44. /1994.72123138x
- 419 34. Turner KE, McClure KE, Weiss WP, Borton RJ, Foster JG. Alpha-tocopherol concentrations and case
  420 life of lamb muscle as influenced by concentrate or pasture finishing1. J Anim Sci. 2002;80:2513-21.
  421 /2002.80102513x
- 422 35. Notter DR, Kelly RF, McClaugherty FS. Effects of ewe breed and management system on efficiency of
  423 lamb production: Ii. Lamb growth, survival and carcass characteristics. J Anim Sci. 1991;69:22-33. /1991.69122x
- 424 36. Fisher AV, Enser M, Richardson RI, Wood JD, Nute GR, Kurt E, et al. Fatty acid composition and eating
  425 quality of lamb types derived from four diverse breed × production systems. Meat Sci. 2000;55:141-7.
  426 <u>http://dx.doi.org/10.1016/S0309-1740(99)00136-9</u>
- 427 37. Resconi VC, Campo MM, Furnols MFi, Montossi F, Sañudo C. Sensory evaluation of castrated lambs
  428 finished on different proportions of pasture and concentrate feeding systems. Meat Sci. 2009;83:31-7.
  429 <u>http://dx.doi.org/10.1016/j.meatsci.2009.03.004</u>
- 430 38. Catherine Greene, Kremen A. U.S. Organic farming in 2000-2001: Adoption of certified systems. USDA
  431 Economic Research Service; 2003. Report No.: AIB-780.

- 432 39. Cheung R, P M. Back to grass: The market potential for u.S. Grassfed beef. Stone Barns Center for Food
   433 and Agriculture [Internet]. 2017.
- 434 40. Van Soest PJ, Ferreira AM, Hartley RD. Chemical properties of fibre in relation to nutritive quality of 435 ammonia-treated forages. Anim Feed Sci Tech. 1984;10:155-64. <u>https://doi.org/10.1016/0377-8401(84)90005-1</u>
- 436 41. Ripoll G, Joy M, Panea B. Consumer perception of the quality of lamb and lamb confit. Foods.
  437 2018;7:80.
- 42. Jacques J, Berthiaume R, Cinq-Mars D. Growth performance and carcass characteristics of dorset lambs
  fed different concentrates: Forage ratios or fresh grass. Small Rumin Res. 2011;95:113-9.
  https://doi.org/10.1016/j.smallrumres.2010.10.002
- 441 43. Phillips WA, Reuter RR, Brown MA, Fitch JQ, Rao SR, Mayeux H. Growth and performance of lambs
  442 fed a finishing diet containing either alfalfa or kenaf as the roughage source. Small Rumin Res. 2002;46:75-9.
  443 <u>http://dx.doi.org/10.1016/S0921-4488(02)00176-1</u>
- 444 44. Aurousseau B, Bauchart D, Faure X, Galot AL, Prache S, Micol D, et al. Indoor fattening of lambs raised
  445 on pasture. Part 1: Influence of stall finishing duration on lipid classes and fatty acids in the longissimus thoracis
  446 muscle. Meat Sci. 2007;76:241-52. 10.1016/j.meatsci.2006.11.005
- 447 45. Nichols ME DH, Fitch GQ, and Phillips WA, editor Feedlot performance and carcass characteristics:
  448 Comparison of small, medium, and large frame wethers backgrounded on wheat pasture. Proc West Sec Amer Soc
  449 Anim Sci; 1992.
- 450 46. Jones SDM, Burgess TD, Dupchak K, Pollock E. The growth performance and carcass composition of
  451 ram and ewe lambs fed on pasture or in confinement and slaughtered at similar fatness. Canada J Anim Sci.
  452 1984;64:631-40. 10.4141/cjas84-072
- 453 47. Purchas RW, O'Brien LE, Pendleton CM. Some effects of nutrition and castration on meat production
  454 from male suffolk cross (border leicester-romney cross) lambs. New Zealand J Agric Res. 1979;22:375-83.
  455 10.1080/00288233.1979.10430763
- 456 48. Duckett SK, Neel JP, Lewis RM, Fontenot JP, Clapham WM. Effects of forage species or concentrate
   457 finishing on animal performance, carcass and meat quality. J Anim Sci. 2013;91:1454-67. 10.2527/jas.2012-5914
- 458 49. Realini CE, Duckett SK, Brito GW, Dalla Rizza M, De Mattos D. Effect of pasture vs. Concentrate
  459 feeding with or without antioxidants on carcass characteristics, fatty acid composition, and quality of uruguayan
  460 beef. Meat Sci. 2004;66:567-77. <u>https://doi.org/10.1016/S0309-1740(03)00160-8</u>
- 50. Santos-Silva J, Mendes IA, Portugal PV, Bessa RJB. Effect of particle size and soybean oil
  supplementation on growth performance, carcass and meat quality and fatty acid composition of intramuscular lipids
  of lambs. Livestock Prod Sci. 2004;90:79-88. https://doi.org/10.1016/j.livprodsci.2004.02.013
- 46451.Gaweł E, Grzelak M. The effect of a protein-xanthophyll concentrate from alfalfa (phytobiotic) on465animal production a current review. Annals Anim Sci. 2012;12:281-9. <a href="https://doi.org/10.2478/v10220-012-0023-5">https://doi.org/10.2478/v10220-012-0023-5</a>
- 466 52. Baker NF, Cook EF, Douglas JR, Cornelius CE. The pathogenesis of trichostrongyloid parasites. Iii.
  467 Some physiological observations in lambs suffering from acute parasitic gastroenteritis. J Parasitol. 1959;45:643-51.
- 468 53. Urquhart GM, Armour, J., Duncan, J.L., Dunn, A.M. and Jennings, F.W. Vet parasitol. 2nd ed.
  469 Oxford1996.
- 470 54. Kowalczuk-Vasilev E KR, and Patkowski K. Effect of px concentrate
- of alfalfa (medicago sativa) on haematological indices of lambs' blood. . In: E.R. G, editor. Alfalfa in human and
   animal nutrition. Dzierdziówka-Lublin: Assoc Reg Loc Develop "Progress"; 2010;186-7.
- 473 and

474 55. Asma Z, Sylvie C, Laurent C, Jérôme M, Christophe K, Olivier B, et al. Microbial ecology of the rumen
475 evaluated by 454 gs flx pyrosequencing is affected by starch and oil supplementation of diets. FEMS Micro Eco.
476 2013;83:504-14. 10.1111/1574-6941.12011

477 56. Bo Trabi E, Seddik H-e, Xie F, Wang X, Liu J, Mao S. Effect of pelleted high-grain total mixed ration on
478 rumen morphology, epithelium-associated microbiota and gene expression of proinflammatory cytokines and tight
479 junction proteins in hu sheep. Anim Feed Sci Tech. 2020;263:1144-53.

- 480 <u>https://doi.org/10.1016/j.anifeedsci.2020.114453</u>
- 481 57. Fernando SC, Purvis HT, Najar FZ, Sukharnikov LO, Krehbiel CR, Nagaraja TG, et al. Rumen microbial
  482 population dynamics during adaptation to a high-grain diet. App Environ Micro. 2010;76:7482-90.
  483 10.1128/aem.00388-10
- 484
  58. Khafipour E, Li S, Plaizier JC, Krause DO. Rumen microbiome composition determined using two
  485 nutritional models of subacute ruminal acidosis. App Environ Micro. 2009;75:7115-24. 10.1128/aem.00739-09

486 59. Tajima K, Aminov RI, Nagamine T, Matsui H, Nakamura M, Benno Y. Diet-dependent shifts in the
487 bacterial population of the rumen revealed with real-time pcr. App Environ Micro. 2001;67:2766-74.
488 10.1128/aem.67.6.2766-2774.2001

489 60. Marley CL, Fraser MD, Fychan R, Theobald VJ, Jones R. Effect of forage legumes and anthelmintic
490 treatment on the performance, nutritional status and nematode parasites of grazing lambs. Vet Parasitol.
491 2005;131:267-82. <u>https://doi.org/10.1016/j.vetpar.2005.04.037</u>

492 61. Niezen J, Waghorn T, Raufaut K, Robertson H, McFarlane R. Lamb weight gain and faecal egg count
493 when grazing one of seven herbages and dosed with larvae for six weeks. Proc New Zealand Soc Anim Prod; Jan:
494 New Zealand Soc Anim Prod; 1994;15-8.

495 62. Marley CL, Cook R, Keatinge R, Barrett J, Lampkin NH. The effect of birdsfoot trefoil (lotus
496 corniculatus) and chicory (cichorium intybus) on parasite intensities and performance of lambs naturally infected
497 with helminth parasites. Vet Parasitol. 2003;112:147-55. 10.1016/s0304-4017(02)00412-0

498 63. Scales GH, Knight TL, Saville DJ. Effect of herbage species and feeding level on internal parasites and
499 production performance of grazing lambs. New Zealand J Agric Res. 1995;38:237-47.
500 10.1080/00288233.1995.9513124

- 50164.Athanasiadou S, Kyriazakis I, Jackson F, Coop RL. Effects of short-term exposure to condensed tannins502on adult trichostrongylus colubriformis. Vet Rec. 2000;146:728-32. 10.1136/vr.146.25.728
- 504 505 506 507 508

- 509
- .....
- 510
- 511
- 512
- 513
- 514

#### 

## **Tables and Figures**

	Ŋ	Yr 1		/r 2	Yr 3		
	Trea	atment <sup>1</sup>	Trea	tment <sup>1</sup>	Treatment <sup>1</sup>		
Item	ALF <sup>2</sup>	BAR <sup>3</sup>	$ALF^2$	BAR <sup>3</sup>	ALF <sup>2</sup>	BAR <sup>3</sup>	
DM, %	89.91	90.14	90.13	89.66	89.71	89.38	
CP, %	20.6	17.5	18	18	21.3	16.3	
ADF, %	30.6	15.6	33.3	21	32.5	16.9	
NDF, %	43	28.7	42.4	31.9	39.8	30.4	
TDN, %	67.7	84.8	64.6	78.6	63.6	83.3	
NE <sub>m</sub> , Mcal/lbs	0.68	0.89	0.64	0.81	0.63	0.87	
NEg, Mcal/lbs	0.41	0.59	0.37	0.55	0.36	0.58	
Cost/ton, \$	452.00	407.20	411.60	366.40	406.40	341.60	

Table 1. Nutrient Concentration of Treatment Diets, % DM basis, and Cost of Treatment Diets, \$/ton

<sup>1</sup>Diets were provided for ad libitum intake.

<sup>2</sup>ALF = pellet containing 71% alfalfa, 18% barley, 5% molasses,0.013% Bovatec, and 6.1% vitamin/mineral package; fed in both confinement and in the field.

<sup>3</sup>BAR = pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, 0.016% Bovatec, and 7.4% vitamin/mineral package; fed in both confinement and in the field.

	Treatment <sup>2,3</sup>					<i>p</i> -value				
	Confine	ement <sup>4</sup>	hent <sup>4</sup> Field <sup>5</sup>		-					
Item	CALF	CBAR	FALF	FBAR	SEM	Model	Location <sup>6</sup>	Feed <sup>7</sup>	Location*Feed	
Ending BW <sup>8</sup> , kg	50.6	50.5	51.3	49.5	0.78	0.49	0.85	0.24	0.30	
ADG, kg/d	0.25	0.25	0.26	0.23	0.01	0.49	0.83	0.21	0.30	
DMI, kg/d	2.09 <sup>a</sup>	1.67 <sup>b</sup>	2.09 <sup>a</sup>	1.62 <sup>b</sup>	0.05	< 0.01	0.59	< 0.01	0.64	
Gain to feed	0.12 <sup>b</sup>	0.15 <sup>a</sup>	0.13 <sup>b</sup>	0.14 <sup>a</sup>	0.01	0.01	0.74	0.01	0.36	
Cost of gain, \$/kg	4.33 <sup>a</sup>	3.09 <sup>b</sup>	3.95 <sup>a</sup>	3.14 <sup>b</sup>	0.21	0.00	0.42	< 0.01	0.29	

Table 2. Performance of Crossbred lambs consuming either a forage or grain based diet while in confinement or on wheat stubble fields in yr 1<sup>1</sup>

<sup>a, b</sup> Least Square Means within a row with different superscripts differ (p < 0.10).

<sup>1</sup>Start date for all treatments was September 26th.

<sup>2</sup>Pen is the experimental unit, 3 sheep per replicate confinement, 6 sheep per replicate field; six replicates per treatment.

<sup>3</sup>Diets were provided to allow for ad libitum intake.

<sup>4</sup>Confinement treatments were: CALF = pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; CBAR = pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>5</sup>Field treatments were: FALF = field fed pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; FBAR = field fed pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>6</sup>Location = confinement or field finish.

 $^{7}$ Feed = alfalfa or barley pellets.

<sup>8</sup>End date for field and confinement treatments was November 25th.

Treatment <sup>2,3</sup>						<i>p</i> -value				
	Confinement <sup>4</sup> Field <sup>5</sup>									
Item	CALF	CBAR	FALF	FBAR	SEM	Model	Location <sup>6</sup>	Feed <sup>7</sup>	Location*Feed	
Ending BW <sup>8</sup> , kg	46.0 <sup>b</sup>	45.8 <sup>b</sup>	49.6 <sup>a</sup>	49.2ª	0.87	0.01	< 0.01	0.74	0.84	
ADG, kg/d	0.24 <sup>b</sup>	0.23 <sup>b</sup>	0.30 <sup>a</sup>	0.29 <sup>a</sup>	0.01	< 0.01	< 0.01	0.72	0.71	
DMI, kg/d	2.03 <sup>b</sup>	1.58 <sup>d</sup>	2.24 <sup>a</sup>	1.74 <sup>c</sup>	0.07	<0.01	0.01	< 0.01	0.66	
Gain to feed	0.11 <sup>d</sup>	0.15 <sup>b</sup>	0.13 <sup>c</sup>	0.16 <sup>a</sup>	0.01	< 0.01	< 0.01	< 0.01	0.85	
Cost of gain, \$/kg	4.00 <sup>a</sup>	2.74°	3.42 <sup>b</sup>	2.00 <sup>d</sup>	0.21	<0.01	< 0.01	< 0.01	0.32	

Table 3. Performance of Crossbred lambs consuming either a forage or grain based diet while in confinement or on wheat stubble fields in yr 2<sup>1</sup>

<sup>a, b, c, d</sup> Least Square Means within a row with different superscripts differ (p < 0.10).

<sup>1</sup>Start date for all treatments was September 4th.

<sup>2</sup>Pen is the experimental unit, 3 sheep per replicate confinement, 6 sheep per replicate field; six replicates per treatment.

<sup>3</sup>Diets were provided to allow for ad libitum intake.

<sup>4</sup>Confinement treatments were: CALF = pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; CBAR = pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>5</sup>Field treatments were: FALF = field fed pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; FBAR = field fed pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>6</sup>Location = confinement or field finish.

 $^{7}$ Feed = alfalfa or barley pellets.

<sup>8</sup>End date for field and confinement treatments was November 4<sup>th</sup>.

		Treatr	ment <sup>2,3</sup>			<i>p</i> -value				
	Confinement <sup>4</sup> Field <sup>5</sup>									
Item	CALF	CBAR	FALF	FBAR	SEM	Model	Location <sup>6</sup>	Feed <sup>7</sup>	Location*Feed	
Ending BW <sup>8</sup> , kg	51.75 <sup>a</sup>	50.51ª	52.84 <sup>b</sup>	53.73 <sup>b</sup>	0.75	< 0.01	< 0.01	0.78	0.09	
ADG, kg/d	0.22 <sup>a</sup>	0.21 <sup>a</sup>	0.24 <sup>b</sup>	0.25 <sup>b</sup>	0.01	< 0.01	< 0.01	0.81	0.08	
DMI, kg/d	$2.08^{a}$	1.86 <sup>b</sup>	2.14 <sup>a</sup>	2.00 <sup>b</sup>	0.07	0.03	0.11	< 0.01	0.57	
Gain to feed	0.13	0.10	0.10	0.11	0.01	0.53	0.65	0.70	0.17	
Cost of gain, \$/kg	4.15 <sup>a</sup>	3.36 <sup>b</sup>	3.72 <sup>a</sup>	2.75°	0.22	<0.01	< 0.01	< 0.01	0.65	

Table 4. Performance of Crossbred lambs consuming either a forage or grain based diet while in confinement or on wheat stubble fields in yr 3<sup>1</sup>

<sup>a, b, c</sup> Least Square Means within a row with different superscripts differ (p < 0.10).

<sup>1</sup>Start date for all treatments was September 1<sup>st</sup>.

<sup>2</sup>Pen is the experimental unit, 3 sheep per replicate confinement, 6 sheep per replicate field; six replicates per treatment.

<sup>3</sup>Diets were provided to allow for ad libitum intake.

<sup>4</sup>Confinement treatments were: CALF = pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; CBAR = pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>5</sup>Field treatments were: FALF = field fed pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; FBAR = field fed pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>6</sup>Location = confinement or field finish.

 $^{7}$ Feed = alfalfa or barley pellets.

<sup>8</sup>End date for field and confinement treatments was November 5th.

		Treat	ment <sup>2,3</sup>			<i>p</i> -value				
	Confine	Confinement <sup>4</sup>		Field <sup>5</sup>						
Item	CALF	CBAR	FALF	FBAR	SEM	Model	Location <sup>6</sup>	Feed <sup>7</sup>	Location*Feed	
Carcass weight, kg	25.1	24.2	25.9	26.6	1.13	0.46	0.16	0.90	0.44	
Dressing Percent	49.18 <sup>b</sup>	46.45 <sup>b</sup>	51.92 <sup>ab</sup>	52.85ª	1.54	0.02	0.01	0.54	0.22	
Back fat thickness, cm	0.34	0.38	0.43	0.28	0.07	0.56	0.94	0.49	0.21	
Ribeye area, cm <sup>2</sup>	5.30 <sup>b</sup>	5.33 <sup>b</sup>	6.03 <sup>a</sup>	6.32 <sup>a</sup>	0.21	< 0.01	< 0.01	0.45	0.55	
Leg Score <sup>8</sup>	406	413	434	422	8.56	0.13	0.04	0.72	0.28	
Conformation <sup>8</sup>	413	419	434	413	8.08	0.20	0.34	0.34	0.09	
Flank Streaking <sup>9</sup>	294	237	256	259	23.99	0.39	0.73	0.25	0.21	
Quality Grade <sup>8</sup>	425	407	406	409	15.08	0.77	0.57	0.61	0.47	
WBSF <sup>10</sup> , kg	3.8 <sup>a</sup>	2.7 <sup>b</sup>	3.8 <sup>a</sup>	3.1 <sup>b</sup>	0.30	0.03	0.56	< 0.01	0.53	

Table 5. Carcass Characteristics of Crossbred lambs consuming either a forage or grain based diet while in confinement or on wheat stubble fields in yr 1<sup>1</sup>

<sup>a,b</sup> Least Square Means within a row with different superscripts differ (p < 0.10).

<sup>1</sup>Start date for all treatments was September 26th; End date for field and confinement treatments was November 25th.

<sup>2</sup>Pen is the experimental unit, 3 sheep per replicate confinement, 6 sheep per replicate field; six replicates per treatment.

<sup>3</sup>Diets were provided to allow for ad libitum intake.

<sup>4</sup>Confinement treatments were: CALF = pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; CBAR = pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>5</sup>Field treatments were: FALF = field fed pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; FBAR = field fed pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>6</sup>Location = confinement or field finish.

 $^{7}$ Feed = alfalfa or barley pellets.

 $^{8}325 = \text{Good minus}; 350 = \text{Good}; 375 = \text{Good plus}; 425 = \text{Choice minus}; 450 = \text{Choice}; 475 = \text{Choice plus}; 525 = \text{Prime minus}; 550 = \text{Prime}; 575 = \text{Prime plus}.$ 

 $^{9}100$  to 199 = Practically Devoid; 200 to 299 = Traces; 300 to 399 = Slight; 400 to 499 = Small.

<sup>10</sup>Warner-Bratzler shear force (WBSF).

		Treat	ment <sup>2,3</sup>					p-value	
-	Confine	ment <sup>4</sup>	Field	5					
Item	CALF	CBAR	FALF	FBAR	SEM	Model	Location <sup>6</sup>	Feed <sup>7</sup>	Location*Feed
Carcass weight, kg	24.5	23.5	25.6	24.4	1.01	0.54	0.33	0.28	0.90
Dressing Percent	49.71	50.85	51.49	50.31	0.95	0.59	0.52	0.98	0.23
Back fat thickness, cm	0.25	0.24	0.33	0.32	0.04	0.33	0.07	0.79	0.94
Ribeye area, cm <sup>2</sup>	5.91	5.73	6.25	6.32	0.20	0.15	0.03	0.79	0.56
Leg Score <sup>8</sup>	419 <sup>ab</sup>	406 <sup>b</sup>	438 <sup>a</sup>	431 <sup>a</sup>	8.92	0.09	0.02	0.30	0.73
Conformation <sup>8</sup>	419 <sup>ab</sup>	406 <sup>b</sup>	438 <sup>a</sup>	431 <sup>a</sup>	8.92	0.09	0.02	0.30	0.73
Flank Streaking <sup>9</sup>	259	289	324	288	35.39	0.61	0.35	0.92	0.33
Quality Grade <sup>8</sup>	403	419	443	413	21.65	0.59	0.42	0.72	0.28
WBSF <sup>10</sup> , kg	4.2	4.3	4.1	3.7	0.48	0.76	0.45	0.67	0.56

Table 6. Carcass Characteristics of Crossbred lambs consuming either a forage or grain based diet while in confinement or on wheat stubble fields in yr 2<sup>1</sup>

<sup>a,b</sup> Least Square Means within a row with different superscripts differ (p < 0.10).

<sup>1</sup>Start date for all treatments was September 4th; End date for field and confinement treatments was November 4th.

<sup>2</sup>Pen is the experimental unit, 3 sheep per replicate confinement, 6 sheep per replicate field; six replicates per treatment.

<sup>3</sup>Diets were provided to allow for ad libitum intake.

<sup>4</sup>Confinement treatments were: CALF = pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; CBAR = pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>5</sup>Field treatments were: FALF = field fed pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; FBAR = field fed pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>6</sup>Location = confinement or field finish.

 $^{7}$ Feed = alfalfa or barley pellets.

 $^{8}$ 325 = Good minus; 350 = Good; 375 = Good plus; 425 = Choice minus; 450 = Choice; 475 = Choice plus; 525 = Prime minus; 550 = Prime; 575 = Prime plus.  $^{9}$ 100 to 199 = Practically Devoid; 200 to 299 = Traces; 300 to 399 = Slight; 400 to 499 = Small.

<sup>10</sup>Warner-Bratzler shear force (WBSF).

Table 7. Whole blood counts of Crossbred lambs consuming either a forage or grain based diet while in confinement or on wheat stubble fields in yr 1<sup>1</sup>

		ment <sup>2,3</sup>				p	-value		
	Confine	Confinement <sup>4</sup>		eld <sup>5</sup>	-				
Item <sup>6</sup>	CALF	CBAR	FALF	FBAR	SEM	Model	Location <sup>7</sup>	Feed <sup>8</sup>	Location*Feed
White blood cell	7.34	7.69	7.46	7.02	0.43	< 0.01	0.44	0.90	0.28
Red blood cell	12.99	12.66	12.92	12.62	0.32	0.13	0.82	0.23	0.95
Hemoglobin	14.07	20.13	14.21	13.68	2.56	0.26	0.14	0.19	0.12
Hematocrit	42.00 <sup>a</sup>	40.56 <sup>b</sup>	42.19 <sup>a</sup>	41.28 <sup>ab</sup>	0.63	0.01	0.38	0.03	0.62
Mean cell volume	32.36	32.21	41.72	32.83	7.67	0.70	0.43	0.47	0.49
Mean cell hemoglobin	10.83 <sup>ab</sup>	10.62 <sup>b</sup>	11.05 <sup>a</sup>	10.97 <sup>ab</sup>	0.18	< 0.01	0.06	0.34	0.66
Mean cell hemoglobin concentration	33.43 <sup>ab</sup>	33.19 <sup>b</sup>	33.65 <sup>a</sup>	33.25 <sup>b</sup>	0.23	< 0.01	0.45	0.08	0.68
Red cell distribution width	25.80 <sup>b</sup>	25.62 <sup>b</sup>	26.00 <sup>b</sup>	27.26 <sup>a</sup>	0.59	< 0.01	0.06	0.27	0.14

<sup>a, b</sup> Least Square Means within a row with different superscripts differ (p < 0.10).

<sup>1</sup>Start date for all treatments was September 26th; End date for field and confinement treatments was November 25th.

<sup>2</sup>Pen is the experimental unit, 3 sheep per replicate confinement, 6 sheep per replicate field; six replicates per treatment.

<sup>3</sup>Diets were provided to allow for ad libitum intake.

<sup>4</sup>Confinement treatments were: CALF = pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; CBAR = pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>5</sup>Field treatments were: FALF = field fed pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; FBAR = field fed pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>6</sup>All items are ending counts.

 $^{7}$ Location = confinement or field finish.

 $^{8}$ Feed = alfalfa or barley pellets.

		Treatr	nent <sup>2,3</sup>			<i>p</i> -value					
	Confir	inement <sup>4</sup>		Field <sup>5</sup>							
Item <sup>6</sup>	CALF	CBAR	FALF	FBAR	SEM	Model	Location <sup>7</sup>	Feed <sup>8</sup>	Location*Feed		
White blood cell	7.41 <sup>a</sup>	6.62 <sup>ab</sup>	7.21 <sup>a</sup>	6.46 <sup>b</sup>	0.38	< 0.01	0.58	0.02	0.96		
Red blood cell	12.28 <sup>ab</sup>	11.97 <sup>b</sup>	12.55 <sup>a</sup>	12.08 <sup>b</sup>	0.25	0.03	0.37	0.07	0.70		
Hemoglobin	13.26 <sup>b</sup>	12.93 <sup>b</sup>	13.82 <sup>a</sup>	13.34 <sup>b</sup>	0.21	< 0.01	0.01	0.03	0.66		
Hematocrit	38.84 <sup>b</sup>	38.84 <sup>b</sup>	40.10 <sup>a</sup>	39.28 <sup>ab</sup>	0.55	0.06	0.08	0.39	0.38		
Mean cell volume	31.72	32.20	32.20	32.62	0.53	< 0.01	0.33	0.33	0.94		
Mean cell hemoglobin	10.82 <sup>ab</sup>	10.74 <sup>b</sup>	11.04 <sup>a</sup>	11.05 <sup>a</sup>	0.12	< 0.01	0.02	0.75	0.66		
Mean cell hemoglobin concentration	34.15 <sup>ab</sup>	33.39°	34.39 <sup>a</sup>	33.92 <sup>b</sup>	0.24	< 0.01	0.06	< 0.01	0.48		
Red cell distribution width	26.11	25.96	26.10	26.58	0.56	0.22	0.53	0.74	0.52		

Table 8. Whole blood counts of Crossbred lambs consuming either a forage or grain based diet while in confinement or on wheat stubble fields in yr  $2^1$ 

<sup>a, b</sup> Least Square Means within a row with different superscripts differ (p < 0.10).

<sup>1</sup>Start date for all treatments was September 4th; End date for field and confinement treatments was November 4th.

<sup>2</sup>Pen is the experimental unit, 3 sheep per replicate confinement, 6 sheep per replicate field; six replicates per treatment.

<sup>3</sup>Diets were provided to allow for ad libitum intake.

<sup>4</sup>Confinement treatments were: CALF = pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; CBAR = pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>5</sup>Field treatments were: FALF = field fed pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; FBAR = field fed pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>6</sup>All items are ending counts.

<sup>7</sup>Location = confinement or field finish.

 $^{8}$ Feed = alfalfa or barley pellets.

Table 9. Ending counts of internal parasite eggs per gram (EPG) in Crossbred lambs consuming either a forage or grain based diet while in confinement or on wheat stubble fields in yr  $1^1$ 

	Treatment <sup>2,3</sup>					<i>p</i> -value				
	Confinement <sup>4</sup> Field <sup>5</sup>									
Item <sup>6</sup>	CALF	CBAR	FALF	FBAR	SEM	Model	Location <sup>7</sup>	Feed <sup>8</sup>	Location*Feed	
Trichostrongyle type EPG	6.95	3.86	9.98	3.95	0.52	0.05	0.70	0.16	0.73	
Nematodirus spp. EPG	10.57	3.34	12.98	2.67	0.58	0.11	0.98	0.02	0.72	

<sup>1</sup>Start date for all treatments was September 26th; End date for field and confinement treatments was November 25th.

<sup>2</sup>Pen is the experimental unit, 3 sheep per replicate confinement, 6 sheep per replicate field; six replicates per treatment.

S

<sup>3</sup>Diets were provided to allow for ad libitum intake.

<sup>4</sup>Confinement treatments were: CALF = pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; CBAR = pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>5</sup>Field treatments were: FALF = field fed pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; FBAR = field fed pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>6</sup>All Counts had an addition of a constant of 1 then were LOG transformed.

 $^{7}$ Location = confinement or field finish.

 $^{8}$ Feed = alfalfa or barley pellets.

Table 10. Worm burden in the abomasum and first meter of the small intestine in Crossbred lambs consuming either a forage or grain based diet while in confinement or on wheat stubble fields in yr  $1^1$ 

		Treatr	ment <sup>2,3</sup>			<i>p</i> -value				
	Confine	ement <sup>4</sup>	Field <sup>5</sup>		_					
Item <sup>6</sup>	CALF	CBAR	FALF	FBAR	SEM	Model	Location <sup>7</sup>	Feed <sup>8</sup>	Location*Feed	
Abomasum Total count	3	24	111	10	1.09	0.20	0.25	0.88	0.07	
Teladorsagia circumcincta	3	24	111	10	1.09	0.20	0.25	0.88	0.07	
Small Intestine Total Count	$2^{bc}$	18 <sup>b</sup>	176 <sup>a</sup>	$0^{c}$	0.94	0.01	0.48	0.08	< 0.01	
Teladorsagia circumcincta	2	0	0	0	0.59	0.47	0.38	0.38	0.38	
Nematodirus	0°	18 <sup>b</sup>	176 <sup>a</sup>	0°	0.72	< 0.01	0.12	0.12	< 0.01	

<sup>a, b, c</sup> Least Square Means within a row with different superscripts differ (p < 0.10).

<sup>1</sup>Start date for all treatments was September 26th; End date for field and confinement treatments was November 25th.

<sup>2</sup>Pen is the experimental unit, 3 sheep per replicate confinement, 6 sheep per replicate field; six replicates per treatment.

<sup>3</sup>Diets were provided to allow for ad libitum intake.

<sup>4</sup>Confinement treatments were: CALF = pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; CBAR = pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>5</sup>Field treatments were: FALF = field fed pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; FBAR = field fed pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>6</sup>All Counts had an addition of a constant of 1 then were LOG transformed.

 $^{7}$ Location = confinement or field finish.

 $^{8}$ Feed = alfalfa or barley pellets.

Table 11. Ending counts of internal parasite eggs per gram (EPG) in Crossbred lambs consuming either a forage or grain based diet while in confinement or on wheat stubble fields in yr  $2^1$ 

	Treatment <sup>2,3</sup>						I	p-value	
	Confinement <sup>4</sup> Field <sup>5</sup>								
Item <sup>6</sup>	CALF	CBAR	FALF	FBAR	SEM	Model	Location <sup>7</sup>	Feed <sup>8</sup>	Location*Feed
Trichostrongyle type EPG	50.92	10.97	28.51	18.31	0.51	0.29	0.92	0.04	0.24
Nematodirus spp. EPG	9.99 <sup>ab</sup>	0.52°	20.62ª	5.60 <sup>b</sup>	0.53	< 0.01	0.02	< 0.01	0.40

<sup>a, b, c</sup> Least Square Means within a row with different superscripts differ (p < 0.10).

<sup>1</sup>Start date for all treatments was September 4th; End date for field and confinement treatments was November 4th.

<sup>2</sup>Pen is the experimental unit, 3 sheep per replicate confinement, 6 sheep per replicate field; six replicates per treatment.

S

<sup>3</sup>Diets were provided to allow for ad libitum intake.

<sup>4</sup>Confinement treatments were: CALF = pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; CBAR = pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>5</sup>Field treatments were: FALF = field fed pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; FBAR = field fed pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>6</sup>All Counts had an addition of a constant of 1 then were LOG transformed.

 $^{7}$ Location = confinement or field finish.

 $^{8}$ Feed = alfalfa or barley pellets.

Table 12. Worm burden in the abomasum and first meter of the small intestine in Crossbred lambs consuming either a forage or grain based diet while in	
confinement or on wheat stubble fields in yr $2^1$	

		Treat	ment <sup>2,3</sup>				I	o-value	
	Confin	nement <sup>4</sup>	Field <sup>5</sup>		-				
Item <sup>6</sup>	CALF	CBAR	FALF	FBAR	SEM	Model	Location <sup>7</sup>	Feed <sup>8</sup>	Location*Feed
Abomasum Total count	49	15	12	44	1.50	0.88	0.93	0.98	0.44
Teladorsagia circumcincta	41	15	12	44	1.48	0.90	0.98	0.94	0.47
Haemonchus contortus	6 <sup>a</sup>	$0^{b}$	$0^{b}$	$0^{b}$	0.57	0.07	0.11	0.11	0.11
Small Intestine Total Count	29	0	3	3	1.10	0.23	0.74	0.14	0.16
Teladorsagia circumcincta	2	0	0	0	0.58	0.43	0.34	0.34	0.34
Nematodirus	8	0	3	3	1.14	0.60	0.86	0.32	0.37

<sup>a, b</sup> Least Square Means within a row with different superscripts differ (p < 0.10).

<sup>1</sup>Start date for all treatments was September 4th; End date for field and confinement treatments was November 4<sup>th</sup>.

<sup>2</sup>Pen is the experimental unit, 3 sheep per replicate confinement, 6 sheep per replicate field; six replicates per treatment.

<sup>3</sup>Diets were provided to allow for ad libitum intake.

<sup>4</sup>Confinement treatments were: CALF = pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; CBAR = pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>5</sup>Field treatments were: FALF = field fed pellet containing 71% alfalfa, 18% barley, 5% molasses, and 6% vitamin/mineral package; FBAR = field fed pellet containing 60% barley, 26% alfalfa, 4% molasses, 2.5% soybean-hi pro, and 7.5% vitamin/mineral package.

<sup>6</sup>All Counts had an addition of a constant of 1 then were LOG transformed.

 $^{7}$ Location = confinement or field finish.

 $^{8}$ Feed = alfalfa or barley pellets.