

JAST (Journal of Animal Science and Technology) TITLE PAGE
Upload this completed form to website with submission

ARTICLE INFORMATION	Fill in information in each box below
Article Type	Research article
Article Title (within 20 words without abbreviations)	Comparison of pork belly characteristics and weights of primal cuts between gilt and barrow of LYD pigs measured by AutoFomIII
Running Title (within 10 words)	Comparison of gilt and barrow characteristics measured by AutoFomIII
Author	Eunyoung Ko ^{1, a} , Yunhwan Park ^{2, a} , Kwangwook Park ¹ , Changhyun Woo ¹ , Jaeyoung Kim ² , Jungseok Choi ² ^a These authors contributed equally to this work.
Affiliation	1Dodram Pig Farmers Cooperative, Icheon 17405, Korea 2Department of Animal Science, Chungbuk national university, Cheongju 28644, Korea
ORCID (for more information, please visit https://orcid.org)	Eunyoung Ko (https://orcid.org/0000-0003-8686-1762) Yunhwan Park (https://orcid.org/0000-0002-2239-6697) Kwangwook Park (https://orcid.org/0000-0003-2671-0687) Changhyun Woo (https://orcid.org/0000-0003-8103-9817) Jaeyoung Kim (https://orcid.org/0000-0002-2847-1731) Kwansuk Kim (https://orcid.org/0000-0002-5895-4398) Jungseok Choi (https://orcid.org/0000-0001-8033-0410)
Competing interests	No potential conflict of interest relevant to this article was reported.
Funding sources State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available.	This work was supported by the Dodram Quality Control Management Team of Dodram Pig Farmers Cooperative. The Korean Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry and Fisheries (IPET) through the Agri-Bioindustry Technology Development Program, funded by Ministry of Agriculture, Food and Rural Affairs (MAFRA) (Project No. 321028-5), Korea. Grant (715003-07) from the Research Center for Production Management and Technical Development for High Quality Livestock Products through Agriculture, Food and Rural Affairs Convergence Technologies Program for Educating Creative Global Leader, Ministry of Agriculture, Food and Rural Affairs.
Acknowledgements	This work was supported by the Dodram Quality Control Management Team of Dodram Pig Farmers Cooperative. The Korean Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry and Fisheries (IPET) through the Agri-Bioindustry Technology Development Program, funded by Ministry of Agriculture, Food and Rural Affairs (MAFRA) (Project No. 321028-5), Korea. Grant (715003-07) from the Research Center for Production Management and Technical Development for High Quality Livestock Products through Agriculture, Food and Rural Affairs Convergence Technologies Program for Educating Creative Global Leader, Ministry of Agriculture, Food and Rural Affairs.
Availability of data and material	Upon reasonable request, the datasets of this study can be available from the corresponding author.
Authors' contributions Please specify the authors' role using this form.	Conceptualization: Eunyoung Ko, Yunhwan Park, Jaeyoung Kim Data curation: Eunyoung Ko, Changhyun Woo, Jungseok Choi Formal analysis: Yunhwan Park, Jaeyoung Kim Methodology: Yunhwan Park, Jaeyoung Kim Software: Yunhwan Park Validation: Kwangwook Park, Changhyun Woo Investigation: Eunyoung ko, Changhyun Woo, Kwansuk Kim Writing - original draft: Yunhwan Park, Eunyoung Ko Writing - review & editing: Jungseok Choi, Eunyoung Ko, Yunhwan Park

Ethics approval and consent to participate	This article does not require IRB/IACUC approval because there are no human and animal participants.
---------------------------------------------------	------------------------------------------------------------------------------------------------------

CORRESPONDING AUTHOR CONTACT INFORMATION

For the corresponding author (responsible for correspondence, proofreading, and reprints)	Fill in information in each box below
First name, middle initial, last name	Choi, Jungseok
Email address – this is where your proofs will be sent	jchoi@chungbuk.ac.kr
Secondary Email address	
Address	Department of Animal Science, Chungbuk National University, Cheonju, Chungcheongbuk-do Korea, 28644
Cell phone number	+82-10-3235-2127
Office phone number	+82-43-261-2551
Fax number	+82-43-261-2773

ACCEPTED

Abstract (up to 350 words)

Currently, pigs breed in Korea are LYD (Landrace x Yorkshire x Duroc) crossbred pigs. Pigs used as fresh meat are gilts and barrows. However, the current supply of pork is not satisfying Korean consumers. In addition, the comparison of carcasses between gilts and barrows only studies carcass weight, backfat thickness, or meat quality, and there are very few studies comparing carcass characteristics. The purpose of this study was to compare characteristics of 7 primal cuts of gilts and barrows as measured by AutoFom III. A total of 350,179 pigs were used, including 176,461 gilts and 173,718 barrows. Characteristics of seven primal cuts were measured using AutoFom III. In the case of carcass weight, there was no significant difference in grade 1+. For all other survey items except for grade 2, gilts showed significantly ($p < 0.05$) higher values. For all grades except for pork belly, amounts of the remaining six primal cuts were higher in gilts (all $p < 0.05$, Table 6). In addition, the ratio of intermuscular fat in the pork belly of barrows showed a higher value than that in the pork belly of gilts ($p < 0.05$, Table 7). The amount of pork belly, which is the most popular among consumers in Korea, not only produced more production than gilts in barrows, but also showed a higher value than gilts in barrows for the ratio of intermuscular fat affecting taste. In summary, gilts produced higher yields than barrows in all parts except pork belly. For the production of only pork belly, barrows showed higher production than gilts.

Keywords (3 to 6): LYD, Gilt, Barrow, Grade, Belly, AutoFom III

Introduction

Currently, the most popular finishing pig in Korea is LYD species, a three-way crossbred of Landrace, Yorkshire, and Duroc. This LYD species created as a three-way crossbred has advantages of the three species by crossing a Landrace species with a high fertility number, a fast-growing Yorkshire species, and a Duroc species with a high meat yield [4, 5]. Among LYD

species produced in this way, gilts and barrows are used as meat [6]. Barrows refer to boars that have been surgically castrated at a young age. Castration is performed because it lowers levels of androsterone and skatole in the meat. Androsterone and skatole are the main causes of boar odor. They can cause an unpleasant taste and odor in the meat of boars. Because of this, male piglets are usually surgically castrated within the first week of life to reduce boar taint and aggressive behavior and to eliminate sexual behavior [7, 17].

In Korea, grilling pork is the most popular culture. Cuts such as pork belly and Boston butt with high fat are popular, whereas cuts such as hams and loin with low fat are less popular. Therefore, there is a large difference in the price of primal cut of pork in Korea [11]. Among pork cuts with high fat, pork belly is one of the most preferred cuts [3]. Since fat in the pork belly can enhance the flavor and juiciness of the meat [8], it is related to meat quality [9]. Pork belly accounts for about 13-18% of pig carcass weight [10]. The supply of pork belly is important in Korea. However, the supply of pork belly could not satisfy Korean consumers [11]. Also, since the amount of pork belly can only be known after the pig carcass grade is evaluated, the amount of pork belly cannot be known immediately [12].

Since the income of Korean people is increasing, the per capita meat consumption is also increasing. In 2018, out of 53.9 kg of meat consumption per capita in Korea, the amount of pork was 27 kg, accounting for more than 50% of all meat consumption [1]. At the same time, the number of pigs slaughtered in Korea increased from 15 million in 2014 to 17 million in 2018 [2]. As so many pigs are slaughtered, an automatic grading machine is needed in the slaughterhouse. AutoFom III is one of automatic grading machines using ultrasound. AutoFom III is currently used in slaughterhouses in 14 overseas countries including North America and Europe. AutoFom III showed very high accuracy that predicting the amount of pork belly and ham of pig carcass, regression coefficients were respectively 0.98, 0.93. Also Choi et al. (2018) [21] indicated that the prediction accuracy was high when the predicted value and the actual value of Korean pigs measured by AutoFom

III were compared.

In Korea, pig carcass grade is measured according to 21 criteria such as sex, appearance, meat quality, and defects based on carcass weight and backfat thickness [16] [Table 1]. According to KAPE (Korea Institute for Animal Products Quality Evaluation, 2019), the price per kg of pork was 4,222 won for grade 1+, 4,134 won for grade 1, and 3,960 won for grade 2. Therefore, pig carcass grade is an important factor that shapes the final market price of finishing pigs.

Big data analysis technology is an essential element when analyzing large-capacity data, both structured and unstructured, to extract available information and to actively respond to or predict changes in the pig industry based on analyzed data [22]. It has been predicted that the number of livestock slaughtered will increase in the future. Thus, accumulated data will increase accordingly. Most papers that have compared gilts and barrows so far have compared carcass weight and backfat thickness as well as meat quality. However, very few studies have compared seven primal cuts. In addition, the number of samples investigated was small. One study has reported few data comparing carcass yields from gilts and barrows [23]. Therefore, the purpose of this study was to compare characteristics of pork belly of gilts and barrows, weights of seven primal cuts, and differences in carcass characteristics according to the Korean pig carcass grading system based on data measured with an AutoFom III.

Materials and Methods

Animals

A total of 350,719 LYD crossbred pigs (Landrace, Yorkshire, and Duroc) were used as experimental materials, including 176,461 gilts and 173,718 barrows. All pigs were bred and slaughtered under the same environment from January 2019 to December 2019. And the feed was provided by the Dodram pig farmers' association in accordance with specific standards of

Dodram union. All pigs were transferred to Dodram LPC in Anseong, Gyeonggi Province, Korea and slaughtered in accordance with the Livestock Sanitation Management Act.

Measurement of pig carcass weight and backfat thickness

Pig carcass weight was measured with an electronic scale and expressed in unit of kg. Backfat thicknesses between the last thoracic vertebra and the first lumbar vertebrae and between the 11th and 12th lumbar vertebrae of the left half carcass were measured with a ruler and expressed in unit of mm. The average of two backfat thickness values was calculated.

Measurement of primal cut weight and content of pork belly using AutoFom III

AutoFom III (Frontmatec Somerum A/S, Denmark) was used after the head was removed and before evisceration of the carcass. AutoFom III is an equipment that performs measurement using ultrasound. It is largely divided into four parts: an ultra-sound transducer array, an acquisition module, a data processing workstation, and a personal computer. It also has 16 ultrasound transducers that are excited in turn with a repeated frequency of about 5 kHz. As the carcass is scanned, an ultrasound image is generated, and 48 image parameters are generated that provide information on lean meat and fat yield [21]. Further methods of using AutoFom III have been described in our previous study [23]. Carcass weight, backfat thickness, pork belly characteristics, and weights of seven primal cuts (belly, Boston butt, ribs, tenderloin, loin, picnic shoulder, ham) were measured with the AutoFom III equipment based on the data classified by sex of the pig.

Statistical analysis

A t-test was used to test the significance of each investigation. Skewness and Kurtosis were measured to know the disperse of each investigated graph. The regression equation was

calculated using weights of seven parts by dividing the carcass by gender and placing the carcass weight as the dependent variable (y_i). Afterwards, coefficient of determination and correlation were calculated. Data of gender and grade were investigated using the following multiple linear regression model.

$$y_i = \mu + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3 + \beta_4 x_4 + \beta_5 x_5 + \beta_6 x_6 + \beta_7 x_7$$

where μ was the intercept and β was the slope of the independent variable x_i , which was the weight of 7 cut meats measured by AutoFom III. x_1 was pork belly, x_2 was pork loin, x_3 was rib, x_4 was sirloin, x_5 was tenderloin, x_6 was forelimb, and x_7 was hind leg. All statistical analyses were performed using SPSS software version 25.0 (SPSS Inc., Chicago, IL, USA).

Results and Discussion

In Table 2, skewness and kurtosis of carcass weight and backfat thickness were measured to investigate characteristics and distribution according to sex and grade. Figure 2 shows the distribution of carcass weight and backfat thickness through a box plot. A normal distribution is a case where the mean, median, and mode all agree with skewness and kurtosis of 0. However, in this study, skewness showed negative values for carcass weight and backfat thickness of all sexes with grade 1+, carcass weights of gilts with grade 1, backfat thickness of all sexes, and barrows with grade 2 (Table 2). Except for this, skewness and kurtosis of carcass weight and backfat thickness showed positive values ($k > 0$, Table 2). If the skewness has a negative value, it is interpreted that the distribution is skewed to the left. If it has a positive value, it is interpreted to be skewed to the right. However, according to Kline (2005), if the absolute value of skewness is less than 3 and the absolute value of Kurtosis is less than 7, then normal distribution is suitable. Results showed that the absolute value of skewness was less than 3 and the absolute value of kurtosis was less than 7, conforming to normality (Table 2).

There was no significant difference in carcass weight or backfat thickness for grade 1+ between gilts and barrows (both $p > 0.05$). On the other hand, for grade 1, the carcass weight of gilts showed a significantly higher value than that of barrows. For grade 2, the carcass weight of gilts showed a significantly lower value than that of barrows. Except for grade 1+, backfat thickness of barrow was significantly thicker than that of gilts. Overall, gilts showed a tendency to be heavier than that of barrows in the case of carcass weight, whereas barrows tended to have thicker backfat thickness than gilts ($p < 0.05$, Table 3). These results were contrary to results of Larzul (1997) [13] and Kim (2006) [18]. They showed that barrows had higher carcass weight and thicker backfat. Kim et al. (2006) [18] have found that barrows have heavier carcass weight than gilts without showing significant difference ($p > 0.05$), although barrows have thicker backfat than gilts ($p < 0.05$). The present study showed different results of carcasses of gilts and barrows compared to results of Kim et al. (2006) [18]. This might be due to differences in breeding area, feed, and age at marketing. According to the study of Larzul et al. (1997) [13], gilts grew later than barrows. Thus, their carcass weights were less. However, backfat thickness in the study of Larzul et al. (1997) [13] was similar to that of the present study. The reason why carcass weight values of gilts and barrows were opposite from those in the study of Larzul et al. (1997) [13] might be due to differences in breed, breeding area, and feed.

Gilts and barrows were classified according to the Korean pig carcass grading system. Results are presented in Table 4. According to the Korean pig carcass grading system, in the case of gilts ($n = 176,461$), grade 1+ showed the largest proportion at 39.13% ($n = 69,049$), followed by grade 1 at 37.32% ($n = 65,850$) and grade 2 at 23.55% ($n = 41,562$). On the other hand, in the case of barrows ($n = 173,718$), grade 1 showed the highest proportion at 37.97% ($n = 65,989$), followed by grade 1+ at 33.88% ($n = 58,862$) and grade 2 at 28.13% ($n = 48,867$). Results shown in Table

4 were consistent with results of Kim et al. (2012) [22]. Both showed that gilts had the highest proportion of grade 1+ and barrows had the highest proportion of grade 1. A similar trend was found for the percentage of pigs slaughtered in 2019 based on the data released by the Livestock Products Quality Assessment Service in 2019 [1]. These results were also similar to results of Kevin et al. (2020) [35], where S, the highest grade of gilts, showed a higher ratio than that of barrows. Although pig carcass grade used in Europe differs from that in Korea in terms of meat production, the high proportion of the highest grade in gilts cannot be ignored.

In Table 5, skewness and kurtosis were measured to investigate the disperse of seven primal cuts according to sex and grade. Measurements of weights for the seven primal cuts shown in Table 5 were estimated with AutoFom III. In the above result, in the case of comparison between gilts and barrows by grade, as grade decreased, the value of skewness increased, showing a tendency to approach 0. Also, most values of kurtosis were negative, showing a less sharpening shape. In the case of overall comparison between gilts and barrows, the value of skewness was negative for loin and ham of gilts and barrows and for tenderloin of gilts. Seven primal cuts showing negative values can be interpreted as skewing the distribution to the left. On the other hand, in the case of kurtosis, all areas of gilts and barrows showed positive values ($k > 0$, Table 5). Skewness and kurtosis did not indicate 0 because the mean, median, and mode did not match. However, the normality was verified according to Kline (2005).

As a result of measuring weights of seven primal cuts with AutoFom III, for all grades of pork belly, the most popular cut in Korea, barrows showed significantly higher values than gilts ($p < 0.05$). Except for pork belly, amounts of the remaining six primal cuts showed higher values in gilts than in barrows. In the study of Kevin et al. (2020) [35], gilts showed significantly higher values of areas such as ham and shoulder than barrows. In addition, similar results were found

for pork belly, which showed a higher value in barrows than in gilts. In Table 1 carcass weights of gilts showed significantly higher values than those of barrows, which might be the reason why all parts except pork belly showed high values in gilts than in barrows ($p < 0.05$). The reason why gilts showed higher values of carcass weight and lean meat content than barrows was similar to that in one study [33] that found that estrogen of gilt hormone could improve protein deposition, similar to research results of Tanghe et al. (2015) [19] and other dissection experiments [34]. Also, compared to gilts and slow-growing pigs, barrows and fast-growing pigs might have higher amounts of pork belly. The fact that loin, picnic shoulder, and Boston butt were produced more in gilts than in barrows was consistent with a previous study [27], suggesting that there were also differences according to growth rate. In the case of the picnic shoulder, it was mentioned that it increased with the slaughter weight. It can also be explained by growth difference between pork belly and other six parts cut of meat due to allometric growth [14].

As a result of measuring characteristics of pork belly, the most popular cut in Korea, with AutoFom III, gilts showed significantly higher values of lean meat weight than barrows. However, barrows showed significantly higher values of subcutaneous fat and intermuscular fat than gilts ($p < 0.05$, Table 7). According to gender and growth rate, the amount of pork belly and fat content of pork belly could be different. Fast-growing pigs have higher amounts of pork belly and fat contents in pork belly than slow-growing pigs [26]. In addition, a higher fat accumulation has been observed in barrows than in gilts in a previous study [25].

This can be explained by the deposition of more protein by estrogen of gilt hormone. According to Berit et al. (2010) [31], castration almost eliminates testosterone. However, testosterone deficiency can increase food intake and body fat, resulting in a large amount of fat deposition. Combining this with the study of Hector et al. (2014) [32] showing that low testosterone levels in

men could cause abdominal obesity, it can be seen that amounts of pork belly and fat contents in barrows are higher than those in gilts. Newcom et al. (2002) [24] have shown a trend consistent with the finding of higher production of pork belly from barrows than from gilts. However, in the study of Kevin et al. (2020) [35], gilts showed no significant difference value compared to barrows ($p < 0.0001$). Such differences in results between the study of Kevin et al. (2020) [35] and the present study might be due to different breed of pigs used in the experiment. From results shown in Table 7, proportions of pork belly from gilts and barrows were consistent results of a previous study [10] showing that pork belly accounted for 13-18% of carcass weight. Lean meat percentage of pork belly was 22-55%. Fat percentage was 20-60%, consistent with above results. In addition, the ratio of subcutaneous fat in pork belly fat was double that of intermuscular fat. According to Van-Ba Hoa et al. (2021) [10], Pork belly with a high fat ratio (fat > 31%) was found to have higher meat quality, fatty acid content, and flavor compound, which were important for eating quality. Therefore, both gilts and barrows showed a high-fat content (over 31%) of pork belly. Thus, it could be considered that the taste of pork belly from both sexes is good. According to Kim et al. (2015) [27], when sensory of pork belly with less than 40% fat was compared to that of pork belly with 40% to 45% fat, pork belly with more than 40% fat showed higher juiciness, tenderness, flavor, fattiness, and overall Palatability. However, there was no significant difference value. However, based on the data that the preference for pork belly with intermuscular fat content of 13 to 15% is the highest [28] and that the highest successful bid price is formed [29], It is thought that the pork belly of barrows is more preferred. It was found that the amount of intermuscular fat had a significant correlation with both taste and flavor of meat [30].

Correlation analysis was conducted to investigate the relationship between carcass weight and weights of seven primal cuts according to grade and between backfat thickness and weights of seven primal cut [Tables 8-1, 8-2]. In both gilts and barrows, correlations between carcass

weight and seven primal cuts showed generally high values. However, correlations between backfat thickness and seven primal cuts showed low values. In the case of gilts, the correlation between the amount of pork belly and carcass weight was 0.725 for grade 1+, 0.901 for grade 1, and 0.960 for grade 2, indicating a high correlation coefficient ($p < 0.01$). In the case of barrows, the correlation between the amount of pork belly and carcass weight was 0.714 for grade 1+, 0.888 for grade 1, and 0.9492 for grade 2, showing a high correlation coefficient ($p < 0.01$). Correlations between carcass weight and weights of seven primal cuts showed generally high correlation coefficients ($p < 0.01$, Table 8), whereas correlations between backfat thickness and weights of seven primal cuts showed low values, including negative values ($p > 0.01$, Table 8). Overall, correlation coefficients between carcass weight and seven primal cuts showed higher values as the grade decreased. On the other hand, correlation coefficients between backfat thickness and the seven primal cuts showed a tendency to increase with negative values as the grade decreased.

A regression analysis was performed to investigate relationships between carcass weight and seven primal cuts. Multiple regression analysis was performed with carcass weight as the dependent variable and the weight of each of the seven primal cuts as the independent variable. The linear regression equation between carcass weight and seven primal cuts by grade according to gender is shown as follows:

Gilts:

Grade 1+:

$$y = 6.723 - 1.578x_1 - 0.892x_2 + 52.001x_3 + 5.507x_4 + 14.578x_5 - 14.123x_6 - 1.660x_7;$$

Grade 1:

$$y = 2.164 - 1.202x_1 - 20797x_2 + 54.464x_3 + 5.766x_4 + 19.602x_5 - 14.98x_6 - 1.7x_7;$$

Grade

2:

$$y = -2.23518 + 1.016x_1 - 9.906x_2 + 49.406x_3 + 4.869x_4 + 31.954x_5 - 13.012x_6 - 1.33x_7$$

Barrows:

Grade

1+:

$$y = 5.195 - 1.131x_1 - 2.053x_2 + 50.754x_3 + 4.964x_4 + 17.16x_5 - 13.867x_6 - 1.36x_7$$

Grade

1:

$$y = 0.37 - 0.276x_1 - 5.46x_2 + 51.419x_3 + 4.943x_4 + 24.984x_5 - 13.958x_6 - 1.351x_7$$

Grade

2:

$$y = -3.841 + 1.888x_1 - 12.546x_2 + 46.687x_3 + 4.215x_4 + 37.071x_5 - 12.013x_6 - 1.074x_7$$

The coefficient of determination is divided into grades according to gender. For gilts, it was 0.943 for grade 1+, 0.984 for grade 1, 0.993 for grade 2. For barrows, it was 0.943 for grade 1+, 0.982 for grade 1, and 0.993 for grade 2, showing high values. These showed a similar trend to a previous study [21]. Park et al. (2022) [12] have made a regression equation without gender distinction, whereas this study created a regression equation between grades of carcass weight and 7 primal cuts weights by classifying gender. Considering that a fairly high coefficient of determination was obtained through the regression equation, it was thought that weights of the seven primal cuts divided by gender in the Korean pig industry could be easily predicted.

This study analyzed and compared to carcass characteristics of gilts and barrows, the weight of primal cut, and the characteristics of pork belly measured by AutoFom III. AutoFom III data were estimated with a mathematical formula developed based on anatomical experiments [20]. The reliability of the estimate has been proven in a previous study by Choi et al. (2018) [21]. Carcass weights of gilts were significantly higher than those of barrows ($p < 0.05$). For all cuts

except for the pork belly, gilts showed higher values ($p < 0.05$) than barrows. Also, the increase in the amount of pork belly and the increase in body fat due to the lack of testosterone in barrows resulted in higher amounts of pork belly of barrows and higher muscle fat content than gilts. In other words, it is considered that the pork belly, which consumers prefer, appears more in barrows than in gilts. The proportion of intermuscular fat was also higher in the pork belly of barrows. Thus, it is thought that the pork belly of barrows is better than that of gilts.

Acknowledgments

This work was supported by the Dodram Quality Control Management Team of Dodram Pig Farmers Cooperative.

The Korean Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry and Fisheries (IPET) through the Agri-Bioindustry Technology Development Program, funded by Ministry of Agriculture, Food and Rural Affairs (MAFRA) (Project No. 321028-5), Korea. Grant (715003-07) from the Research Center for Production Management and Technical Development for High Quality Livestock Products through Agriculture, Food and Rural Affairs Convergence Technologies Program for Educating Creative Global Leader, Ministry of Agriculture, Food and Rural Affairs.

References

1. Ministry of Agriculture F, Rural Affairs SK. Major Statistics of Agriculture, Food and Rural Affairs during 2019. Ministry of Agriculture, Food and Rural Affairs Sejong, Korea; 2019.
2. Evaluation KIfAPQ. Animal Products Grading Statistical Yearbook. Korea Institute for Animal Products Quality Evaluation Sejong, Korea; 2019.
3. Choe J-H, Yang H-S, Lee S-H, Go G-W. Characteristics of pork belly consumption in South Korea and their health implication. *Journal of Animal Science and Technology*. 2015;57(1):1-7.
4. Cho S, Park B, Kim J, Kim M, Seong P, Kim Y, et al. Carcass yields and meat quality by live weight of Korean native black pigs. *Journal of animal science and technology*. 2007;49(4):523-30.
5. Hong K, Kim B, Son Y, Kom B. Effects of the mating system on fattening performance and meat quality in commercial pigs. *J Anim Sci Technol*. 2001;43:139-48.
6. 한국식육과학연구회. MeatScience. 1st ed. Sjprinting; 2018
7. Font-i-Furnols M, Skrlep M, Aluwé M, editors. Attitudes and beliefs of consumers towards pig welfare and pork quality. IOP Conference Series: Earth and Environmental Science; 2019: IOP Publishing.
8. Fernandez X, Monin G, Talmant A, Mourot J, Lebret B. Influence of intramuscular fat content on the quality of pig meat—2. Consumer acceptability of m. longissimus lumborum. *Meat science*. 1999;53(1):67-72.
9. Hausman GJ, Basu U, Du M, Fernyhough-Culver M, Dodson MV. Intermuscular and intramuscular adipose tissues: bad vs. good adipose tissues. *Adipocyte*. 2014;3(4):242-55.
10. Hoa V-B, Seol K-H, Seo H-W, Seong P-N, Kang S-M, Kim Y-S, et al. Meat quality characteristics of pork bellies in relation to fat level. *Animal Bioscience*. 2021;34(10):1663.
11. Oh S-H, See M. Pork preference for consumers in China, Japan and South Korea. *Asian-Australasian journal of animal sciences*. 2012;25(1):143.
12. Park Y, Ko E, Park K, Woo C, Kim J, Lee S, et al. Correlation between the Korean pork grade system and the amount of pork primal cut estimated with AutoFom III. *Journal of Animal Science and Technology*. 2022;64(1):135.
13. Larzul C, Lefaucheur L, Ecolan P, Gogue J, Talmant A, Sellier P, et al. Phenotypic and genetic parameters for longissimus muscle fiber characteristics in relation to growth, carcass, and meat quality traits in large white pigs. *Journal of animal science*. 1997;75(12):3126-37.
14. Walstra P. Growth and carcass composition from birth to maturity in relation to feeding level and sex in Dutch Landrace pigs: Wageningen University and Research; 1980.

15. Soladoye P, Shand P, Aalhus J, Gariépy C, Juárez M. Pork belly quality, bacon properties and recent consumer trends. *Canadian Journal of Animal Science*. 2015;95(3):325-40.
16. Kim J, Han H-D, Lee WY, Wakholi C, Lee J, Jeong Y-B, et al. Economic analysis of the use of vcs2000 for pork carcass meat yield grading in Korea. *Animals*. 2021;11(5):1297.
17. Prunier A, Bonneau M, Von Borell E, Cinotti S, Gunn M, Fredriksen B, et al. A review of the welfare consequences of surgical castration in piglets and the evaluation of non-surgical methods. *ANIMAL WELFARE-POTTERS BAR THEN WHEATHAMPSTEAD*-. 2006;15(3):277.
18. Kim G-W, In B-S. Carcass grade and characteristics by carcass weight and backfat thickness of pigs. *Food Science of Animal Resources*. 2006;26(2):183-8.
19. Tanghe S, Millet S, Hellebuyck S, Van Meensel J, Buys N, De Smet S, et al. Effect of sex and sire on lean meat percentage and weight of primal cuts of pork using Autofom data. 66th EAAP book of abstracts2015. p. 166.
20. Brøndum J, Egebo M, Agerskov C, Busk H. On-line pork carcass grading with the Autofom ultrasound system. *Journal of Animal Science*. 1998;76(7):1859-68.
21. Choi JS, Kwon KM, Lee YK, Joeng JU, Lee KO, Jin SK, et al. Application of AutoFom III equipment for prediction of primal and commercial cut weight of Korean pig carcasses. *Asian-Australasian journal of animal sciences*. 2018;31(10):1670.
22. Kim J-S. Consideration of big data utilization and related technologies. *Review of Korea Contents Association*. 2012;10(1):34-40.
23. Boler D, Puls C, Clark D, Ellis M, Schroeder A, Matzat P, et al. Effects of immunological castration (Improvest) on changes in dressing percentage and carcass characteristics of finishing pigs. *Journal of Animal Science*. 2014;92(1):359-68.
24. Newcom DW, Baas T, Mabry J, Goodwin R. Genetic parameters for pork carcass components. *Journal of animal science*. 2002;80(12):3099-106.
25. Fredeen H. Yields and dimensions of pork bellies in relation to carcass measurements. *Journal of Animal Science*. 1980;51(1):59-68.
26. Correa J, Faucitano L, Laforest J, Rivest J, Marcoux M, Gariépy C. Effects of slaughter weight on carcass composition and meat quality in pigs of two different growth rates. *Meat Science*. 2006;72(1):91-9.
27. Kim HS. Shape and Characteristics of Korean's Favorite Pork Belly. *Food Science of Animal Resources and Industry*. 2015;4(2):30-44.
28. <https://www.mk.co.kr/news/economy/view/2022/05/386053/>, 2022
29. <https://www.pignpork.com/news/articleView.html?idxno=740>, 2021

30. Frank D, Joo S-T, Warner R. Consumer acceptability of intramuscular fat. *Korean journal for food science of animal resources*. 2016;36(6):699.
31. Christoffersen BO, Gade LP, Golozoubova V, Svendsen O, Raun K. Influence of castration-induced testosterone and estradiol deficiency on obesity and glucose metabolism in male Göttingen minipigs. *Steroids*. 2010;75(10):676-84.
32. Escobar-Morreale HF, Álvarez-Blasco F, Botella-Carretero JI, Luque-Ramírez M. The striking similarities in the metabolic associations of female androgen excess and male androgen deficiency. *Human Reproduction*. 2014;29(10):2083-91.
33. Claus R, Weiler U. Endocrine regulation of growth and metabolism in the pig: a review. *Livestock Production Science*. 1994;37(3):245-60.
34. Nautrup BP, Van Vlaenderen I, Aldaz A, Mah C. The effect of immunization against gonadotropin-releasing factor on growth performance, carcass characteristics and boar taint relevant to pig producers and the pork packing industry: A meta-analysis. *Research in Veterinary Science*. 2018;119:182-95.
35. Kress K, Hartung J, Jasny J, Stefanski V, Weiler U. Carcass characteristics and primal pork cuts of gilts, boars, immunocastrates and barrows using AutoFOM III data of a commercial abattoir. *Animals*. 2020;10(10):1912.
36. Janiszewski P, Borzuta K, Lisiak D, Grześkowiak E, Stanisławski D. Prediction of primal cuts by using an automatic ultrasonic device as a new method for estimating a pig-carcass slaughter and commercial value. *Anim Prod Sci*. 2019;59:1183–9.

Tables and Figures




Figure 1. Box plot of characteristics of carcass weight and backfat thickness of gilts and barrows.




Figure 2. Box plot of lean amount by part of gilts and barrows measured by AutoFom III

Table 1. Korean pig carcass grading system

Primary carcass grade	Carcass weight (kg)	Backfat thickness (mm)
1 ⁺	83 - 92	17 - 24
1	80 - 82	15 - 27
	83 - 92	15 - 16
	83 - 92	25 - 27
	93 - 97	15 - 27
2	Neither 1 ⁺ nor 1	

Korean Ministry of Agriculture, Food and Rural Affairs Notification 2014-4.

Table 2. Disperse characteristics of carcass weight and backfat thickness of gilts and barrows

	Skewness	Kurtosis		Skewness	Kurtosis
1+ grade carcass weight			1+ grade backfat thickness		
Gilts	-0.117	-1.014	Gilts	-0.184	-0.978
Barrow	-0.017	-1.053	Barrows	-0.624	-0.511
1 grade carcass weight			1 grade backfat thickness		
Gilts	-0.054	-1.414	Gilts	-0.006	-1.374
Barrows	0.110	-1.352	Barrows	-0.585	-0.899
2 grade carcass weight			2 grade backfat thickness		
Gilts	0.206	-1.173	Gilts	0.382	-0.489
Barrows	0.051	-1.042	Barrows	-0.440	-0.804
All of grades Carcass weight			All of grades Backfat thickness		
Gilts	0.222	0.357	Gilts	0.164	1.170
Barrows	0.285	0.350	Barrows	0.108	0.409

Population number for gilts and barrows were 176,461 and 173,718, respectively. Skewness means that the degree to which the distribution is skewed out of symmetry. Kurtosis means that the sharpness of the frequency distribution.

Table 3. Pig carcass weight and backfat thickness

	Gilt	Barrow
Carcass weight		
1+ Grade	87.57±2.65	87.83±2.62
1 Grade	88.65±5.41 ^a	88.01±5.32 ^b
2 Grade	88.16±9.63 ^b	88.54±9.19 ^a
All of Grades	88.21±5.97 ^a	88.10±6.13 ^b
Backfat thickness		
1+ Grade	20.87±2.04	21.68±1.91
1 Grade	20.69±3.92 ^b	22.43±3.64 ^a
2 Grade	20.07±6.91 ^b	24.51±6.70 ^a
All of Grades	20.61±4.33 ^b	22.78±4.47 ^a

Each value is presented as mean ± SD. Population numbers for Gilts and Barrows were 176,461 and 173,718, respectively. ^{a-b} Means within the same row with different superscripts are significantly different ($p < 0.05$).

Table 4. Grade percentage of sex according to the Korean pig meat grading system

	1+	1	2
Gilt	39.13%	37.32%	23.55%
Barrow	33.88%	37.97%	28.13%

Population numbers for gilts and barrows were 176,461 and 173,718, respectively. Population numbers for grades 1+, 1, and 2 were 127,911, 131,839, and 90,429, respectively.

ACCEPTED

Table 5. Disperse of lean amount by part of barrows and gilts measured by AutoFom III

	skewness	kurtosis		skewness	kurtosis
Gilt belly			Barrow belly		
1+	0.093	-0.21	1+	0.061	-0.26
1	0.009	-0.99	1	0.061	-0.93
2	0.204	-1.13	2	-0.086	-0.97
All Grades	0.227	0.339	All Grades	0.176	0.325
Gilt Boston butt			Barrow Boston butt		
1+	-0.18	-0.14	1+	-0.13	-0.16
1	-0.10	-0.79	1	0.10	-0.72
2	0.21	-0.79	2	0.30	-0.63
All Grades	0.014	0.408	All Grades	0.048	0.369
Gilt rib			Barrow rib		
1+	-0.10	-0.57	1+	-0.02	-0.59
1	-0.05	-1.17	1	0.16	-1.08
2	0.23	-0.99	2	0.26	-0.85
All Grades	0.226	0.329	All Grades	0.294	0.328
Gilt loin			Barrow loin		
1+	-0.25	0.28	1+	-0.19	0.18
1	-0.24	0.04	1	-0.11	-0.02
2	-0.07	-0.01	2	0.05	-0.02
All Grades	-0.131	0.484	All Grades	-0.106	0.425
Gilt tenderloin			Barrow tenderloin		
1+	-0.23	0.201	1+	-0.18	0.13
1	-0.18	-0.28	1	-0.01	-0.24
2	0.11	-0.45	2	0.23	-0.33
All Grades	-0.025	0.449	All Grades	0.008	0.411
Gilt Picnic Shoulder			Barrow Picnic Shoulder		
1+	-0.13	-0.37	1+	-0.19	0.18
1	-0.08	-0.98	1	-0.11	-0.02
2	0.23	-0.90	2	0.30	-0.75
All Grades	0.059	0.372	All Grades	0.092	0.325
Gilts ham			Barrow ham		
1+	-0.23	0.05	1+	-0.17	-0.02
1	-0.14	-0.56	1	0.05	-0.52
2	0.18	-0.65	2	0.28	-0.51
All Grades	-0.119	0.509	All Grades	-0.076	0.289

Population numbers for gilts and barrows were 176,461 and 173,718, respectively. Skewness means that the degree to which the distribution is skewed out of symmetry. Kurtosis means that the sharpness of the frequency distribution.

ACCEPTED

Table 6. Lean meat weights of seven primal cuts as measured by AutoFom III

		Gilt	Barrow
Belly			
	1+ Grade	12.12±0.52 ^b	12.29±0.54 ^a
	1 Grade	12.24±0.90 ^b	12.40±0.90 ^a
	2 Grade	12.19±1.57 ^b	12.64±1.54 ^a
	All Grades	9.94±0.70 ^b	10.10±0.68 ^a
Boston Butt			
	1+ Grade	4.55±0.20 ^a	4.46±0.20 ^b
	1 Grade	4.59±0.32 ^a	4.45±0.32 ^b
	2 Grade	4.56±0.49 ^a	4.43±0.48 ^b
	All Grades	3.89±0.32 ^a	3.76±0.33 ^b
Rib			
	1+ Grade	3.67±0.11 ^a	3.65±0.12 ^b
	1 Grade	3.70±0.21 ^a	3.65±0.21 ^b
	2 Grade	3.68±0.35 ^a	3.65±0.34 ^b
	All Grades	3.67±0.23 ^a	3.62±0.23 ^b
Tenderloin			
	1+ Grade	0.99±0.05 ^a	0.96±0.06 ^b
	1 Grade	1.00±0.08 ^a	0.96±0.08 ^b
	2 Grade	0.99±0.11 ^a	0.95±0.11 ^b
	All Grades	0.99±0.08 ^a	0.96±0.08 ^b
Loin			
	1+ Grade	5.83±0.48 ^a	5.60±0.49 ^b
	1 Grade	5.89±0.61 ^a	5.55±0.62 ^b
	2 Grade	5.84±0.78 ^a	5.45±0.82 ^b
	All Grades	6.43±0.64 ^a	6.11±0.68 ^b
Picnic shoulder			
	1+ Grade	8.22±0.28 ^a	5.60±0.49 ^b
	1 Grade	8.29±0.48 ^a	5.55±0.62 ^b
	2 Grade	8.27±0.78 ^a	8.09±0.75 ^b
	All Grades	7.98±0.54 ^a	7.75±0.57 ^b
Ham			
	1+ Grade	17.31±0.77 ^a	16.95±0.79 ^b
	1 Grade	17.45±1.17 ^a	16.90±1.18 ^b
	2 Grade	17.33±1.74 ^a	16.75±1.74 ^b
	All Grades	16.29±1.19 ^a	15.69±1.27 ^b

Each value is presented as mean \pm SD. Population numbers for gilts and barrows were 176,461 and 173,718, respectively. ^{a-b} Means within the same row with different superscripts are significantly different ($p < 0.05$).

ACCEPTED

Table 7. Weights of lean meat, subcutaneous fat, and muscle fat of pork belly according to sex measured by AutoFom III

	Gilt	Barrow
Lean meat(kg)		
1+ Grade	10.05±0.39a	9.45±0.39
1 Grade	10.14±0.65a	9.95±0.65
2 Grade	10.10±1.02a	9.92±1.00
All Grades	10.10±0.68 ^a	9.94±0.70 ^b
Lean meat percentage		
1+ Grade	55.57±2.17a	54.53±2.28
1 Grade	55.58±2.74a	54.04±2.85
2 Grade	55.56±3.82a	52.80±3.99
All Grades	55.57±2.84 ^a	53.86±3.13 ^b
Subcutaneous fat weight		
1+ Grade	4.89±0.38	5.03±0.39a
1 Grade	4.94±0.57	5.13±0.57
2 Grade	4.92±0.96	5.37±0.95a
All Grades	4.92±0.63 ^b	5.16±0.67 ^a
Subcutaneous fat percentage		
1+ Grade	26.51±1.33	26.98±1.40a
1 Grade	26.49±1.65	27.26±1.70a
2 Grade	26.46±2.26	27.94±2.32a
All Grades	26.49±1.71 ^b	27.36±1.85 ^a
Intermuscular fat weight		
1+ Grade	2.29±0.28	2.41±0.29a
1 Grade	2.32±0.40	2.49±0.42a
2 Grade	2.31±0.66	2.68±0.67a
All Grades	2.30±0.44 ^b	2.52±0.48 ^a
Intermuscular fat percentage		
1+ Grade	12.41±1.28	13.06±1.36a
1 Grade	12.41±1.62	13.36±1.70a
2 Grade	12.45±2.29	14.13±2.40a
All Grades	12.42±1.69 ^b	13.47±1.88 ^a
Belly fat percentage		
1+ Grade	38.89±2.45	40.06±2.58a
1 Grade	38.88±3.10	40.62±3.23a
2 Grade	38.91±4.34	42.06±4.55a

	All Grades	38.89±3.22 ^b	40.83±3.56 ^a
Ratio belly weight to carcass weight			
	1+ Grade	13.77	13.97
	1 Grade	13.75	14.09
	2 Grade	13.85	14.20
	All Grades	13.81	14.11

Each value is presented as mean ± SD. Population numbers for gilts and barrows were 176,461 and 173,718, respectively. A, b Means within a same row with different superscripts are significantly different ($p < 0.05$).

ACCEPTED

Table 8-1. Correlation coefficients between characteristics of carcass and weights of seven primal cuts of gilts

		Belly	Boston Butt	Ribs	Tenderloin	Loin	Picnic Shoulder	Ham
Carcass weight	1+	0.725**	0.718**	0.880**	0.553**	0.401**	0.803**	0.635**
	1	0.901**	0.856**	0.948**	0.733**	0.579**	0.903**	0.799**
	2	0.960**	0.906**	0.968**	0.811**	0.670**	0.938**	0.864**
Backfat thickness	1+	0.239**	-0.137**	-0.066**	-0.169**	-0.168**	-0.124**	-0.155**
	1	0.469**	0.032**	0.149**	-0.058**	-0.127**	0.068**	-0.015**
	2	0.648**	0.244**	0.372**	0.124**	-0.002**	0.290**	0.184**

** $p < 0.01$.

Table 8-2. Correlation coefficients between characteristics of carcass and weights of seven primal cuts of barrows

		Belly	Boston Butt	Ribs	Tenderloin	Loin	Picnic Shoulder	Ham
Carcass weight	1+	0.714**	0.702**	0.874**	0.534**	0.389**	0.389**	0.618**
	1	0.888**	0.841**	0.941**	0.714**	0.562**	0.562**	0.780**
	2	0.949**	0.885**	0.961**	0.778**	0.635**	0.924**	0.836**
Backfat thickness	1+	0.248**	-0.100**	-0.043**	-0.129**	-0.128**	-0.128**	-0.117**
	1	0.443**	-0.001	0.106**	-0.080**	-0.136**	-0.136**	-0.045**
	2	0.616**	0.169**	0.300**	0.056**	-0.052**	0.219**	0.110**

** $p < 0.01$.