# JAST (Journal of Animal Science and Technology) TITLE PAGE Upload this completed form to website with submission

ARTICLE INFORMATION	Fill in information in each box below						
Article Type	Research article						
Article Title (within 20 words without abbreviations)	Effects of various Cooling Methods and Drinking Water						
	Temperatures on Reproductive Performance and Behavior in Heat						
	Stressed Sows						
Running Title (within 10 words)	Cooling Methods and Drinking Temperatures for Sows.						
Author	Habeeb Taiudeen <sup>1,#</sup> Joseph Moturi <sup>1,#</sup> Abdolreza Hosseindoust <sup>1</sup>						
	lin Soo Kim1*						
Affiliation	<sup>1</sup> Department of Bio-health Convergence, Kangwon National						
	University, Chuncheon, 24341, Republic of Korea						
	<sup>2</sup> Swine Division, National Institute of Animal Science, Rural						
	Development Administration, Cheonan 31000, Republic of Korea						
	<sup>3</sup> Poultry Science Division, Livestock Research Development,						
	National Institute of Animal Science, RDA, 4937, Republic of Korea.						
ORCID (for more information, please visit	Habeeb Tajudeen (https://orcid.org/0000-0002-5623-3175)						
https://orcid.org)	Joseph Moturi (https://orcid.org/ 0000-0002-0626-0255)						
	Abdolreza Hosseindoust (https://orcid.org/0000-0001-9191-0613)						
	SangHun Ha (https://orcid.org/0000-0003-3779-1144)						
	JunYoung Mun (https://orcid.org/0000-0002-3075-7157)						
	YoHan Choi ( <u>https://orcid.org/0000-0003-4710-4731</u> )						
	SooJin Sa (https://orcid.org/0000-0002-2634-5109)						
	JinSoo Kim (https://orcid.org/0000-0002-9518-7917)						
Competing interests	The authors declare that they had no conflict of interests during this						
	research.						
Funding sources State funding sources (grants, funding sources,	The project was funded by the Rural Development Administration						
equipment, and supplies). Include name and number of grant if available	Research Project (Detailed Task Number: PJ01479601) and the						
	2021 Rural Development Administration, National Institute of						
	Livestock Science, Republic of Korea.						

Acknowledgements	The Authors of this project appreciates the support received by the
	Rural Development Administration Research Project and the 2021
	Rural Development Administration, National Institute of Livestock
	Science, Republic of Korea.
Availability of data and material	Upon reasonable request, the datasets of this study can be available
	from the corresponding author.
Authors' contributions	All authors contributed to this work.
Please specify the authors' role using this form.	
	Conceptualization: J.S. Kim, H. Tajudeen, S.H. Ha.
	Data curation: J. Moturi, Y.H. Choi
	Formal analysis: A. Hosseindoust, S.J. Sa
	Methodology: J.S. Kim, H. Tajudeen, J. Y. Mun
	Software: A. Hosseindoust, J. Y. Mun
	Validation: J. Moturi, S.J. Sa
	Investigation: S.H. Ha, Y.H. Choi
	Writing - original draft: H. Tajudeen, A. Hosseindoust., and J.S. Kim.
Ethics approval and consent to participate	The animal care and experimental protocols used in the present
	study were approved by the Institution of Animal Care and Use
	Committee, Kangwon National University. (Ethical code: KW-
	210503-6).
4	

## CORRESPONDING AUTHOR CONTACT INFORMATION

For the corresponding author (responsible for correspondence, proofreading, and reprints)	Fill in information in each box below
First name, middle initial, last name	JinSoo Kim
Email address – this is where your proofs will be sent	kjs896@kangwon.ac.kr
Secondary Email address	
Address	Department of Animal Industry Convergence, Kangwon National
	University, Chuncheon, 24341, Republic of Korea

Cell phone number	+82-102566-5961
Office phone number	+82-102566-5961
Fax number	
10	

#### 12 Abstract

13 The purpose of this study is to evaluate the effects of multiple cooling systems and different drinking water 14 temperatures (DWT) on the performance of sows and their hair cortisol levels during heat stress. In this study, the 15 effect of four different cooling systems: air conditioner (AC), cooling pad (CP), snout cooling (SC), and mist spray 16 (MS), and two drinking water treatments (DWT), namely low water temperature (LWT) and high water temperature 17 (HWT) on 48 multiparous sows (Landrace  $\cdot$  Yorkshire; 242.84  $\pm$  2.89 kg) was tested. The experiment is based on the 18 use of eight replicas during a 21-days test. Different behaviors were recorded under different cooling treatments in 19 sows. As a result, behaviors such as drinking, standing, and position change were found to be lower in sows under 20 the AC and CP treatments than in those under the SC and MS treatments. Lying behavior increased under the AC 21 and CP systems as compared with that under the SC and MS, systems. The average daily feed intake (ADFI) in sows 22 and weight at weaning in piglets was higher under the AC, CP, and LWT treatments than under the SC, MS and 23 HWT treatments. Sows subjected to SC and MS treatment showed higher hair cortisol levels, rectal temperature, and 24 respiratory rate during lactation than those under AC and CP treatments. Hair cortisol levels, rectal temperature, and 25 respiratory rate were also higher under the HWT than under the LWT treatment. As per the results of this study, the 26 LWT has no significant effect on any of the behavioral factors. Taken together, the use of AC and CP cooling 27 treatment is highly recommended to improve the behavior and to reduce the stress levels in lactating sows. 

28

#### 29 Keywords:

30 31

32

# Introduction

Temperature-humidity index, Water temperature, Cortisol, Cooling system, Lactation, Piglets

33 The environmental conditions play an important role in maintaining a suitable production output for animals. The 34 temperature is rising globally, as projected by Huddart et al. [1], and therefore, heat stress (HS) is becoming a more 35 common and serious problem. HS can hinder the performance and general welfare of animals leading to a reduction 36 in productivity and food security [2]. The problem of HS is not only restricted to tropical areas but is extended to 37 temperate countries as well, especially during the extreme summer months owing to the acute events of summer 38 heatwaves [3]. Sows are extremely poor in resisting heat, not only because they lack functional sweat glands but 39 also because they have a thick layer of subcutaneous adipose tissue that prevents radiative heat dissipation [5]. HS 40 typically decreases the feed intake in sows [3,4], which results in bodily issues such as loss of body condition, 41 negative energy balance, some reproductive issues related to anestrus, long weaning to estrus intervals, low

42 farrowing rates, and poor litter size [6]. HS can also result in decreased milk production, leading to a negative 43 impact on the overall development and weight of a piglet at the time of weaning during lactation [7]. HS plays an 44 important role during both lactation and gestation such that the gestational HS can increase the mortality in embryos, 45 leading to a direct impact on the farrowing rate and litter size [6]. This also extends to late gestation, wherein HS can 46 increase the number of stillborn piglets [8] and decrease the weight in newborn piglets [9].

47 During hot weather, various environmental changes can aid heat loss by convection, conduction, radiation, and 48 evaporation. On-farm HS in sows can be reduced using a variety of methods. More importantly, several cooling 49 systems can be employed to improve the efficacy of animal welfare. However, there are limited studies that compare 50 the effects of different cooling systems and drinking water temperatures on the performance of sows during HS. The 51 purpose of this study is to evaluate the effects of four different cooling systems, namely air conditioner (AC), 52 cooling pad (CP), snout cooling (SC), and mist spray (MS), on the well-being of sows during HS. In addition to the 53 cooling systems, we combined different drinking water temperatures (DWT):low water temperature (LWT) and high 54 water temperature (HWT), to study their effects on different parameters such as sow performance, litter performance, 55 behavior, hair cortisol level, rectal temperature, and respiratory rate.

56

57

# **Materials and Methods**

58 The animal care and experimental protocols used in this study received approval by the Institutional Animal Care 59 and Use Committee of Kangwon National University. (Ethical code: 210503-6).

60

#### 61 Test Animals, Feed and Experimental Design

62 Forty-eight multiparous sows (Landrace  $\cdot$  Yorkshire; 242.84 $\pm$ 2.89 kg) were used to examine the outcome of 63 various cooling facilities and two DWT in the farrowing house based on the reproductive efficiency and stress score 64 of the sows during HS in the summer period of June to July 2021 at the Teaching and Research Farm of Kangwon National University. Only sows with successful farrowing performance were selected for the experiment, 65 66 and the average litter size was 12.34 pigs/sow. The corn-soybean meal used was formulated to meet or exceed 67 the nutrient recommendation of the National Research Council (10), and it was made available at ad libitum in 68 powdered form as shown in Table 1. This study was conducted as a 2×4 factorial arrangement with 2 drinking water 69 temperatures (LWT; 15°C, HWT; 25°C) and 4 cooling systems (AC, CP, SC, MS). Eight replicas of the six 70 treatments were built, such that there was one sow per replica. The test was conducted during the lactation period

71 (21 days upon delivery), and the animals were selected after 112 days of gestation. The floor of each stall was 72 equipped with a cuboidal cast iron pad (56 cm  $\cdot$  56 cm  $\cdot$  5 cm; Cooling Sow System, Nooyen Corporation, The 73 Netherlands), which served as a surface where sows could rest while lying down. The serpentine conduits installed 74 beneath the floor pads enabled the circulation of cool water via an appropriate channel. The water temperature in the 75 reserve buffer tank was maintained at around 15 °C using a liquid-to-liquid heat pump (Carrier Corp, Indianapolis, 76 IN; Model GW62wl). Pumps were used to deliver 1.9 L/min of chilled water to each farrowing stall using the 77 Tichelmann settings [11], which allows each stall to receive cool water at a corresponding temperature. The other 78 rooms with water at 25 °C had identical cast iron flooring pads. In order to ensure that the temperature of water at 79 nipple drinkers is consistent, chilled water at 15 °C was circulated repeatedly via insulated water lines to each nipple 80 drinker. In the second method of treatment, the drinking water was neither chilled nor circulated. The treatments 81 carried out in separate farrowing rooms, had the same delivery frame, with a length and width of 2,400 mm and 82 1,800 mm, respectively. Each delivery frame in the farrowing room was installed with a heating box and a lamp. 83 The cooling systems were programmed to activate when the average temperature reached 25°C in the farrowing 84 house.

85

#### 86 Temperature and Humidity Index

87 Temperature and humidity data monitoring devices (TENMARS, TM-305U, Taiwan) were installed in each
88 farrowing room to examine the environmental temperature during the experiment. Temperature and humidity were
89 evaluated in terms of temperature and humidity index (THI) [12]. The mathematical formula for THI is as follows:

90 THI = (1.8 X T + 32) - [(0.55 - 0.0055 X RH) X (1.8 X T - 26))].

91 where T = temperature in degree (°C)

92 RH = relative humidity in percent (%)

93

#### 94 Sow and Litter Reproductive Performance

95 After 112 days of pregnancy and weaning, the body weight and backfat thickness of sows were thoroughly 96 examined. The backfat thickness of the sow was evaluated three times through the use of an ultrasonic measuring 97 tool (SONG KANG GLC, Anyscan BF, Korea) at 6.5 cm from the 10th rib, and the mean values were determined. 98 The average daily feed intake (ADFI) was estimated by recording the feed intake during the weaning phase and 99 estrus relapse period. The number and weight of the piglets at farrowing and weaning of each sow were used to

100 estimate the reproductive performance.

101

### 102 Cortisol Concentration, Rectal Temperature, and Respiratory Rate

In our final experiment, freshly grown hair from individual animals were collected and used for the hair cortisol analysis. Before that, a part of the dorsal hair of the sows and piglets was removed at farrowing. The hair were washed three times with isopropanol, followed by drying in a vacuum dryer at 35°C, and then placed in a EML plastic tube containing steel pellets and a bead beater (tacoTMPrep, 50/60 Hz 2A, GeneReach). Hair cortisols were extracted using methanol after crushing at the Biotechnology Corp. in Taiwan. A cortisol ELISA kit (ADI-900-071, Enzo Life Sciences, Inc., US) was used to determine the concentration of the extracted sample [13].

During the course of this experiment, the rectal temperatures and respiratory rates of these animals were measured twice a day (11:30 and 13:00 hours) using a digital thermometer (SATO, SK-1260, Japan) which was inserted into the rectum and a stopwatch which helped to visually observe the abdominal motions per 1 minute in sow, respectively. The mean of these recorded values was used.

113

#### 114 Statistical Analysis

The statistical analysis system (SAS) and general linear model (GLM) was employed to analyze the data collected in the experiment at  $2\times4$  factorial arrangement in a completely randomized design [14]. The Tukey test was employed for post hoc testing wherein the difference was considered to be statistically significant when the value of p was less than 0.05 (p<0.05) in the experimental units, with the main effects for separating treatments to be AC, CP, MS, SC, as well as the LWT and the HWT with individual sow and her litter as the repeated experimental unit.

- 120
- 121

## Results

122 Temperature and Humidity Index

The temperature in the pig's pen ranged from 24.71 to 31.49°C at a THI of 74 to 84°C, with AC and CP treatments showing lower scores ranging from 74.7 to 77.0°C in comparison to the scores observed under SC and MS treatments which were higher than 80°C (**Fig. 1**).

126

#### 127 Behavior Characteristics

- 128 A significant increase in drinking, standing, and position change behaviors was observed when sows were under
- 129 SC and MS (p<0.05) treatments in comparison to when they were under AC and CP treatment (Table 1). Lying
- behavior was higher in sows under AC and CP (p < 0.05) treatments than the ones under SC and MS treatments.
- 131

#### 132 **Reproductive Performances**

- 133 During lactation, the sows under AC and CP treatments showed a significantly higher ADFI (p < 0.05) than the ones
- under SC and MS treatments. The LWT (p < 0.05) treatment also showed an increase in ADFI as compared to the HWT treatment (Table 2).
- 136 The weaning weight of piglets under the AC and CP treatment was higher (p < 0.05) than that of piglets under SC
- 137 and MS treatment (Table 3). Also, there was an increase (p<0.05) in piglet weight during weaning in LWT as
- 138 compared with the weaning weight of the ones under HWT.
- 139

#### 140 Hair Cortisol, Respiratory Rate, and Rectal Temperature

141 Sows under SC and MS treatment had significantly higher cortisol levels (p < 0.05) than those under AC and CP

142 treatment (Fig. 2). Cortisol levels were also higher under HWT treatment (p<0.05) than those under LWT treatment.

143 During the exposure to HS, the rectal temperature and respiratory rate of the sows was recorded, as illustrated in Fig.

144 **3 and Fig. 4.** As a result, the rectal temperature and respiratory rate were found to be significantly higher under SC

- 145 and MS treatment than those under AC and CP treatment (p < 0.05).
- 146
- 147

# Discussion

The results for THI in this experiment is in accordance with a previous study where the air cooling systems showed a reduced THI ranging from 76.1-80.1 as compared to that (THI: 79.6-82.9) under water drip [15]. A satisfactory range of THI for pigs should be less than 74 [8]. A THI value of 75-78 is considered as mild whereas a value of 79-83 is considered to be dangerous for pigs [15,16]. If the THI value is equal to or greater than 84, it is an emergency condition for pigs [17]. As per our results, the THI values under AC and CP treatment ranged from 74.7 to 77.0, which indicates that the AC and CP treatments are preferable over SC and MS in order to mitigate HS. The increased drinking behavior of sows under SC and MS treatments supported a previous report, according to

155 which the floor cooling pads decreased drinking behavior in lactating sows [18]. During HS, Sows are generally

156 known to consume more water to retrieve the water loss [11] and to reduce discomfort due to body temperature [19],

157 as compared to the consumption under thermoneutral conditions. It is possible that an effective cooling system 158 would minimize the drinking behavior in the sows during extreme heat. The results of this study demonstrated that 159 the AC and CP treatments led to alleviation of drinking behavior in lactating sows.

Lactating sows under SC and MS treatment spent more time standing and changed position at intervals, but showed an opposite trend under the AC and CP treatments, wherein the sows spent more time lying. Results similar to AC and CP treatments were observed while using a cooling pad system for lactating sows [18]. The drip cooling system proposed by Barbari et al. [20] also showed similar results of increase in lying behavior and decrease in position alteration. Lying behavior requires less energy, and the posture allows more heat exchange between the sows and the cooling system, which helps to reduce the adverse effects of HS [21]. Therefore, the use of AC and CP is beneficial because it uses the advantage of thermoregulatory behavior.

167 In our results, the ADFI was higher in lactating sows under AC, CP, and LWT treatments. Food consumption by 168 lactating sows has always been a challenge during HS, presumably to alleviate the metabolic heat produced during 169 digestion [22,23,26], which clarifies the reason for lower feed consumption in sows during HS and an increase in 170 feed intake at a favorable thermal zone. HS can also alter milk yield by compromising dietary nutrients in the 171 mammary gland through peripheral vasodilation [23,25]. A report by Ribeiro et al. [7] says that temperature above 172 the thermal comfort zone negatively affects feed intake and milk production in lactating sows and the weaning 173 weight in piglets. The improved thermal condition of sows in our study explains the reason for an increase in piglet 174 weight at weaning under AC, CP, and LWT treatment because HS must be sufficiently reduced in sows to reach a 175 unique potential for minimizing weight loss and improving milk production during lactation [24,25]. We agree that 176 AC, CP, and LWT systems helped to alleviate HS in lactating sows.

The lower cortisol levels in our study are in agreement with a recent study from our laboratory, which says that the cortisol levels in the lactating sows are lower under the AC and CP treatment than under the SC and MS treatments [27]. Cortisol levels increase as the environmental temperature increases [27,28,29]. We hypothesized that the optimum environment for a sow is determined by the appropriate temperature provided to them under AC and CP systems, as well as under the LWT treatment. Therefore, these systems/treatments help to minimize HS thereby reducing the cortisol levels in the sows.

Rectal temperature is higher in sows particularly during lactation due to an increase in their metabolic activities that generate more heat, leading to an increase in respiratory rate [27,30,31]. The lower rectal temperature and respiratory rate observed by employing the AC, CP, and LWT systems is consistent with the report by Watanabe et al. [32], according to which a reduced rectal temperature and respiratory frequency is observed in farrowing sows

188 with cold water (10 or 15°C) decreases respiration rate and rectal temperature by about 20% and 0.8%, respectively.

while using an evaporative cooling system. A similar study by Jeon et al. [33] reported that supplying lactating sows

189 Therefore, we opined that the rectal temperature and respiratory rate is alleviated in pigs under HS by the AC, CP,

190 and LWT treatments, which minimizes the overall heat production.

191

187

#### 192 CONCLUSION

193 The use of AC and CP systems decreased drinking, standing, and position change times, and increased 194 lying behavior as compared to the sows under SC and MS systems. The AC and CP treatments led to an increase in 195 ADFI in sows and piglet weight at weaning in comparison to the SC and MS treatments. LWT also led to an 196 increase in ADFI levels in sows during lactation and piglet weight at weaning. Hair cortisol levels, rectal 197 temperature, and respiratory rates were reduced in sows under the AC, CP and LWT systems in comparison to the 198 the SC, MS, and HWT systems. However, the LWT had no significant effect on the drinking, standing, position 199 change times, and lying behaviors of the sows. Therefore, based on behavioral responses in sows during HS, we 200 recommend the use of AC and CP systems for better performance of sows.

201

202

# Acknowledgments

The Authors of this project appreciates the support fund received by the Rural Development Administration
Research Project (Detailed Task Number: PJ01479601) and the 2021 Rural Development Administration, National
Institute of Livestock Science, Republic of Korea.

207		References
208 209 210	1	Huddart D, Stott T. Climate Change and Adventure Tourism. In Adventure Tourism. Palgrave Mac Cham. 2020; 437-469. DOI: <u>https://doi.org/10.1007/978-3-030-18623-4_13</u>
211 212	2	Johnson JS. Heat stress: impact on livestock well-being and productivity and mitigation strategies to alleviate the negative effects. Anim Prod Sci. 2018; 58:1404-1413. DOI: <u>https://doi.org/10.1071/AN17725</u>
213 214 215	3	Renaudeau D, Collin A, Yahav S, De Basilio V, Gourdine JL, Collier, RJ. Adaptation to hot climate and strategies to alleviate heat stress in livestock production. Anim. 2012; 6:707–728. DOI: <u>https://doi.org/10.1017/s1751731111002448</u>
216 217 218	4	Kim K, Hosseindoust A, Choi Y, Kim M, Lee J, Kim T, Chae B. Hot-Melt Extruded Selenium: a Highly Absorbable Nano-Selenium in Lactating Sows Exposed to High Ambient Temperature. Biol Trace Element Res. 2021; 199:3345-53. DOI: <u>https://doi.org/10.1007/s12011-020-02459-3</u>
219 220	5	Ross JW, Hale BJ, Gabler NK, Rhoads RP, Keating AF, Baumgard LH. Physiological consequences of heat stress in pigs. Anim Prod Sci. 2015; 55:1381-1390. DOI: <u>https://doi.org/10.1071/AN15267</u>
221 222	6	Nardone A, Ronchi B, Lacetera N, Bernabucci U. Climatic effects on productive traits in livestock. Vet Research Comm. 2006; 30:75-81. DOI: <u>http://10.1007/s11259-006-0016-x</u>
223 224 225	7	Ribeiro BP, Lanferdini E, Palencia JY, Lemes MA, de Abreu ML, de Souza Cantarelli V, Ferreira RA. Heat negatively affects lactating swine: A meta-analysis. Journal of Therm Bio. 2018; 74:325-330. DOI: <u>https://doi.org/10.1016/j.jtherbio.2018.04.015</u>
226 227 228	8	Wegner K, Lambertz C, Das G, Reiner G, Gauly M. Effects of temperature and temperature-humidity index on the reproductive performance of sows during summer months under a temperate climate. Anim Sci Journ. 2016; 87:1334–1339. DOI: <u>https://doi.org/10.1111/asj.12569</u>
229 230	9	Lucy MC, Safranski TJ, Rhoades JN, Ross JW, Gabler NK, Rhoads RP, Baumgard L.H. Effect of heat stress (HS) on thermal regulation during pregnancy in first parity sows. Journ of Anim Sci. 2012; 90:574.
231 232	10	National Research Council (NRC), "Nutrient Requirements of swine", Livestock Production Science. Nap 2012; 208-238.
233 234 235	11	Zhu Y, Johnston LJ, Reese MH., Buchanan ES, Tallaksen JE, Hilbrands AH, Li YZ. Effects of cooled floor pads combined with chilled drinking water on behaviour and performance of lactating sows under heat stress. Journ of amim sci. 2021;99: skab066. DOI: <u>https://doi.org/10.1093/jas/skab066</u>
236 237 238	12	Dikmen SE, Alava E, Pontes E, Fear JM, Dikmen BY, Olson TA, Hansen PJ. Differences in thermoregulatory ability between slick-haired and wild-type lactating Holstein cows in response to acute heat stress. Journ of dairy sci. 2008; 91:3395-3402. DOI: https://doi.org/10.3168/jds.2008-1072

- Ataallahi M, Nejad J G, Takahashi J, Song Y, Sung K, Yun J, Park K. Effects of environmental changes during different seasons on hair cortisol concentration as a biomarker of chronic stress in Korean native cattle. Int Journ of Agric and Biol. 2019; 21:1166-1172. DOI: <u>http://doi.org/10.17957/IJAB/15.1007</u>
- 242 14 SAS. 2012, SAS Software for PC. Release 9.3, SAS Institute. Ins, Cart, NC, USA.
- Jeong Y, Choi Y, Kim D, Min Y, Cho E, Kim J. Effects of cooling systems on physiological responses and intestinal microflora in early gestating sows exposed to high-temperature stress. Journ of Anim Sci and Technology. 2021; 63:904. DOI: <u>https://doi.org/10.5187/jast.2021.e79</u>
- Haeussermann A, Hartung E, Jungbluth T, Vranken E, Aerts JM, Berckmans D. Cooling effects and evaporation characteristics of fogging systems in an experimental piggery. Biosys Eng. 2007; 97:395–405.
  DOI: <u>http://dx.doi.org/10.1016/j.biosystemseng.2007.03.019</u>
- Vega RS, Lontoc CA, Punay LC, Cajano PJ. Comparative performance of sows housed with and without evaporative cooling system at temperature humidity index of 73-83. Philip Journ of Vet and Anim Sci. 2016; 42:1-1
- Parois SP, Cabezón FA, Schinckel AP, Johnson JS, Stwalley RM, Marchant-Forde JN. Effect of floor cooling
  on behavior and heart rate of late lactation sows under acute heat stress. Frontiers in Veterinary Science.
  2018:223. DOI: https://dx.doi.org/10.3389%2Ffvets.2018.00223
- Pang ZZ, Li BM, Zheng WC, Lin BZ, Liu ZH. Effects of water-cooled cover on physiological and production parameters of farrowing sows under hot and humid climates. Int J Agric Biol Eng. 2016; 9:178–84. DOI: http://dx.doi.org/10.3965/j.ijabe.20160904.1858
- 20 Barbari M, Bianchi M, Guerri FS. Preliminaryanalysis of Different Cooling Systems of Sows In Farrowing
   Room. Journ of Agric Eng. 2007; 38:53-8. DOI: <u>https://doi.org/10.4081/jae.2007.1.53</u>
- Huynh TT, Aarnink AJ, Spoolder HA, Verstegen MW, Kemp B. Effects of floor cooling during high ambient temperatures on the lying behavior and productivity of growing finishing pigs. Transactions of the ASAE. 2004; 47:1773. DOI: <u>http://dx.doi.org//10.13031/2013.17620</u>
- 263 22 Baumgard LH, Rhoads Jr RP. Effects of heat stress on postabsorptive metabolism and energetics. Annu. Rev.
   264 Anim. Biosci. 2013; 1:311-37. DOI: <u>https://doi.org/10.1146/annurev-animal-031412-103644</u>
- Renaudeau D, Noblet J, Dourmad JY. Effect of ambient temperature on mammary gland metabolism in lactating sows. Journ of Anim Sci. 2003; 81:217-31. DOI: <u>https://doi.org/10.2527/2003.811217x</u>
- 267 24 Cabezon FA, Schinckel AP, Marchant-Forde JN, Johnson JS, Stwalley RM. Effect of floor cooling on late 268 lactation sows under acute heat stress. Livestock Science. 2017; 206:113-20. DOI: 269 https://doi.org/10.1016/j.livsci.2017.10.017
- 25 Mendes M F, Oliveira DH, Cruz FL, Mendes MA, Ribeiro BP, Ferreira RA. Evaporative cooling system for gestating and lactating sows: a systematic review. Ciência Rural. 2020; 50. DOI: <u>https://doi.org/10.1590/0103-</u>

### 272 <u>8478cr20190830</u>

- 26 Choi Y, Moturi J, Hosseindoust A, Kim M, Kim K, Lee J, Song C, Kim Y, Chae B. Night feeding in lactating
  274 sows is an essential management approach to decrease the detrimental impacts of heat stress. Journal of Animal
  275 Science and Technology. 2019; 61:333. DOI: https://doi.org/10.5187/jast.2019.61.6.333
- 27 Oh SM, Choi YH, Kim DW, Ha SH, Kim JE, Jung HJ, Kim JS. Effect of Different Cooling System on
  277 Performance and Hair Cortisol on Sows under Heat Stress. Journ of the Kor Academ-Indust coop Soc.
  278 2021; 22:160-168. DOI: https://doi.org/10.5762/KAIS.2021.22.3.160
- 28 Elez D, Vidović S, Matić G. The influence of hyperthermic stress on the redox state of glucocorticoid receptor.
   280 Stress. 2000; 3:247-55. DOI: <u>http://dx.doi.org/10.3109/10253890009001129</u>
- 281 29 Mahmoud KZ, Edens FW, Eisen EJ, Havenstein GB. Ascorbic acid decreases heat shock protein 70 and plasma corticosterone response in broilers (Gallus gallus domesticus) subjected to cyclic heat stress. Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology. 2004;137:35-42.DOI: http://doi.org/10.1016/j.cbpc.2003.09.013
- Williams AM, Safranski TJ, Spiers DE, Eichen PA, Coate EA, Lucy MC. Effects of a controlled heat stress during late gestation, lactation, and after weaning on thermoregulation, metabolism, and reproduction of primiparous sows. J Anim Sci. 2013;91:2700-14. DOI: <u>https://doi.org/10.2527/jas.2012-6055</u>
- Kim K, Choi Y, Hosseindoust A, Kim M, Hwang S, Bu M, Lee J, Kim Y, Chae BJ. Evaluation of high nutrient diets and additional dextrose on reproductive performance and litter performance of heat-stressed lactating sows. Anim Sci Journ. 2019; 9:1212-9. DOI: https://doi.org/10.1111/asj.13214
- Watanabe PH, Azevedo TA, do Nascimento Silva MA, Oliveira NM, Gomes TR, Andrade TS, Barbosa Filho
   JA. Cooling ventilation at farrowing for sows from first to third parturition. Comunicata Scientiae. 2018; 9:556 64. DOI: https://doi.org/10.14295/cs.v9i4.1098
- Jeon JH, Yeon SC, Choi YH, Min W, Kim S, Kim PJ, Chang HH. Effects of chilled drinking water on the
   performance of lactating sows and their litters during high ambient temperatures under farm conditions. Livest
   Scie. 2006; 105:86-93. DOI:<u>https://doi.org/10.1016/j.livsci.2006.04.035</u>
- 297

## 

# 

# 301 Table 1. Chemical composition of experimental diet

Item	Basal lactation diet				
Calculated composition	%				
ME, kcal/kg	3.350				
СР	20.13				
Ca	0.75				
Av. P	0.32				
Lys	1.15				
Met + Cys	0.72				

Tables

Me, metabolizable energy; CP, crude protein; Ca, calcium; P, phosphorus; Lys, Lysine; Met, methionine; Cys, cysteine.

## Table 2. Effects of cooling system and drinking temperature on behavior characteristics of sows

Ű

305

304

201
3116

Item 1	Coling system			DWT (°C)		SEM <sup>1</sup>		<i>p</i> -value		
Item 2 (°C)	AC	CP	SC	MS	LWI	HWI		C	D	$C \times D$
Drinking	5.86 <sup>b</sup>	6.06 <sup>b</sup>	10.39ª	10.2ª	8.18	8.10	0.21	<0.01	0.621	0.868
Standing, %	9.51 <sup>b</sup>	9.82 <sup>b</sup>	14.18ª	15.1ª	12.26	12.06	0.46	<0.01	0.539	0.483
Position change, times	5.38 <sup>b</sup>	5.98 <sup>b</sup>	8.21ª	8.31ª	7.01	6.93	0.25	<0.01	0.634	0.995
Lying, %	80.80ª	80.90ª	75.21 <sup>b</sup>	74.9 <sup>b</sup>	77.87	78.06	0.64	<0.01	0.69	0.719
Sitting, %	9.68	9.28	10.60	9.92	9.86	9.88	0.53	0.111	0.961	0.753
Feeding	0.58	0.63	0.60	0.59	0.61	0.59	0.02	0.181	0.462	0.923
Nursing	1.96	1.84	1.83	1.96	1.97	1.83	0.19	0.859	0.333	0.499

<sup>1</sup>SEM, standard error of means; DWT, drinking water temperature; AC, air conditioner; CP, cooling pad; SC, snout cooling; MS, mist spray; LWT : low water temperature; HWT, high water temperature; C, cooling system; D, drink temperature. 

307

308

## 

#### Table 3. Effects of cooling system and drink temperature on sow performance

ltem 1	Coling system			DWT(°C)			<i>p</i> -value			
Item 2 (°C)	AC	СР	SC	MS	LWT	HWT	SEM <sup>1</sup>	С	D	$C \times D$
ADFI, kg/d										
During lactation	5.76ª	5.70 <sup>a</sup>	5.26 <sup>b</sup>	5.26 <sup>b</sup>	5.73ª	5.26 <sup>b</sup>	0.07	<0.001	<0.001	0.973
Weaning to estrus interval, d	4.75	4.91	5.42	5.67	5.00	5.37	0.38	0.072	0.172	0.966
BW, kg										
d 112	255.8	253.6	256.7	254.7	255.7	254.7	5.96	0.957	0.798	0.989
Weaning	222.7	219.9	223.1	220.5	222.5	220.5	6.09	0.938	0.641	0.988
Loss during Lactation	33.1	33.7	33.6	34.3	33.2	34.1	0.76	0.527	0.09	0.933
BF, mm										
d 112	20.7	20.5	20.2	20.8	20.6	20.5	0.37	0.427	0.728	0.284
Weaning	15.9	15.4	14.9	15.4	15.6	15.2	0.39	0.13	0.16	0.467
Loss during Lactation	4.83	5.03	5.31	5.36	4.98	5.29	0.23	0.109	0.071	0.892

<sup>1</sup>SEM, standard error of means; DWT; drinking water temperature; AC, air conditioner; CP, cooling pad; SC, snout cooling; MS, mist spray; LWT : low water temperature; HWT, high water temperature; C, cooling system; D, drink temperature; BW, body weight; BF, backfat thickness; ADFI, average daily feed intake.



Fig. 1. Temperature-humidity index during experimental period



**Fig. 2.** Effects of cooling system and drink temperature on hair cortisol in sow. a, b means with different superscript on the bar differ significantly (p<0.05); AC, air conditioner; CP, cooling pad; SC, snout cooling; MS, mist spray; LWT, low water temperature; HWT, high water temperature; C, cooling system.



Fig. 3. Effects of cooling system and drink temperature on rectal temperature of sows. Asterisks (\*) Means with different superscript on the bar differs significantly (p<0.05) where Snout cooling and Mist spray are higher in rectal temperature compared with Air condition and Cooling pad treatment; L means at low water temperature; H, at high water temperature



Fig. 4. Effects of cooling system and drink temperature on respiratory rate of sows. Asterisks (\*) Means with different superscript on the bar differs significantly (p<0.05) where Snout cooling and Mist spray are higher in respiratory rate compared with Air condition and Cooling pad treatment; L means at low water temperature; H, at hight water temperature.