

**JAST (Journal of Animal Science and Technology) TITLE PAGE**

**Upload this completed form to website with submission**

ARTICLE INFORMATION	Fill in information in each box below
<b>Article Type</b>	Research article
<b>Article Title (within 20 words without abbreviations)</b>	Feeding dietary non-starch polysaccharides supplemented with xylanase could improve the performance of broilers
<b>Running Title (within 10 words)</b>	Growth performance in response to dietary NSP in broiler
<b>Author</b>	Venuste Maniraguha <sup>1,a</sup> , Jun Seon Hong <sup>1,a</sup> , Myunghwan Yu <sup>1</sup> , Elijah Ogola Oketch <sup>1</sup> , Young-Joo Yi <sup>2</sup> , Hyeonho Yun <sup>3</sup> , Dinesh D. Jayasena <sup>4</sup> , Jung Min Heo <sup>1,*</sup>  <sup>a</sup> Venuste Maniraguha and Jun Seon Hong contributed equally
<b>Affiliation</b>	<sup>1</sup> Department of Animal Science and Biotechnology, Chungnam National University, Daejeon 34134, Korea  <sup>2</sup> Department of Agricultural Education, College of Education, Suncheon National University, Suncheon 57922, Korea  <sup>3</sup> CJ CheilJedang BIO, Seoul 04560, Korea  <sup>4</sup> Department of Animal Science, Uva Wellassa University of Sri Lanka, Badulla 90000, Sri Lanka
<b>ORCID (for more information, please visit <a href="https://orcid.org">https://orcid.org</a>)</b>	Venuste Maniraguha ( <a href="https://orcid.org/0000-0002-5615-3036">https://orcid.org/0000-0002-5615-3036</a> )  Jun Seon Hong ( <a href="https://orcid.org/0000-0003-2142-9888">https://orcid.org/0000-0003-2142-9888</a> )  Myunghwan Yu ( <a href="https://orcid.org/0000-0003-4479-4677">https://orcid.org/0000-0003-4479-4677</a> )  Elijah Ogola Oketch ( <a href="https://orcid.org/0000-0003-4364-460X">https://orcid.org/0000-0003-4364-460X</a> )  Young-Joo Yi ( <a href="https://orcid.org/0000-0002-7167-5123">https://orcid.org/0000-0002-7167-5123</a> )  Hyeonho Yun ( <a href="https://orcid.org/0009-0001-3215-822X">https://orcid.org/0009-0001-3215-822X</a> )  Dinesh Darshaka Jayasena ( <a href="https://orcid.org/0000-0002-2251-4200">https://orcid.org/0000-0002-2251-4200</a> )  Jung Min Heo ( <a href="https://orcid.org/0000-0002-3693-1320">https://orcid.org/0000-0002-3693-1320</a> )

<b>Competing interests</b>	The author Hyeonho Yun is an employee of CJ CheilJedang BIO and declares no conflicts of interest.
<b>Funding sources</b> State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available.	This work was supported by the CJ CheilJedang BIO, Seoul 04560, Korea
<b>Acknowledgements</b>	The authors are grateful to CJ CheilJedang BIO, Seoul 04560, Korea, for the financial support and supply of xylanase enzyme used in this study.
<b>Availability of data and material</b>	Upon reasonable request, the datasets of this study can be available from the corresponding author.
<b>Authors' contributions</b> Please specify the authors' role using this form.	Conceptualization: Hong JS, Yun H and Heo JM Data curation: Maniraguha V, Hong JS and Heo JM Formal analysis: Yi YJ Methodology: Hong JS and Yu M Software: Hong JS Validation: Hong JS, Yu M, Jayasena DD and Heo JM Investigation: Yu M and Oketch EO Writing - original draft: Maniraguha V Writing - review & editing: Maniraguha V, Hong JS, Yu M, Oketch EO, Yi YJ, Yun H, Jayasena DD and Heo JM
<b>Ethics approval and consent to participate</b>	The experimental protocol and procedures for this study were reviewed and approved by the Animal Care and Use Committee of Chungnam National University (Protocol Number; 202103A-CNU-029).

5 **CORRESPONDING AUTHOR CONTACT INFORMATION**

<b>For the corresponding author (responsible for correspondence, proofreading, and reprints)</b>	<b>Fill in information in each box below</b>
First name, middle initial, last name	Jung Min Heo
Email address – this is where your proofs will be sent	<u>jmheo@cnu.ac.kr</u>
Secondary Email address	None
Address	Department of Animal Science and Biotechnology, Chungnam National University, Daejeon 34134, Republic of Korea
Cell phone number	
Office phone number	+ 82 42-821-5777
Fax number	

6

ACCEPTED

7 **Abstract**

8           The impact of dietary non-starch polysaccharides (NSP) on performance and carcass traits of broilers fed  
9 wheat-bran substituted into corn-soybean meal-based diets supplemented with xylanase was investigated. A total of  
10 280 (7-day-old) Ross 308 broilers were randomly allotted to one of five dietary treatments with 8 replicates, 7 chicks  
11 per pen. Treatments were; i) CON: Control diet, ii) CON-X (CON + 3,000 U/kg xylanase), iii) L-X: low NSP (2%  
12 wheat bran in CON + 3,000 U/kg xylanase), iv) M-X: medium NSP (4% wheat bran in CON + 3,000 U/kg xylanase),  
13 v) H-X: higher NSP (8% wheat bran in CON+ 3,000 U/kg xylanase). Birds fed the H-X diet increased ( $p < 0.05$ ) daily  
14 gains, and average daily feed intake and had marginally improved body weights ( $p = 0.074$ ) on day 35. Relatively, the  
15 H-X diet tended to increase the average daily gains ( $p = 0.053$ ;  $p = 0.073$ ) of birds during the grower phase (d 24-35)  
16 and the entire experimental period (d 8-35), respectively. Moreover, there were no significant differences among  
17 treatments in the feed conversion ratio of birds throughout the entire experiment period. Birds fed diets CON-X, L-X,  
18 and M-X had improved ( $p < 0.05$ ) the ileal digestibility of energy on d 24 and 35 compared to those fed the H-X diet.  
19 Furthermore, birds fed diet CON-X improved ( $p < 0.05$ ) N digestibility on d 24. Improved carcass moisture content  
20 and lowered crude fat of leg meat ( $p < 0.05$ ) were noted in birds fed the diet M-X and H-X on d 35, respectively. The  
21 intestinal viscosity was reduced ( $p < 0.05$ ) in xylanase-supplemented treatments CON-X, L-X, M-X, and H-X diets  
22 when compared to CON. Our results suggest that supplementing 3,000 U/kg xylanase in a higher NSP (8% wheat bran  
23 substituted level) diet could improve the intestinal viscosity and growth performance of broilers.

24 **Keywords:** Broiler, digestibility, non-starch polysaccharide, wheat bran, xylanase

25

## Introduction

The provision of nutrient-dense diets that aim to utilize the full genetic potential of modern fast-growing broilers is a costly affair. It is well known that feeds constitute the largest portion of farm expenditure [1, 2]. Thus, it has been a fixation for nutritionists to investigate strategies to lower feed costs while including other feed alternative resources and additives that might be more sustainable and relatively inexpensive. In the face of high feed costs and erratic supply, wheat bran has been considered an alternative energy source in poultry diets because of its higher starch level and reasonable N content [3, 4].

Nevertheless, the maximum utilization of nutrients from wheat bran by monogastric is constrained by the presence of anti-nutritive, water-soluble non-starch polysaccharides[5] in the form of arabinoxylan and beta-glucan fractions that could have detrimental impacts on the productive performance of broilers [6]. The anti-nutritive effect of water-soluble NSP is attributed to the increase in the viscosity of the intestinal digesta. Increased digesta viscosity has been reported to impede enzymatic digestion by endogenous enzymes, resultantly limiting nutrient digestion and bioavailability [7]. However, dietary non-starch polysaccharides (NSP) have received attention for their capacity to elicit positive impacts on broiler performance and normal gut function [8, 9]. As a result, monogastric diets are formulated with a 2-3% fiber level to strike a balance between the deleterious effects of high fiber in diets and the desired gut-modulating effects of dietary fiber.

It is also well-established that specific enzyme supplementation in the form of xylanase could improve the nutritional value of dietary fiber. Xylanase works to degrade arabinoxylans, and in turn, reduce chyme viscosity, facilitate enzymatic nutrient digestion, and potentially release xylo-oligosaccharides that could induce beneficial prebiotic effects to the gut. Xylanase supplementation in higher dietary NSP diets has also been reported to improve the performance of broilers by increasing the availability of nutrients [6, 10]. Nonetheless, the degradation ability of xylanase differs with the inclusion levels of dietary NSP in the diet. It is imperative to evaluate the effect of xylanase supplementation on broilers fed different NSP levels to determine the optimal inclusion level for improved performance. In this study, we hypothesized that feeding broilers with a diet containing higher dietary NSP levels supplemented with xylanase would lower intestinal viscosity, improve nutrient utilization, and improve broiler performance.

## Materials and Methods

The experiment was conducted at the Daejeon Poultry Research Unit of Chungnam National University, Daejeon, Korea. The rearing practices followed the recommendations of the Ross 308 Specification Management Guide [11]. The experimental protocol and procedures for the study were reviewed and approved by the Animal Care and Use Committee of Chungnam National University (Protocol Number; 202103A-CNU-029). The novel xylanase enzyme used in the current study was supplied by CJ CheilJedang BIO, Seoul, South Korea.

### Animals and housing

A total of 280 one-day-old Ross 308 were purchased from a commercial hatchery to be used in the current study. The birds were allowed a one-week adaptation period and then randomly allocated based on the average body weight ( $115.01\text{g} \pm 1.55$ ) on day 8. Seven birds were housed in each raised wire floor enriched cage ( $85 \times 55 \times 35 \text{ cm}^3$ ). The room was lit continuously during the whole experimental period and the room temperature started at  $32^\circ\text{C}$  for the first day and lowered by  $2^\circ\text{C}$  per week until a final temperature of  $28^\circ\text{C}$  on day 35. Each pen was equipped with three nipple drinkers and a metal feeding trough for the *ad libitum* access to clean drinking water and feed.

### Experimental design and dietary treatments

Birds were assigned to one of five dietary treatments with 8 replicates per treatment in a completely randomized design. Following the Ross 308 Nutritional Specifications, a corn-soybean meal basal diet was formulated for the five dietary treatments with different NSP levels (i.e. low, medium, and high NSP) using wheat bran and supplemented with xylanase. The mash diets were as follows; i) CON: Control diet, ii) CON-X (CON + 3000 U/kg xylanase), iii) L-X: low NSP (corn replaced to 2% wheat bran in CON + 3000 U/kg xylanase), iv) M-X: medium NSP (corn replaced to 4% wheat bran in CON + 3000 U/kg xylanase), v) H-X: high NSP (corn replaced to 8% wheat bran in CON+ 3000 U/kg xylanase). Chromium oxide powder (99.9% purity, Sigma-Aldrich, St. Louis, MO, USA) was added as an external indigestible marker at the proportion of 0.3% in all experimental diets.

In addition, diet samples were collected and analyzed for the gross energy, and crude protein to verify closeness to the standard values. NSP fractions in dietary treatments were balanced using the dietary NSP profile estimator (iNSPect: inspect.canadianbio.com; Canadian Bio-system). The experimental diets were provided in two distinct phases, the starter phase (8 to 24 days) and the grower phase (25 to 35 days). The dietary composition of the

80 experimental diet utilized during all phases is illustrated in (Table 1). The amount of xylanase unit in the experimental  
81 diet was determined using Xylazyme AX tablets (Megazyme International Ltd., Ireland) according to the method  
82 described by Wickramasuria [12]. One xylanase unit is defined as the amount of enzyme that liberates 1 $\mu$ mol of xylose  
83 per minute under the conditions of the assay (pH 5.3 and 50°C). The assay is based on a 5-minute hydrolysis of xylan  
84 substrate which is stopped by the addition of dinitrosalicylic acid.

85

### 86 **Growth performance evaluation**

87 The body weight and feed consumption of birds were recorded weekly. Using the feed consumed and  
88 recorded body weights, the average daily gain (ADG), mortality-corrected average daily feed intake (ADFI), and the  
89 feed conversion ratio (FCR) to depict the efficiency of converting feed supplied to lean muscle was conducted.

90

### 91 **Nutrient digestibility, viscosity, and carcass characteristics**

92 One bird per cage (closer to the mean body weight) was selected and sacrificed by cervical dislocation for  
93 ileal digesta sample collection on days 24 and 35. Following the procedure of Wickramasuriya et al. [13], the collected  
94 samples were pooled and stored at -20°C until further analysis for nutrient digestibility and viscosity. The contents of  
95 the chromium oxide, crude protein, and energy contents of the feed and digesta samples were analyzed according to  
96 the standard procedures [14]. The apparent ileal digestibility of gross energy and crude protein was determined to  
97 estimate the rate of nutrient disappearance at the terminal ileum using the following equation stipulated by Oketch et  
98 al. [15].

99 Digestibility % =  $100 - \left[ 100 \times \left( \frac{M_{\text{diet}} \times N_{\text{digest}}}{M_{\text{digest}} \times N_{\text{diet}}} \right) \right]$ , whereby,

100  $M_{\text{diet}}$  is the marker concentration in the diet,  $N_{\text{digest}}$  is the nutrient concentration in ileal digesta,  $M_{\text{digest}}$  is the  
101 marker concentration in ileal digesta, and  $N_{\text{diet}}$  is the nutrient concentration in the diet. For carcass characteristics  
102 analysis, samples of leg meat were collected from each selected bird per cage on day 35. Thereafter, the proximate  
103 analysis of leg meat samples was conducted to determine the composition of moisture content, crude protein, crude  
104 fat, and ash according to the standard methods of AOAC [14].

105

### 106 **Statistical analyses**

107 All data for growth performance, digestibility, and carcass characteristics with respect to the treatments were  
108 analyzed by using the general linear model (GLM) of the SPSS 26.0 (SPSS Inc., Chicago, IL, USA). The pen was  
109 used as the experimental unit for the assessment of the growth performance parameters such as body weight (BW),  
110 average daily gain (ADG), average daily feed intake (ADFI), and feed conversion ratio (FCR). The individual bird  
111 was used as the statistical unit for ileal digestibility, digesta viscosity, and carcass characteristics. The results were  
112 presented by mean values and the standard error of the mean (SEM). All statements of significance were declared on  
113 a probability value of less than 0.05. Marginal effects were measured at  $0.05 < p < 0.10$ . Significant means were  
114 separated using Tukey's Multiple Range Test.

115

116

## Results

### 117 Growth performance

118 Birds fed the highest NSP levels plus xylanase in the H-X diet recorded marginally higher BW ( $p = 0.074$ )  
119 on day 35 compared to birds fed with other diets (Table 2). Feeding diet H-X increased ( $p < 0.05$ ) ADG of birds on  
120 day 35. Moreover, marginal improvements of the ADG ( $p = 0.053$ ;  $p = 0.073$ ) were noted from birds fed the H-X diet  
121 during the grower phase (day 24-35) and for the entire experimental period (day 8-35), respectively. Birds fed the H-  
122 X diet also increased ( $p < 0.05$ ) ADFI on day 35. Neither significant effects nor trends were noted for the FCR of the  
123 birds over the entire experimental period.

124

### 125 Nutrient digestibility and viscosity

126 Regarding the ileal digestibility of nutrients and energy, birds fed on diet CON-X, L-X, and M-X improved  
127 ileal digestibility of energy ( $p < 0.05$ ) on days 24 and 35 than birds fed the H-X diet (Table 3). However, an increase  
128 in protein digestibility ( $p < 0.05$ ) was only noted at day 24 within birds fed on diet CON-X when compared to the  
129 other treatments. Feeding different levels of NSP did not show any increase in protein digestibility on day 35.  
130 Considering the ileal digesta viscosity, reductions were noted ( $P < 0.05$ ) in birds fed the H-X diet when compared to  
131 the other diet-fed birds (Table 3). There was a general reduction in digesta viscosity in the xylanase-supplemented  
132 diets when compared to the non-supplemented CON diet.

133

### 134 Carcass composition



135 Data showing the effects of different dietary NSP levels on carcass composition of broilers are summarized  
136 in Table 4. Feeding M-X and H-X diets had improved the moisture content ( $p < 0.05$ ) of leg meat on day 35 than other  
137 treatments. Furthermore, birds fed on diet H-X lowered the crude fat content of leg meat on day 35 ( $p < 0.05$ ). However,  
138 feeding different NSP levels showed no effects on crude protein and ash content of leg meat.

ACCEPTED

139 **Discussion**

140 Dietary non-starch polysaccharides (NSPs) have gained more interest in monogastric feed formulation due  
141 to their influence on nutrient availability, digesta viscosity, gut health function, and overall productive performance.  
142 The current study investigated the impact of dietary NSPs supplemented with xylanase on the performance and  
143 nutrient utilization of broilers. It was hypothesized that feeding broilers with a diet containing higher dietary NSP  
144 content supplemented with xylanase would satisfy a necessary enzyme-substrate concentration and in turn, enhance  
145 the broiler's performance. As hypothesized, body weights, average daily gain, and daily feed intake increased when  
146 birds were fed a higher NSP level diet supplemented with xylanase in the present study. These results are consistent  
147 with several previous studies which indicated positive results on growth performance by feeding higher NSP diets  
148 supplemented with xylanase [6, 16]. The improvement in the broiler's performance is affected by the varying nature  
149 and the quantities of NSP present in monogastric diets, and thus careful consideration should be taken to achieve  
150 improved performance.

151 It is well documented that the total fiber content of wheat bran ranges from 36.5 to 52.4 %, with the most  
152 constituent of this fiber being NSP and the main NSPs present are arabinoxylan, cellulose, and beta-glucan at  
153 proportions of 70%, 20%, and 6% respectively [17]. Wheat bran is also known to have both insoluble and soluble  
154 fiber content which ranges from 35.0 to 48.4% and 1.5 to 4.0% respectively [18]. Considering the NSP constituents  
155 present in wheat bran, broilers are unable to optimally utilize nutrients in wheat bran due to insufficiency of  
156 endogenous NSP degrading enzymes [19]. It is well-known that, despite their adverse effects on nutrient and energy  
157 digestibility, dietary NSPs can elicit nutritional value by directly providing energy; and indirectly by improving gut  
158 health and immune function. The growth-improving capability of NSP digestion with xylanase is linked to the role  
159 played by xylanase to internally cleave the xylan molecule, releasing oligosaccharides such as xylose and arabinose;  
160 and resultantly mitigating the deleterious impacts of dietary NSPs in the diets. The improved body weights and feed  
161 intake of birds fed higher dietary NSP supplemented with xylanase has been reported elsewhere [20, 21]. A higher  
162 NSP level diet was advantageous to satisfy an ideal enzyme-substrate concentration for optimal enzymatic activity. In  
163 addition, the fraction of water-soluble NSP in the diet can function as a source of beneficial fermentable substrates for  
164 gut microbiota through selective fermentation, which promote the integrity of the intestinal lining, improving the  
165 digestibility of nutrient and energy, and reducing the ability of pathogenic bacteria to proliferate therefore improving  
166 the growth performance of birds [22, 23].

167           Moreover, nutrient digestibility in poultry is closely related to the amount of digested protein, fat, and starch  
168 that disappears at the ileum [24]. A couple of studies have demonstrated that the addition of xylanases to wheat-bran-  
169 based diets could increase nutrient digestibility by partially hydrolyzing NSPs available in wheat bran, resulting in  
170 improvements in nutrient utilization and growth performance of broilers [25, 26]. In the current study, birds fed dietary  
171 NSP supplemented with xylanase recorded improved ileal digestibility of energy and crude protein. The improvements  
172 in nutrient digestibility are caused by reduced intestinal viscosity and the breaking down of the arabinoxylan backbone  
173 into arabinose and xylose at different segments of the broiler gastrointestinal tract [3]. As earlier reported, both the  
174 nature and the level of NSP present in wheat bran may also impair endogenous amino acid release and suppress amino  
175 acid digestibility [27]. The current findings, along with those results from Esmailipour et al. [28] suggest that  
176 xylanase can reduce nutrient entrapment and increase the digestibility of mostly starch and protein at the early stages  
177 of a chick's life. It was not unusual to have no improvement in crude protein digestibility on day 35 as was also  
178 previously reported by Esmailipour et al. [28]. The efficacy of the exogenous xylanase to the diet was significantly  
179 observed in broiler chickens fed wheat bran-based diets during the early growth stage. The improvements could be  
180 attributed to the capacity of xylanase to mitigate the deleterious effects of low levels of endogenous protease that have  
181 been reported at the early stages of a chick's life that coincides with the common use of a higher crude protein diet at  
182 the starter stage as compared to the grower stage.

183           Concerning the ileal digesta viscosity, a reduction in ileal digesta's viscosity with xylanase supplementation  
184 was noted. This is in agreement with Gorenz et al. [29] who reported that xylanase supplementation in broiler diets  
185 both linearly and quadratically decreased ileal digesta viscosity. These results are also consistent with the work of  
186 Barasch and Grimes, [30] who reported that the xylanase enzyme supplementation generally reduces digesta viscosity  
187 and increases nutrient digestibility. Additionally, xylanase reduces intestinal viscosity, increases endogenous amino  
188 acid secretion, releases nutrients, and increases cell wall permeability to absorb utilizable nutrients. According to Van  
189 Hoeck et al. [20], compared to the control diets, xylanase decreased viscosity and pH at various intestinal segments;  
190 and was beneficial for the carcass traits. In this current study, xylanase supplementation significantly reduced the  
191 intestinal viscosity in broilers at both days 24 and 35 of age. The highest decrease in digesta viscosity was observed  
192 with a higher NSP diet. This reduction in intestinal viscosity was believed to be associated with available substrates  
193 of dietary NSP concerning the enzyme.

194 It was also observed that feeding dietary NSP supplemented with xylanase exerted an impact on the carcass  
195 composition of broilers in this study. Our findings showed that birds fed a higher NSP diet plus xylanase had the lowest  
196 crude fat content compared to other dietary treatments. In addition, a significant improvement in the moisture content  
197 of leg meat was noted in birds fed the M-X and H-X diet. The results of this study are in line with the findings of Alam  
198 et al. [31]. The addition of xylanase enzymes in wheat bran-based diets improved carcass yield and composition of  
199 broiler chicken and this was attributed to higher muscle development. This is consistent with another report from  
200 Williams et al. [32] which indicated a reduction in carcass crude fat content of birds when a wheat bran and corn diet  
201 was supplemented with carbohydrase enzyme. However, the present study contradicts the findings of Hussein et al.  
202 [33] who revealed that carcass yield and retail cuts chemical composition showed no significant difference with dietary  
203 xylanase supplementation. In this experiment, there were no notable major effects or marginal effects of xylanase  
204 supplementation on crude protein and ash content of leg meat. This can be attributed to the amount of energy level  
205 that was available concerning the amount of xylanase level supplemented. Another potential evidence is that feeding  
206 a fibrous diet moderates nutrient density which provides an opportunity to maintain the nutrient requirements thus  
207 resulting in reduced fat deposition in muscles [34, 35].

208 In conclusion, our results showed that the impact of dietary NSPs on broiler's performance can be alleviated  
209 by xylanase supplementation, and could differ depending on the formulated NSP proportions in the diet. In this study,  
210 it was determined that higher NSP levels in diets supplemented with 3000 U/kg xylanase led to improved body weights  
211 and feed intake of birds. The improved performance is associated with an influence of higher NSP levels to satisfy a  
212 requisite enzyme-substrate concentration that ultimately facilitates the lowering of intestinal viscosity and improves  
213 nutrient utilization. From a practical point of view, elevated dietary NSP levels from the substitution of corn with  
214 wheat bran (up to 8%) in the control diet indicated the potential of higher inclusion of alternative feed resources with  
215 specific enzyme supplementation to replace a fraction of corn in diets without compromising broiler performance.

216

217 **REFERENCES**

218

- 219 1. Samuel Olugbenga O, Olusegun Abayomi O, Adebimpe Oluseye A, Akinbowale Taiwo T. Optimized nutrients  
220 diet formulation of broiler poultry rations in nigeria using linear programming. *J Nutr Food Sci S.* 2015;14:1-6.  
221 <https://doi.org/10.4172/2155-9600.s14-002>
- 222 2. Munezero O, Kim IH. Effects of protease enzyme supplementation in weanling pigs' diet with different crude  
223 protein levels on growth performance and nutrient digestibility. *J Anim Sci Technol.* 2022;64:854-62.  
224 <https://doi.org/10.5187/jast.2022.e51>
- 225 3. Zhang L, Xu J, Lei L, Jiang Y, Gao F, Zhou G. Effects of xylanase supplementation on growth performance,  
226 nutrient digestibility and non-starch polysaccharide degradation in different sections of the gastrointestinal tract  
227 of broilers fed wheat-based diets. *Asian-Australas J Anim Sci.* 2014;27:855-61.  
228 <https://doi.org/10.5713/ajas.2014.14006>
- 229 4. Del Alamo AG, Verstegen M, Den Hartog L, De Ayala PP, Villamide M. Effect of wheat cultivar and enzyme  
230 addition to broiler chicken diets on nutrient digestibility, performance, and apparent metabolizable energy content.  
231 *Poult Sci.* 2008;87:759-67. <https://doi.org/10.3382/ps.2007-00437>
- 232 5. Bedford MR, Apajalahti JH. The role of feed enzymes in maintaining poultry intestinal health. *J Sci Food Agric.*  
233 2022;102:1759-70. <https://doi.org/10.1002/jsfa.11670>
- 234 6. Chiang C-C, Yu B, Chiou PW-S. Effects of xylanase supplementation to wheat-based diet on the performance  
235 and nutrient availability of broiler chickens. *Asian-Australas J Anim Sci.* 2005;18:1141-6.  
236 <https://doi.org/10.5713/ajas.2005.1141>
- 237 7. Choi WJ, Kim JH, Kim HW, Kim KE, Kil DY. Effects of dietary palm kernel meal and  $\beta$ -xylanase on productive  
238 performance, fatty liver incidence, and excreta characteristics in laying hens. *J Anim Sci Technol.* 2021;63:1275.  
239 <https://doi.org/10.5187/jast.2021.e111>
- 240 8. Branton S, Lott B, Deaton J, Maslin W, Austin F, Pote L, et al. The effect of added complex carbohydrates or  
241 added dietary fiber on necrotic enteritis lesions in broiler chickens. *Poult Sci.* 1997;76:24-8.  
242 <https://doi.org/10.1093/ps/76.1.24>
- 243 9. Choct M. Managing gut health through nutrition. *Br Poult Sci.* 2009;50:9-15.  
244 <https://doi.org/10.1080/00071660802538632>
- 245 10. Adibmoradi M, Navidshad B, Faseleh Jahromi M. The effect of moderate levels of finely ground insoluble fibre  
246 on small intestine morphology, nutrient digestibility and performance of broiler chickens. *Ital J Anim Sci.*  
247 2016;15:310-7. <https://doi.org/10.1080/1828051X.2016.1147335>

- 248 11. Aviagen W. Ross 308 broiler nutrition specifications. Aviagen: Huntsville, AL, USA. 2014.
- 249 12. Wickramasuriya SS, Macelline SP, Cho HM, Hong JS, Patterson R, Heo JM. Multi-carbohydrase application  
250 into energy and amino acid deficient broiler diets: A strategy to enhance performance of broiler chickens. *J Anim*  
251 *Sci Technol.* 2021;63:1328. <https://doi.org/10.5187%2Fjast.2021.e104>
- 252 13. Wickramasuriya SS, Macelline SP, Cho HM, Hong JS, Park SH, Heo JM. Physiological effects of a tallow-  
253 incorporated diet supplemented with an emulsifier and microbial lipases on broiler chickens. *Front Vet Sci.*  
254 2020;7:583998. <https://doi.org/10.3389/fvets.2020.583998>
- 255 14. AOAC G. Official methods of analysis of AOAC International. Rockville, MD: AOAC International, ISBN: 978-  
256 0-935584-87-5. 2016
- 257 15. Oketch EO, Lee JW, Yu M, Hong JS, Kim YB, Nawarathne SR, et al. Physiological responses of broiler chickens  
258 fed reduced-energy diets supplemented with emulsifiers. *Anim Biosci.* 2022;35:1929-39.  
259 <https://doi.org/10.5713/ab.22.0142>
- 260 16. Kwak M-J, Ha D-J, Park MY, Eor JY, Whang K-Y, Kim Y. Comparison Study between Single Enzyme and  
261 Multienzyme Complex in Distiller's Dred Grains with Soluble Supplemented Diet in Broiler Chicken. *J Anim*  
262 *Sci Technol.* 2023. <https://doi.org/10.5187/jast.2023.e90>
- 263 17. Stevenson L, Phillips F, O'sullivan K, Walton J. Wheat bran: its composition and benefits to health, a European  
264 perspective. *Int J Food Sci Nutr.* 2012;63:1001-13. <https://doi.org/10.3109/09637486.2012.687366>
- 265 18. Vitaglione P, Napolitano A, Fogliano V. Cereal dietary fibre: a natural functional ingredient to deliver phenolic  
266 compounds into the gut. *Trends Food Sci Technol.* 2008;19(9):451-63. <https://doi.org/10.1016/j.tifs.2008.02.005>
- 267 19. Ko H, Kang HK, Moturi J, Ingale SL, Kim J. Supplementation of enzyme cocktail in chickens diet is an effective  
268 approach to increase the utilization of nutrient in wheat-based diets. *J Anim Sci Technol.* 2021;63:69.  
269 <https://doi.org/10.5187%2Fjast.2021.e1>
- 270 20. Van Hoeck V, Papadopoulos GA, Giannenas I, Lioliopoulou S, Tsiouris V, Mantzios T, et al. New Intrinsically  
271 Thermostable Xylanase Improves Broilers' Growth Performance, Organ Weights, and Affects Intestinal  
272 Viscosity and pH. *Agriculture.* 2021;11:1235. <https://doi.org/10.3390/agriculture11121235>
- 273 21. Lee JJ, Choi SH, Cho JH, Choe J, Kang J, Kim S, et al. Effects of dietary carbohydrases on productive  
274 performance and immune responses of lactating sows and their piglets. *J Anim Sci Technol.* 2019;61:359.  
275 <https://doi.org/10.5187/jast.2019.61.6.359>
- 276 22. Munezero O, Cho S, Kim IH. The effects of synbiotics-glyconutrients on growth performance, nutrient  
277 digestibility, gas emission, meat quality, and fatty acid profile of finishing pigs. *J Anim Sci Technol.* 2023.  
278 <https://doi.org/10.5187/jast.2023.e52>

- 279 23. Kim E, Morgan N, Moss A, Solbak A, Li L, Ader P, et al. In vitro degradation of non-starch polysaccharide  
280 residues in the digesta of broilers offered wheat-soy or maize-soy diets by feed enzymes. *J Appl Anim Nutr.*  
281 2022;10:71-81. <https://doi.org/10.3920/JAAN2022.0001>
- 282 24. Yacoubi N, Van Immerseel F, Ducatelle R, Rhayat L, Bonnin E, Saulnier L. Water-soluble fractions obtained by  
283 enzymatic treatment of wheat grains promote short chain fatty acids production by broiler cecal microbiota. *Anim*  
284 *Feed Sci Technol.* 2016;218:110-9. <https://doi.org/10.1016/j.anifeedsci.2016.05.016>
- 285 25. Choct M, Kocher A, Waters D, Pettersson D, Ross G. A comparison of three xylanases on the nutritive value of  
286 two wheats for broiler chickens. *Br J Nutr.* 2004;92:53-61. <https://doi.org/10.1079/BJN20041166>
- 287 26. Gao F, Jiang Y, Zhou G, Han Z. The effects of xylanase supplementation on performance, characteristics of the  
288 gastrointestinal tract, blood parameters and gut microflora in broilers fed on wheat-based diets. *Anim Feed Sci*  
289 *Technol.* 2008;142:173-84. <https://doi.org/10.1016/j.anifeedsci.2007.07.008>
- 290 27. Wealleans A, Ashour R, Abu Ishmais M, Al-Amairh S, Gonzalez-Sanchez D. Comparative effects of proteases  
291 on performance, carcass traits and gut structure of broilers fed diets reduced in protein and amino acids. *J Anim*  
292 *Sci Technol.* 2023. <http://dx.doi.org/10.5187/jast.2023.e20>.
- 293 28. Esmailipour O, Moravej H, Shivazad M, Rezaian M, Aminzadeh S, Van Krimpen M. Effects of diet  
294 acidification and xylanase supplementation on performance, nutrient digestibility, duodenal histology and gut  
295 microflora of broilers fed wheat based diet. *Br J Nutr.* 2012;53:235-44.  
296 <https://doi.org/10.1080/00071668.2012.681771>
- 297
- 298 29. Gorenz B, Iseri V, Rubach J, Dilger RN. Xylanase supplementation of pelleted wheat-based diets increases  
299 growth efficiency and apparent metabolizable energy and decreases viscosity of intestinal contents in broilers.  
300 *Poult Sci.* 2022;101:102220. <https://doi.org/10.1016/j.psj.2022.102220>
- 301 30. Barasch I, Grimes J. The effect of a heat-stable xylanase on digesta viscosity, apparent metabolizable energy and  
302 growth performance of broiler chicks fed a wheat-based diet. *Poult Sci.* 2021;100:101275.  
303 <https://doi.org/10.1016/j.psj.2021.101275>
- 304 31. Alam M, Howlider M, Pramanik M, Haque M. Effect of exogenous enzyme in diet on broiler performance. *Int J*  
305 *Poult Sci.* 2003;2:168-73. <https://doi.org/10.3923/ijps.2003.168.173>
- 306 32. Williams M, Klein J, Wyatt C, York T, Lee J. Evaluation of xylanase in low-energy broiler diets.  
307 *J Appl Poult Res.* 2014;23:188-95. <http://dx.doi.org/10.3382/japr.2013-00856>

- 308 33. Hussein E, Suliman G, Alowaimer A, Ahmed S, Abd El-Hack M, Taha A, et al. Growth, carcass characteristics,  
309 and meat quality of broilers fed a low-energy diet supplemented with a multienzyme preparation. *Poult Sci.*  
310 2020;99:1988-94. <https://doi.org/10.1016/j.psj.2019.09.007>
- 311 34. Fouad A, El-Senousey H. Nutritional factors affecting abdominal fat deposition in poultry: a review. *Asian-*  
312 *Australas J Anim Sci.* 2014;27:1057. <https://doi.org/10.5713%2Fajas.2013.13702>
- 313 35. Park T-W, Lee E-Y, Jung Y, Son YM, Oh S-H, Kim D-H, et al. Effects of lysine concentration of the diet on  
314 growth performance and meat quality in finishing pigs with high slaughter weights. *J Anim Sci Technol.* 2023.  
315 <http://dx.doi.org/10.5187/jast.2023.e49>

ACCEPTED



Tables

**Table 1.** Composition (g/kg, as-fed basis) of the experimental diets<sup>1</sup>

Items	Starter phase (d 8-24)					Grower phase (d 25-35)				
	CON	CON-X	L-X	M-X	H-X	CON	CON-X	L-X	M-X	H-X
<b>Ingredient (%)</b>										
Corn	61.45	61.45	59.46	57.48	53.50	63.79	63.79	58.01	52.22	40.65
Wheat bran	-	-	2.00	4.00	8.00	-	-	2.00	4.00	8.00
Soybean meal (SBM, 48 %)	31.62	31.62	30.94	30.25	28.88	27.44	27.44	27.62	27.81	28.17
Fish meal	1.30	1.30	1.60	1.90	2.50	1.90	1.90	2.05	2.20	2.50
Vegetable oil	1.20	1.20	1.78	2.35	3.50	2.45	2.45	2.84	3.23	4.00
Corn starch	-	-	-	-	-	-	-	3.25	6.50	13.00
Limestone	1.78	1.78	1.59	1.39	1.00	1.84	1.84	1.67	1.50	1.15
Mono-calcium phosphate	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50	1.50
Salt	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30
Vitamin-mineral premix <sup>2</sup>	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30
Lys-HCl	0.25	0.25	0.25	0.25	0.25	0.20	0.20	0.19	0.18	0.15
DL-methionine	0.30	0.30	0.29	0.28	0.27	0.28	0.28	0.28	0.28	0.28
<b>Calculated composition</b>										
Metabolizable energy (kcal/kg)	3,050	3,050	3,050	3,050	3,050	3,150	3,150	3,150	3,150	3,150
Crude protein (%)	21.50	21.50	21.50	21.50	21.50	20.00	20.00	20.00	20.00	20.00

Non-starch polysaccharides (%)	8.33	8.33	8.70	9.07	9.82	7.89	7.89	8.15	8.40	8.91
Lysine (%)	1.32	1.32	1.32	1.32	1.33	1.20	1.20	1.20	1.20	1.20
Methionine +Cystine (%)	0.99	0.99	0.99	0.99	0.99	0.94	0.94	0.94	0.94	0.94

**Analyzed composition**

Analyzed xylanase unit (U/g)	-	3.01	3.00	3.00	3.01	-	3.00	2.99	3.00	3.00
------------------------------	---	------	------	------	------	---	------	------	------	------

<sup>1</sup>Treatments; i) CON (Control diet), ii) CON-X (CON + 3000 U/kg xylanase), iii) L-X (corn replaced to 2% wheat bran in CON + 3000 U/kg xylanase), iv) M-X: (corn replaced to 4% wheat bran in CON + 3000 U/kg xylanase), v) H-X (corn replaced to 8% wheat bran in CON+ 3000 U/kg xylanase).

<sup>2</sup>Vitamin and mineral mixture provided the following nutrients per kg of diet: vitamin A, 24,000 IU; vitamin D3, 6,000 IU; vitamin E, 30 IU; vitamin K, 4 mg; thiamin, 4 mg; riboflavin, 12 mg; pyridoxine, 4 mg; folacine, 2 mg; biotin, 0.03 mg; vitamin B8 0.06 mg; niacin, 90 mg; pantothenic acid, 30 mg; Fe, 80 mg (as FeSO<sub>4</sub> · H<sub>2</sub>O); Zn, 80 mg (as ZnSO<sub>4</sub> · H<sub>2</sub>O); Mn, 80 mg (as MnSO<sub>4</sub> · H<sub>2</sub>O); Co, 0.5 mg (as CoSO<sub>4</sub> · H<sub>2</sub>O); Cu, 10 mg (as CuSO<sub>4</sub> · H<sub>2</sub>O); Se, 0.2 mg (as Na<sub>2</sub>SeO<sub>3</sub>); I, 0.9 mg (as Ca (IO<sub>3</sub>) · 2H<sub>2</sub>O).

**Table 2:** Effects of different dietary non-starch polysaccharide (NSP) level diets on the growth performance of broilers<sup>1</sup>

Items	Treatments <sup>2</sup>					SEM <sup>3</sup>	P value
	CON	CON-X	L-X	M-X	H-X		
<b>Body weight (g)</b>							
Day 8	114.8	115.1	115.1	114.7	115.4	1.555	0.992
Day 14	268.8	276.6	275.7	283.0	281.1	9.655	0.626
Day 21	597.2	598.0	597.1	587.7	615.3	23.476	0.832
Day 24	769.4	776.1	799.6	779.4	814.8	30.246	0.550
Day 28	1052.1	1064.7	1120.2	1077.6	1131.9	44.355	0.310
Day 35	1630.6	1650.6	1708.9	1669.5	1806.7	64.405	0.074
<b>Average daily gain (g/day)</b>							
Day 14	22.00	23.06	22.94	24.04	23.68	1.376	0.633
Day 21	46.91	45.93	45.91	43.53	47.75	2.803	0.641
Day 28	70.67	72.16	80.14	74.54	79.27	0.562	0.360
Day 35	82.64 <sup>b</sup>	83.70 <sup>ab</sup>	84.11 <sup>ab</sup>	84.55 <sup>ab</sup>	96.40 <sup>a</sup>	4.537	0.026
Day 8-24	38.51	38.88	40.26	39.10	41.15	1.790	0.567
Day 25-35	78.29	79.50	82.67	80.91	90.17	4.117	0.053
Day 8-35	54.14	54.84	56.92	55.53	60.41	2.287	0.073
<b>Average daily feed intake (g/day)</b>							
Day 14	31.36	32.47	32.91	32.65	31.97	1.194	0.715
Day 21	67.52	64.98	66.73	64.21	61.56	2.031	0.049
Day 28	105.05	101.42	115.88	105.55	112.48	8.208	0.400
Day 35	116.97 <sup>ab</sup>	108.51 <sup>b</sup>	124.58 <sup>ab</sup>	113.99 <sup>ab</sup>	125.96 <sup>a</sup>	6.060	0.041
Day 8-24	56.61	55.99	57.57	55.91	54.73	1.530	0.316
Day 25-35	112.74	106.53	115.74	110.85	121.73	5.817	0.141
Day 8-35	78.50	75.68	80.26	77.74	80.95	2.992	0.426
<b>Feed conversion ratio (g/g)</b>							
Day 14	1.44	1.42	1.44	1.37	1.36	0.070	0.621
Day 21	1.46	1.44	1.47	1.50	1.30	0.107	0.404
Day 28	1.49	1.41	1.46	1.42	1.44	0.104	0.950
Day 35	1.42	1.30	1.49	1.35	1.31	0.072	0.334

Day 8-24	1.47	1.44	1.43	1.43	1.33	0.066	0.269
Day 25-35	1.44	1.34	1.40	1.37	1.35	0.058	0.388

<sup>1</sup>Values are the mean of eight replicates per treatment

<sup>2</sup>Treatments; i) CON(Control diet), ii) CON-X (CON + 3000 U/kg xylanase), iii) L-X (corn replaced to 2% wheat bran in CON + 3000 U/kg xylanase), iv) M-X: (corn replaced to 4% wheat bran in CON + 3000 U/kg xylanase), v) H-X (corn replaced to 8% wheat bran in CON+ 3000 U/kg xylanase).

<sup>3</sup>Pooled standard error of the mean, <sup>a-b</sup>Values in a row with different superscripts differ significantly ( $p < 0.05$ )

ACCEPTED

**Table 3:** Effects of different dietary non-starch polysaccharide (NSP) level diets on nutrient digestibility and viscosity of broilers on day 24 and 35<sup>1</sup>

Items	Treatments <sup>2</sup>					SEM <sup>3</sup>	P-value
	CON	CON-X	L-X	M-X	H-X		
<b>Energy digestibility (%)</b>							
d 24	86.86 <sup>a</sup>	83.80 <sup>ab</sup>	86.27 <sup>a</sup>	86.62 <sup>a</sup>	80.21 <sup>b</sup>	2.011	<0.001
d 35	87.72 <sup>a</sup>	88.15 <sup>a</sup>	86.45 <sup>ab</sup>	83.70 <sup>bc</sup>	82.75 <sup>c</sup>	1.013	0.002
<b>Protein digestibility (%)</b>							
d 24	73.37 <sup>b</sup>	75.25 <sup>a</sup>	72.47 <sup>a</sup>	72.22 <sup>b</sup>	72.95 <sup>b</sup>	1.605	<0.001
d 35	75.58	76.82	75.62	75.43	75.71	0.710	0.596
<b>Viscosity, mPa/s</b>							
d 24	2.63 <sup>a</sup>	2.50 <sup>ab</sup>	2.45 <sup>abc</sup>	2.37 <sup>bc</sup>	2.28 <sup>c</sup>	0.467	<0.001
d 35	2.58 <sup>a</sup>	2.56 <sup>a</sup>	2.41 <sup>b</sup>	2.30 <sup>c</sup>	2.18 <sup>d</sup>	0.326	<0.001

<sup>1</sup>Values are the mean of eight replicates per treatment

<sup>2</sup>Treatments; i) CON(Control diet), ii) CON-X (CON + 3000 U/kg xylanase), iii) L-X (corn replaced to 2% wheat bran in CON + 3000 U/kg xylanase), iv) M-X: (corn replaced to 4% wheat bran in CON + 3000 U/kg xylanase), v) H-X (corn replaced to 8% wheat bran in CON+ 3000 U/kg xylanase).

<sup>3</sup>Pooled standard error of the mean

<sup>a,b-d</sup> Values in a row with different superscripts differ significantly ( $p < 0.05$ )

**Table 4:** Effects of different dietary non-starch polysaccharide (NSP) level diets on carcass composition of broilers on day 35<sup>1</sup>

Items	Treatments <sup>2</sup>					SEM <sup>3</sup>	P value
	CON	CON-X	L-X	M-X	H-X		
<b>Leg meat (%)</b>							
Moisture	70.51 <sup>e</sup>	71.42 <sup>c</sup>	70.78 <sup>d</sup>	73.15 <sup>a</sup>	72.42 <sup>b</sup>	1.072	<0.001
Crude protein	19.14	19.48	20.15	19.55	18.99	0.638	0.463
Crude fat	8.19 <sup>a</sup>	7.79 <sup>b</sup>	7.67 <sup>b</sup>	7.72 <sup>b</sup>	6.16 <sup>c</sup>	1.087	<0.001
Ash	1.13	1.17	1.24	1.20	1.21	0.712	0.587

<sup>1</sup>Values are the mean of eight replicates per treatment

<sup>2</sup>Treatments; i) CON(Control diet), ii) CON-X (CON + 3000 U/kg xylanase), iii) L-X (corn replaced to 2% wheat bran in CON + 3000 U/kg xylanase), iv) M-X: (corn replaced to 4% wheat bran in CON + 3000 U/kg xylanase), v) H-X (corn replaced to 8% wheat bran in CON+ 3000 U/kg xylanase).

<sup>3</sup>Pooled standard error of the mean

<sup>a,b-d</sup> Values in a row with different superscripts differ significantly ( $p < 0.05$ )