

Review article

Interaction between host cell proteins and open reading frames of porcine circovirus type 2

Running title: Molecular pathogenesis of porcine circovirus type 2

Si-Won Park[#], In-Byung Park[#], Seok-Jin Kang, Joonbeom Bae and Taehoon Chun*

Department of Biotechnology, School of Life Sciences and Biotechnology, Korea University, Seoul 02841, Republic of Korea

[#]These authors contributed equally to this work.

***Corresponding author:** Taehoon Chun, Department of Biotechnology, College of Life Sciences and Biotechnology, Korea University, Seoul, 02841, Korea. Tel: +82-2-3290-3069
E-mail: tchun@korea.ac.kr

E-mail addresses of authors

Si-Won Park: kkd01119@korea.ac.kr

In-Byung Park: tnquddls@korea.ac.kr

Seok-Jin Kang: laughingkang@korea.ac.kr

Joonbeom Bae: joonbeombae@korea.ac.kr

Taehoon Chun: tchun@korea.ac.kr

Abbreviations: AMPK, AMP-activated protein kinase; ANT3, adenine nucleotide translocase 3; ATF6, activating transcription factor 6; COX8A, cytochrome c oxidase subunit 8A; CYP1A1, cytochrome P450 1A1; C1QBP, complement component 1, q subcomponent binding protein; DDX21, DExD-Box Helicase 21; EIF4A2, eukaryotic translation initiation factor 4A2; ER, endoplasmic reticulum; FHC, ferritin heavy chain; GPNMB, glycoprotein-NMB; HIV, human immunodeficiency virus; hnRNPC, heterogeneous nuclear ribonucleoproteins C1/C2; Hsp40, heat shock protein 40; IC1, plasma protease C1 inhibitor; IFN, interferon; IPO5, Importin-5; IRE1, inositol requiring enzyme 1; MKRN1, makorin-1 RING zinc finger protein; mTOR, mammalian target of rapamycin; MX2, MX Dynamin Like

GTPase 2; NAP1, nucleosome assembly protein-1; NEF, nucleotide-exchange factor; NPM1, nucleophosmin-1; ORF, open reading frame; Par-4, prostate apoptosis response-4; PCV2, porcine circovirus type 2; PCVAD, PCV2 infection causes porcine circovirus-associated diseases; PCV2-SD, PCV2-systemic disease; PDNS, porcine dermatitis and nephropathy syndrome; PERK, RNA-activated protein kinase-like ER kinase; PMWS, postweaning multisystemic wasting syndrome; RGS16, regulator of G protein signaling 16; SNAI2, snail family transcriptional repressor 2; SNRPN, lamin C and small nuclear ribonucleoprotein polypeptide N; SRSF3, serine and arginine rich splicing Factor 3; TDG, thymine DNA glycosylase; VG5Q, angiogenic factor with G patch and FHA domains 1; YWHAB, tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein b; ZNF, zinc finger protein.

Total words in abstract: 229

Total words in text: 4304

Abstract

Postweaning multisystemic wasting syndrome (PMWS) is caused by a systemic inflammation after porcine circovirus type 2 (PCV2) infection. It was one of the most economically important pathogens affecting pig production worldwide before PCV2 vaccine was first introduced in 2006. After the development of a vaccine against PCV2a type, pig farms gradually restored enormous economic losses from PMWS. However, vaccine against PCV2a type could not be fully effective against several different PCV2 genotypes (PCV2b – PCV2h). In addition, PCV2a vaccine itself could generate antigenic drift of PCV2 capsid. Therefore, PCV2 infection still threatens pig industry worldwide. PCV2 infection was initially found in local tissues including reproductive, respiratory, and digestive tracks. However, PCV2 infection often leads to a systemic inflammation which can cause severe immunosuppression by depleting peripheral lymphocytes in secondary lymphoid tissues. Subsequently, a secondary infection with other microorganisms can cause PMWS. Eleven putative open reading frames (ORFs) have been predicted to encode PCV2 genome. Among them, gene products of six ORFs from ORF1 to ORF6 have been identified and characterized to estimate its functional role during PCV2 infection. Acquiring knowledge about the specific interaction between each PCV2 ORF protein and host protein might be a key to develop preventive or therapeutic tools to control PCV2 infection. In this article, we reviewed current understanding of how each ORF of PCV2 manipulates host cell signaling related to immune suppression caused by PCV2.

Keywords: host cell protein, open reading frame, pathogenesis, pig, porcine circovirus type 2

INTRODUCTION

Postweaning multisystemic wasting syndrome (PMWS) is now referred to as a porcine circovirus type 2 (PCV2) systemic disease (PCV2-SD) because PCV2 infection is evidently associated with PMWS [1, 2]. PMWS was first found in the middle of 1990's. It was one of the most disastrous diseases in pig industry worldwide before PCV2 vaccine was first commercialized in 2006 [1-3]. After the introduction of a vaccine against PCV2a type, pig producers gradually recovered from economic losses caused by PMWS [4, 5]. However, several evidence indicates that vaccination against PCV2a type cannot completely remove circulating PCV2 after an infection [6, 7]. Besides, PMWS still occurs occasionally in vaccinated piglets [8, 9]. Several speculations have been raised to explain why PCV2a vaccine is not effective to eradicate PCV2 infection. First, PCV2a vaccine might not cover all genetic variations of PCV2. At least eight different PCV2 genotypes (PCV2a – PCV2h) have been identified based on their capsid structures [10]. Indeed, the field studies in US and Korea have reported the failure of PCV2a vaccine during outbreaks of PCV2b mutant infections [8, 9]. Second, vaccination against PCV2 itself can be a driven force to induce antigenic variation of PCV2 capsid [11, 12]. Therefore, developing multivalent vaccines against PCV2 is currently ongoing [13]. Overall, PCV2 infection is still considered as one of the most critical problems in pig industry worldwide [14].

Clinical symptoms after PCV2 infection can vary from regional inflammation targeting reproductive, respiratory, or digestive tracks to systemic inflammation including PMWS [15]. Although the mechanism is not clear, systemic infection of PCV2 in piglets often causes an immune suppression by depleting peripheral lymphocytes [16]. Therefore, PMWS caused by systemic PCV2 infection can gradually decrease the immunity of piglets and cause a secondary infection by bacteria or other viruses [17]. Defining which open reading frame (ORF) of PCV2 modifies a host cell physiology by interacting with host cell proteins is very helpful for understanding the molecular pathology of PCV2. This is particularly significant when we exam the pathology of PMWS related to immune suppression, which is a well-known strategy used by PCV2 to escape from host immune surveillance [18]. In this review, we will summarize current knowledge of interactions between each ORF of PCV2 and host proteins and introduce several theories on how PCV2 causes immune suppression after infection.

GENOTYPES AND ASSOCIATED DISEASES OF PCV

PCV belongs to the genus *Circovirus* in the family *Circoviridae* which has a non-enveloped icosahedral capsid [17]. The genome of PCV is composed of a single-stranded DNA with a closed circular shape (Fig. 1). It contains very small amounts of genetic information with a length of 1.7 to 2.0 kb [19]. Four PCV genotypes have been identified so far [20]. Recent phylogenetic analysis based on amino acid sequences of replicase (ORF1) of each PCV genotype has indicated that PCV1, PCV2, and PCV3 might be derived from bat circovirus because PCV1 and PCV2 share a common ancestor with bat-clade 2 circoviruses and replicase of PCV3 is most similar to bat-clade 1 circoviruses [21, 22]. However, PCV4 shares a common ancestor with mink circoviruses having a distinct origin from PCV1 - PCV3 [22]. Therefore, pig circovirus might have emerged from cross-species transmission of other mammalian species [21].

PCV1 was originally identified in the genome of immortalized porcine kidney cells (PK-15 cells) [19]. PCV1 infection does not cause any types of swine diseases [19]. In contrast to PCV1, PCV2 infection causes porcine circovirus-associated diseases (PCVAD) including porcine dermatitis and nephropathy syndrome (PDNS), PMWS, and porcine reproductive diseases [23-25]. PCV3 genome was first identified in 2015 [26]. PCV3 infection is also associated with various swine diseases including gastrointestinal, neurological, reproductive, and respiratory disorders [26-28]. PCV4 was first identified by serotyping as early as 2008, but the genome of PCV4 was fully characterized in 2019 [29]. Recent observations have indicated that PCV4 infection is also correlated with clinical symptoms of pigs such as gastrointestinal and respiratory distresses, and PDNS [22, 30, 31]. Therefore, all identified types of PCV except PCV1 are considered as major pathogens to threat pig industry worldwide.

PROPOSED FUNCTION OF EACH PCV2 ORF

Eleven ORFs have been predicted to encode PCV2 genome [32]. Among them, ORF 1, 5, 7, and 10 are likely to be located on the sense strand of PCV genome and the rest of ORFs are encoded by the anti-sense strand of PCV genome (Fig. 1) [32]. Until now, only six ORFs from ORF1 to ORF6 have been confirmed to produce polypeptides with their putative functions characterized [33, 34]. The ORF1 encodes enzymes essential for the replication of PCV2 [35]. Several mRNA splicing variants have been detected from ORF1. These splicing variants are also involved in the replication of PCV2 [35]. The ORF2 encodes capsid, a major structural protein and the major source of PCV2 vaccine. It serves as a target of neutralizing antibodies to block viral entry to host cells [36]. The rest of identified ORFs (ORF3 – ORF6)

are not essential for the replication of PCV2. They mainly modulate host cell physiology by interacting with host cell proteins [34, 37-39].

The pathogenesis of PCV2 infection is not fully understood yet. Although the infection route of PCV2 is currently unclear, detectable amounts of PCV2 have been found in secretions and excretions of infected pigs [40-43]. Therefore, oral, nasal, and fecal contaminations might mediate horizontal transmission [44-46]. Vertical transmission could also occur through maternal-fetal interface [47, 48]. One of the symptoms of PCVAD is abortion or mummification of fetus because infected fetus cannot survive due to PCV2 replication in heart, liver, lung, or lymphoid tissues of fetus [48-51].

PCV2 mainly infects hematopoietic lineage cells such as lymphocytes and macrophages [52, 53]. Especially, massively proliferating lymphoblasts are known to be major host cells of PCV2 [53]. Since these major host cells of PCV2 are immune cells, PCV2 particles are preferentially found in lymphoid tissues [24]. However, several reports have provided evidence that PCV2 replication occurs in non-lymphoid tissues such as lungs, livers, and kidneys of piglets [54, 55]. This may indicate that macrophages in non-lymphoid tissues can be infected when PCV2 infection is systemically disseminated in the pig body, or PCV2 itself can infect other cell types besides immune cells. Other cell types reported to be host cells of PCV2 include endothelial cells, epithelial cells, fetal cardiomyocytes, and primary hepatocytes [56-61]. However, these results are mainly acquired from *in vitro* infection studies using several cell lines or primary cells [56-61].

To study the life cycle of PCV2 or the interaction between PCV2 ORFs and host proteins, two types of pig cell lines, 3D4/31 and PK-15, have been extensively used. 3D4/31 cells are derived from pig alveolar macrophages and PK-15 cells are pig kidney epithelial cells [62, 63]. Genetically engineered artificial PCV2 or natural PCV2 originally isolated from pigs with PMWS can infect and replicate within both cell lines [64, 65].

Previous study using 3D4/31 or PK-15 cells has shown that the capsid of PCV2 can recognize heparan sulphate or chondroitin sulphate B glycosaminoglycans on host cells [66]. PCV2 can attach to these receptors expressed on porcine macrophages within 30 minutes [65]. A motif containing “⁹⁸IRKVKV¹⁰³” residues within PCV2 capsid has been identified as a putative heparin binding motif based on structural analysis using crystallography [67]. After attachment, PCV2 particle is internalized via clathrin-dependent endocytosis. Nucleic acid of PCV2 is then released by endosomal acidification [65]. These processes are further confirmed by treatment with a serine protease inhibitor which can abolish penetration of PCV2 genome into cytoplasm of both 3D4/31 and PK-15 cells [68]. After penetration, single-stranded DNA of PCV2 is transported into nucleus, converted to

double-stranded intermediates, and replicated via a rolling-circle replication method [69].

The hallmark of PMWS mediated by systemic PCV2 infection is immunosuppression in pigs [14, 24]. Major leukocytes including helper T cells, B cells and dendritic cells are significantly decreased in pigs with PMWS [70-72]. These observations have been further supported by the fact that PCV2 titers in periphery of pigs are correlated with the severity of immune cell destruction and antibody deficiency against PCV2 [72]. Immunosuppression due to depletion of immune effector cells can cause secondary infections in pigs with PMWS [16, 73-75]. Therefore, pigs with PMWS, similar to immunocompromised people, are not directly killed by PCV2 infection, but sacrificed by secondary infections with other microbial pathogens targeting lung or digestive tracks [76, 77].

HOST CELL MODULATION BY PCV2 ORFs

Virus infection can significantly change physiological conditions of infected animals. Since host inflammatory responses are dependent on the titer and antigenicity of viral antigens and/or the ability of infiltrated immune cells in certain infectious sites, the clue explaining how virus can modulate host physiology might be found in specific interactions between viral proteins and host proteins. There are many examples indicating the virus itself has evolved to have several strategies to downregulate or escape immune surveillance of its host. Otherwise, viruses cannot survive inside their hosts. Several evidence has shown that PCV2 can also use this tactic inside its host. Each ORF's function related to modulation of host cell physiology or immune surveillance by having a crosstalk with host proteins will be discussed below.

PCV2 ORF1

Since PCV2 ORF1 encodes replicase of PCV2, its main role is focus on generation of progeny virus rather than interaction with host cell proteins [35]. Only two studies have identified PCV2 ORF1 interacting proteins from host cells by screening bacterial or yeast two hybrid libraries using PCV2 ORF1 as a bait [78, 79]. Five PCV2 ORF1 binding partners have been identified in host cells (Table 1). Two of them, syncoilin and c-myc, have been identified in bacterial two hybrid libraries and the rest of them, thymine DNA glycosylase (TDG), angiogenic factor with G patch and FHA domains 1 (VG5Q), and zinc finger protein 265 (ZNF265), have been identified in yeast two hybrid libraries [78, 79]. Especially, TDG, VG5Q, and ZNF265 can bind to both PCV1 and PCV2 replicase [79].

Syncoilin is an intermediate filament protein found in various cell types including skeletal and cardiac muscle, hepatic stellate cells, and neurons [80-82]. It is valuable to test whether blocking syncoilin's function can inhibit PCV2 replication because vimentin, another intermediate filament protein, is the target for a novel drug to block viral protein transport to specific organelle in host cells [83-85]. As a well-known transcriptional factor and an oncogene, c-myc can facilitate cell cycle progress [86]. It can also initiate DNA replication non-transcriptionally by binding with members of the prereplication protein complex [87, 88]. Therefore, PCV2 replicase could bind with c-myc for the initiation of DNA replication in host cells.

TDG is involved in G/T mismatch repairs by removing thymine residues from sugar-phosphate DNA backbone [89]. VG5Q is highly expressed in activated endothelial cells. It can promote angiogenesis by interacting with proteins involved in vascular development [90]. Overexpression of VG5Q can lead to poor prognosis by increasing cancer metastasis [91, 92]. ZNF265 is a zinc finger protein that contains an arginine/serine-rich domain and a subunit of pre-mRNA spliceosome [93]. Therefore, ZNF265 is involved in mRNA splicing [93]. Since TDG and ZNF265 participate in DNA replication and mRNA splicing, PCV2 replicase may interact with these proteins during replication or splicing of viral DNA or mRNA. However, the reason why PCV2 replicase binds with VG5Q, an angiogenetic factor, is still elusive.

PCV2 ORF2

Initial binding partners of PCV2 capsid, an ORF2 encoding protein, have been screened using bacterial two hybrid libraries and PK-15 cDNA [78]. P-selectin and complement component 1, q subcomponent binding protein (C1QBP or gC1qR) have been found to be able to bind to capsid (Table 1) [78]. P-selectin is one of the selectins that belong to C-type lectin, a carbohydrate binding protein [94]. P-selectin is mainly expressed on activated endothelial cells and platelets [95]. Although additional example showing that viral capsid can bind to P-selectin has not been documented yet, L-selectin, its closest paralog, is critical for human immunodeficiency virus (HIV)-entry into host cells by interacting with gp120, an HIV spike protein [96]. Therefore, PCV2 capsid might interact with P-selectin during attachment to host cells.

C1QBP is a receptor of globular heads of C1q which is expressed on a broad range of cell types [97]. C1QBP is mainly expressed in the mitochondrial matrix. It is also detected in nucleus, endoplasmic reticulum (ER), and cell surface [98]. Therefore, C1QBP could play multifunctional roles in various cellular events based on its multi-subcellular distributions

[98]. Indeed, C1QBP can mediate various inflammatory responses including blood clotting, recruitment of leukocytes, phagocytosis, and production of reactive oxygen intermediates [99-102]. Another study has also identified C1QBP as a binding protein of PCV2 capsid protein through screening using a yeast two-hybrid system [79]. Further study has revealed that PCV2 capsid can facilitate phagocytic activity and secretions of proinflammatory cytokine in porcine macrophages by inhibiting ubiquitin-mediated proteasomal degradation of C1QBP via direct interactions [103]. This finding is further supported by additional reports demonstrating that the interaction between C1QBP and PCV2 capsid can induce IL-10 production by activating PI3K/Akt and p38 MAPK pathways in porcine alveolar macrophages [104]. Additional host proteins identified as binding partners of PCV2 capsid by screening a yeast two-hybrid library include heat shock protein 40 (Hsp40), makorin-1 RING zinc finger protein (MKRN1), nucleosome assembly protein-1 (NAP1), nucleophosmin-1 (NPM1), and prostate apoptosis response-4 (Par-4) (Table 1) [79]. Hsp40 is a molecular chaperone that can assist the function of Hsp70 with nucleotide-exchange factor (NEF) [105, 106]. Hsp40 can bind to misfolded proteins and deliver these proteins to Hsp70, and NEF can catalyze the dissociation of ADP from Hsp70 to allow ATP binding to Hsp70 [106]. Therefore, Hsp40 and NEF are essential for the function of Hsp70, the common molecular chaperone for protein folding in mammalian cells [106]. In agreement with this observation, an independent study has demonstrated that PCV2 capsid can interact with Hsp70 to enhance progeny virus production of PCV2 by increasing PCV2 genome replication in 3D4/31 cells [107].

Among host proteins that can bind to PCV2 capsid, MKRN1, NAP1, and NPM1 are nuclear proteins or proteins with nucleotide binding motifs [108-114]. MKRN1 is an RNA-binding E3 ubiquitin ligase with an RNA binding property and conserved zinc finger domains [108]. MKRN1 also has an E3 ubiquitin ligase activity that can regulate cell cycle or cell senescence by regulating target protein turnover rates [108-110]. NAP1 is an evolutionarily conserved protein. It might be involved in a nucleosome assembly because it has been found in several different chromatin remodeling complexes [111, 112]. NPM1 is often found in nucleolar ribonucleoprotein structures. It serves as an abundant nucleus-cytoplasmic shuttling protein to transport nucleic acid [113, 114].

Par-4 serves as a caspase-3 substrate that can enhance apoptosis under various apoptotic signals [115]. Since PCV2 infection can lead to lymphocyte depletion by inducing host cell apoptosis, how the interaction between PCV2 capsid and Par-4 regulates apoptosis in host cells might be a necessary research subject [70-72].

A recent study using co-immunoprecipitation with anti-PCV2 capsid antibody in PK-15 cells has revealed that 222 additional host proteins might interact with PCV2 capsid [116].

Host proteins that interact with PCV2 capsid might participate in a broad range of cell physiology including nucleic acid homeostasis, transcriptional regulation, several metabolic pathways, and innate immunity [116]. Among them, host proteins such as DExD-Box Helicase 21 (DDX21), eukaryotic translation initiation factor 4A2 (EIF4A2), heterogeneous nuclear ribonucleoproteins C1/C2 (hnRNP), plasma protease C1 inhibitor (IC1), Importin-5 (IPO5), MX Dynamin Like GTPase 2 (MX2), nucleophosmin 1 (NPM1), and snail family transcriptional repressor 2 (SNAI2) have been confirmed to be able to bind with PCV2 capsid in PK-15 by pull-down assays (Table 1) [116].

Several PCV2 capsid binding proteins including DDX21, EIF4A2, hnRNP, IPO5, NPM1, and SNAI2 are often found in the nucleus [117-122]. DDX21 is a putative RNA helicase that can modify secondary RNA structure during RNA splicing or translational initiation [117]. EIF4A2 is a component of eIF4F translation initiation complex that exhibits RNA helicase activity [118]. The hnRNP has a putative RNA-binding domain. It might participate in RNA homeostasis such as processing heterogeneous nuclear RNAs and stabilizing mRNAs [119]. IPO5 is a member of the importin beta family known to transport materials from cytosol into nucleus [120]. NPM1 is a nuclear protein. It is also identified as a PCV2 capsid binding protein in a previous study using a yeast two-hybrid library [79]. SNAI2 is a transcription factor that possesses a zinc finger domain. It can promote differentiation and migration of mammalian cells [121].

IC1 and MX2 are two proteins that can interact with PCV2 capsid. They are involved in innate immunity [122, 123]. IC1 belongs to the serpin family of protease inhibitors that can suppress local inflammation by decreasing vascular permeability [122]. MX2 is one of the dynamin superfamilies of large GTPases. It is an interferon (IFN) induced GTPase that might prevent the assembly of progeny virus in host cells by blocking viral capsid transport [123].

PCV2 ORF3

The ORF3 gene is about 315 bp in length. It is located within the ORF1 gene and transcribed in the opposite direction of ORF1 mRNA transcript (Fig. 1) [124]. The ORF3 encoding protein was firstly identified in PCV2 infected PK-15 cells [124]. A further genomic study has revealed that non-virulent PCV1 has a truncated ORF3 genome [125]. This may imply that ORF3 possibly encodes a virulence factor of PCV2. Indeed, three independent studies have demonstrated that PCV2 ORF3 is the major protein which induces apoptosis of host cells after PCV2 infection [37, 124, 126]. The proposed mechanism of how PCV2 ORF3 induces apoptosis is that ORF3 competes with p53, tumor suppressor, for the

binding with Pirh2, an E3 ubiquitin ligase (Fig. 2A) [127, 128]. Therefore, PCV2 ORF3 can increase half-life of p53 by blocking ubiquitin-dependent proteasomal degradation of p53 (Fig. 2A) [127, 128].

Since ORF3 protein can induce apoptosis of PCV2 infected host cells *in vitro*, several studies have attempted to estimate the pathological significance of ORF3 during a real PCV2 infection [129-131]. Initial two studies using PCV2 infected animals have revealed that ORF3-null mutant of PCV2 is less virulent in mice and piglets [129, 130]. However, an independent study of PCV2 infection model later reported that clinical symptoms in piglets were not significantly changed when they were infected by ORF3-null mutant of PCV2 compared to those infected by wild-type PCV2, although ORF3-null PCV2 mutant infected piglets exhibited a delayed antibody production and lower serum levels of viral load [131].

Besides Pirh2, Tn10 transposase (DDE-like transposase) and regulator of G protein signaling 16 (RGS16) have also been identified as host binding proteins of PCV2 ORF3 by screening a bacterial two-hybrid library using cDNA of PK-15 cells (Table 1) [78]. Although the biological meaning of the interaction between Tn10 transposase and PCV2 ORF3 has not been documented yet, two independent studies have tried to find out changes of host cell physiology after interaction between RGS16 and PCV2 ORF3 [132, 133]. The first study regarding interaction between RGS16 and PCV2 ORF3 confirmed the interaction between these two proteins by *in vitro* pull-down assay and showed that these two proteins were co-localized within nucleus [132]. The second study demonstrated that PCV2 ORF3 could facilitate proteasomal degradation of RGS16 by direct interaction [133]. Since RGS16, an inhibitor of large G protein signaling, suppresses the production of proinflammatory cytokines via attenuation of NF κ B signaling, the degradation of RGS16 can lead to enhancement of NF κ B translocation into the nucleus via the ERK1/2 signaling pathway and transcription of IL-6 and IL-8 in porcine epithelial cells, PK-15 cells (Fig. 2B) [133-135].

PCV2 ORF4

The ORF4 gene is about 180 bp in length. It is located within the ORF3 gene and transcribed in the same direction as ORF3 mRNA transcript (Fig. 1) [136]. Initial studies using PCV2 ORF4 deletion mutant have revealed that PCV2 ORF4 might prevent apoptosis of host cells [136, 137]. A further study using a yeast two-hybrid system has identified that PCV2 ORF4 can bind with ferritin heavy chain (FHC) (Table 1) [138]. FHC has a ferroxidase activity. It serves as an anti-apoptotic factor via suppressing the accumulation of reactive oxygen intermediates [139]. Therefore, PCV2 ORF4 might play a role in rescuing host cells

from apoptosis by coordinating with FHC which removes accumulative reactive oxygen stress [139]. However, a recent study has argued that PCV2 ORF4 rather induces host cell apoptosis via interaction with mitochondrial adenine nucleotide translocase 3 (ANT3) [140]. In brief, PCV2 ORF4 possesses amphipathic α -helix targeting mitochondria at *N*-terminus and ANT3 can help ORF4 protein anchor to mitochondria [140]. After migration to mitochondria with an assistance by ANT3, PCV2 ORF4 could directly induce mitochondria-mediated apoptosis of host cells (Fig. 3) [140].

Other host proteins identified as ORF4 binding partners by yeast two-hybrid screening include cytochrome c oxidase subunit 8A (COX8A), lamin C, and small nuclear ribonucleoprotein polypeptide N (SNRPN) (Table 1) [138]. COX8A is the smallest component of COX (Complex IV), the inner membrane attached enzyme complex which participates in hydrogen (electron) transfer from the matrix to the intermembrane space in mitochondria [141]. Although COX8A deficiency can lead to severe impairment of COX stability, the relationship between COX8A and apoptosis is still elusive [142]. Lamin C belongs to type V intermediate filament. It is mainly detected in the nuclear lamina. It might participate in maintaining the shape of nucleus by providing nuclear interior or regulating gene expression via chromatin remodeling [143]. SNRPN is involved in pre-RNA processing including alternative splicing within the nucleus [144]. Further studies are needed to find out the biological significance of the interaction between PCV2 ORF4 and these proteins in host cells.

PCV2 ORF5

The ORF5 gene is about 180 bp in length. It is located within the ORF1 gene. Its mRNA transcript is synthesized in the same direction as ORF1 (Fig. 1) [145]. The function of ORF5 protein is not necessary for PCV2 replication. However, ORF5 contributes to the pathogenesis of PCV2 by inducing activation of NF κ B, induction of autophagy, and inhibition of type I IFN production in host cells [145-149].

An initial study of ORF5 function in host cells has revealed that ORF5 protein can induce ER stress, increase proinflammatory cytokine production by activation of NF κ B, and inhibit host cell growth by increasing the S-phase of the cell cycle [145]. A subsequent study has revealed that PCV2 ORF5 protein can induce ER stress and unfolded protein response via the RNA-activated protein kinase-like ER kinase (PERK), activating transcription factor 6 (ATF6) and inositol requiring enzyme 1 (IRE1) signaling pathways [146]. After induction of ER stress by ORF5, PERK-eIF2 α -ATF4 pathway can induce autophagy through autophagosome formation in host cells (Fig. 4) [147]. ORF5 expression in host cells can also enhance autophagy by inhibiting phosphorylation of mammalian target of rapamycin (mTOR) via activation of AMP-activated protein kinase (AMPK) pathway (Fig. 4) [147]. The reason why certain viruses can induce autophagy in host cells is still under debate. However, PCV2 has an advantage by inducing autophagy because PCV2 ORF5 also promotes viral replication [147, 148]. In addition, PCV2 ORF5 might enhance replication of PCV2 in host cells by inhibiting type I IFN production via transcriptional repression of genes facilitating type I IFN expression or by increasing the expression of RNF128 which down-regulates type I IFN production [149, 150]. Overall, PCV2 ORF5 can facilitate PCV2 replication in host cells by inducing autophagy and inhibiting host immune surveillance [145-150].

Five host proteins including cytochrome P450 1A1 (CYP1A1), glycoprotein-NMB (GPNMB), serine and arginine rich splicing Factor 3 (SRSF3), tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein β (YWHAB), and zinc finger protein 511 (ZNF511) have been found to be able to interact with PCV2 ORF5 by yeast two-hybrid screening (Table 1) [145]. Among them, both GPNMB and YWHAB can inhibit PCV2 replication by binding with ORF5 [151, 152]. GPNMB might be involved in cell proliferation and anti-inflammatory responses [153]. Further studies are needed to elucidate the mechanism of how GPNMB inhibits PCV2 replication after interaction with ORF5. YWHAB can inhibit the function of M-phase inducer phosphatase 2 (CDC25B) which facilitates cell proliferation [154]. By interacting with PCV2 ORF5, YWHAB can prevent PCV2-induced autophagy and ER stress-mediated apoptosis [152].

PCV2 ORF6

PCV2 ORF6 encoding region is located within ORF2. Its transcriptional direction is the same as that of ORF2 [34]. However, the detection of ORF6 encoding protein using an anti-ORF6 polyclonal antibody has confirmed that ORF6 can make its own polypeptide [34]. Therefore, it is important to know the regulatory mechanism of transcriptional variation between ORF2 and ORF6. There is no known host protein that can interact with ORF6. However, the replication of ORF6 deletion mutant of PCV2 is reduced in PK-15 cells compared to that of wild-type PCV2 [34]. Also, ORF6 possibly can regulate caspase activity and down-regulate the expression of proinflammatory cytokines such as type 1 IFN (IFN- β), IL-10, IL-12, and TNF- α in PK-15 cells based on an ORF6 deletion mutant study [34]. Further study is needed to determine which host protein can interact with PCV2 ORF6 and how PCV2 ORF6 modulates host cell physiology.

CONCLUSION

Interactions between PCV2 ORFs 1 ~ 6 and host proteins can trigger various phenotypic changes of host cells, including the induction of apoptosis and autophagy, or blocking anti-inflammatory responses. Currently, PCV2a vaccine can efficiently reduce the incidence and ameliorate the severity of PCVAD. The majority of PCV2 vaccines are based on inactivated PCV2 or subunit vaccines containing PCV2 capsid protein [5, 19]. However, these vaccines could not cover all genotypes of PCV2 infection and might be a driven source of antigenic variation of PCV2. In addition to targeting PCV2 capsid, the identification of specific ORFs that control PCV2 infection by modulating host signaling network will be helpful for designing novel anti-PCV2 drugs that can selectively inhibit multiplication of PCV2 in host cells. These novel anti-PCV2 drugs are expected to aid in controlling PCVAD, which threatens the pig industry. Further studies are needed to define the proposed role of PCV2 ORF7 ~ ORF11 or identify host proteins that interact with them.

Competing Interests

None of the other authors had any conflict of interests.

ACCEPTED

Acknowledgements

This study was supported by a grant from the Next-Generation Bio-Green 21 program (Project No. PJ01327101), Rural Development Administration, Republic of Korea.

ACCEPTED

Author's Contributions

Conceptualization: Chun T. Validation: Park SW, Park IB, Kang SJ, Bae J. Writing - original draft: Park SW, Park IB. Writing - review & editing: Park SW, Park IB, Kang SJ, Bae J, Chun T.

ACCEPTED

Ethics approval and consent to participate

There were no animal or human studies contained in this manuscript.

ACCEPTED

References

1. Nayar GP, Hamel A, Lin L. Detection and characterization of porcine circovirus associated with postweaning multisystemic wasting syndrome in pigs. *Can Vet J*. 1997;38:385-6.
2. Allan GM, McNeilly F, Kennedy S, Daft B, Clarke EG, Ellis JA, et al. Isolation of porcine circovirus-like viruses from pigs with a wasting disease in the USA and Europe. *J Vet Diagn Invest*. 1998;10:3-10. <https://doi.org/10.1177/104063879801000102>
3. Kekarainen T, Segalés J. Porcine circovirus 2 immunology and viral evolution. *Porcine Health Manag*. 2015;19;1:17. <https://doi.org/10.1186/s40813-015-0012-z>
4. Kekarainen T, McCullough K, Fort M, Fossum C, Segalés J, Allan GM. Immune responses and vaccine-induced immunity against Porcine circovirus type 2. *Vet Immunol Immunopathol*. 2010;136:185-93. <https://doi.org/10.1016/j.vetimm.2010.03.025>
5. Chae C. Commercial porcine circovirus type 2 vaccines: efficacy and clinical application. *Vet J*. 2012;194:151-7. <https://doi.org/10.1016/j.tvjl.2012.06.031>
6. Alarcon P, Wieland B, Mateus AL, Dewberry C. Pig farmers' perceptions, attitudes, influences and management of information in the decision-making process for disease control. *Prev Vet Med*. 2014;116:223-42. <https://doi.org/10.1016/j.prevetmed.2013.08.004>
7. Opriessnig T, Karuppannan AK, Halbur PG, Calvert JG, Nitzel GP, Matzinger SR, et al. Porcine circovirus type 2a or 2b based experimental vaccines provide protection against PCV2d/porcine parvovirus 2 co-challenge. *Vaccine*. 2020;38:1975–81. <https://doi.org/10.1016/j.vaccine.2020.01.013>
8. Xiao CT, Halbur PG, Opriessnig T. Complete genome sequence of a novel porcine circovirus type 2b variant present in cases of vaccine failures in the United States. *J Virol*. 2012;86:12412-69. <https://doi.org/10.1128/JVI.02345-12>
9. Seo HW, Park CH, Kang IJ, Choi KH, Jeong JW, Park SJ, et al. Genetic and antigenic characterization of a newly emerging porcine circovirus type 2b mutant first isolated in cases of vaccine failure in Korea. *Arch Virol*. 2014;159:3107-11. <https://doi.org/10.1007/s00705-014-2164-6>
10. Franzo G, Segalés J. Porcine circovirus 2 (PCV-2) genotype update and proposal of a new genotyping methodology. *PLoS One*. 2018;13:e0208585. <https://doi.org/10.1371/journal.pone.0208585>
11. Krakowka S, Allan G, Ellis J, Hamberg A, Charreyre C, Kaufmann E, et al. A nine-base nucleotide sequence in the porcine circovirus type 2 (PCV2) nucleocapsid gene determines viral replication and virulence. *Virus Res*. 2012; 164:90-9. <https://doi.org/10.1016/j.virusres.2011.10.027>

12. Shen HG, Halbur PG, Opriessnig T. Prevalence and phylogenetic analysis of the current porcine circovirus 2 genotypes after implementation of widespread vaccination programmes in the USA. *J Gen Virol.* 2012;93:1345-55. <https://doi.org/10.1099/vir.0.039552-0>
13. Bandrick M, Balasch M, Heinz A, Taylor L, King V, Toepfer J, et al. A bivalent porcine circovirus type 2 (PCV2), PCV2a-PCV2b, vaccine offers biologically superior protection compared to monovalent PCV2 vaccines. *Vet Res.* 2022;53:12. <https://doi.org/10.1186/s13567-022-01029-w>
14. Segalés J. Porcine circovirus type 2 (PCV2) infections: clinical signs, pathology and laboratory diagnosis. *Virus Res.* 2012;164:10-9. <https://doi.org/10.1016/j.virusres.2011.10.007>
15. Harding JC. The clinical expression and emergence of porcine circovirus 2. *Vet Microbiol.* 2004;98:131-5. <https://doi.org/10.1016/j.vetmic.2003.10.013>
16. Nielsen J, Vincent IE, Bøtner A, Ladekaer-Mikkelsen AS, Allan G, Summerfield A, et al. Association of lymphopenia with porcine circovirus type 2 induced postweaning multisystemic wasting syndrome (PMWS). *Vet Immunol Immunopathol.* 2003;92:97-111. [https://doi.org/10.1016/s0165-2427\(03\)00031-x](https://doi.org/10.1016/s0165-2427(03)00031-x)
17. Meng XJ. Porcine circovirus type 2 (PCV2): pathogenesis and interaction with the immune system. *Annu Rev Anim Biosci.* 2013;1:43-64. <https://doi.org/10.1146/annurev-animal-031412-103720>
18. Chathuranga K, Weerawardhana A, Dodantenna N, Lee JS. Regulation of antiviral innate immune signaling and viral evasion following viral genome sensing. *Exp Mol Med.* 2021;53:1647-68. <https://doi.org/10.1038/s12276-021-00691-y>
19. Tischer I, Gelderblom H, Vettermann W, Koch MA. A very small porcine virus with circular single-stranded-DNA. *Nature.* 1982;295:64-6. <https://doi.org/10.1038/295064a0>
20. Opriessnig T, Karuppannan AK, Castro AMMG, Xiao CT. Porcine circoviruses: current status, knowledge gaps and challenges. *Virus Res.* 2020;286:198044. <https://doi.org/10.1016/j.virusres.2020.198044>
21. Li J, Xing G, Zhang C, Yang H, Li G, Wang N, et al. Cross-species transmission resulted in the emergence and establishment of circovirus in pig. *Infect Genet Evol.* 2019;75:103973. <https://doi.org/10.1016/j.meegid.2019.103973>
22. Wang D, Mai J, Yang Y, Xiao CT, Wang N. Current knowledge on epidemiology and evolution of novel porcine circovirus 4. *Vet Res.* 2022;53:38. <https://doi.org/10.1186/s13567-022-01053-w>
23. Thomson J, Smith B, Allan G, McNeilly F, McVicar C. PDNS, PMWS and porcine circovirus type 2 in Scotland. Porcine dermatitis and nephropathy syndrome. Post-weaning multisystemic wasting syndrome. *Vet Rec.* 2000;146:651-2.

24. Opriessnig T, Meng XJ, Halbur PG. Porcine circovirus type 2 associated disease: update on current terminology, clinical manifestations, pathogenesis, diagnosis, and intervention strategies. *J Vet Diagn Invest.* 2007;19:591-615. <https://doi.org/10.1177/104063870701900601>
25. Rodriguez-Carino C, Segales, J. Ultrastructural findings in lymph nodes from pigs suffering from naturally occurring postweaning multisystemic wasting syndrome. *Vet Pathol.* 2009;46:729-35. <https://doi.org/10.1354/vp.08-VP-0141-R-FL>
26. Phan TG, Giannitti F, Rossow S, Marthaler D, Knutson TP, Li L, et al. Detection of a novel circovirus PCV3 in pigs with cardiac and multi-systemic inflammation. *Virol J.* 2016;13:184. <https://doi.org/10.1186/s12985-016-0642-z>
27. Palinski R, Piñeyro P, Shang PC, Yuan FF, Guo R, Fang Y, et al. A novel porcine circovirus distantly related to known circoviruses is associated with porcine dermatitis and nephropathy syndrome and reproductive failure. *J Virol.* 2016;91:e01879-16. <https://doi.org/10.1128/JVI.01879-16>
28. Jiang H, Wang D, Wang J, Zhu S, She R, Ren X, et al. Induction of porcine dermatitis and nephropathy syndrome in piglets by infection with porcine circovirus Type 3. *J Virol.* 2019;93:e02045-18. <https://doi.org/10.1128/JVI.02045-18>
29. Ge M, Hu WQ, Ning KM, Li SY, Xiao CT. The seroprevalence of the newly identified porcine circovirus type 4 in China investigated by an enzyme-linked immunosorbent assay. *Transbound Emerg Dis.* 2021;68:2910-4. <https://doi.org/10.1111/tbed.14184>
30. Hang HH, Hu WQ, Li JY, Liu TN, Zhou JY, Opriessnig T, et al. Novel circovirus species identified in farmed pigs designated as porcine circovirus 4, Hunan province, China. *Transbound Emerg Dis.* 2020;67:1057-61. <https://doi.org/10.1111/tbed.13446>
31. Sun WC, Du Q, Han ZX, Bi JS, Lan T, Wang W, et al. Detection and genetic characterization of porcine circovirus 4 (PCV4) in Guangxi, China. *Gene.* 2021;773:145384. <https://doi.org/10.1016/j.gene.2020.145384>
32. Hamel AL, Lin LL, Nayar GP. Nucleotide sequence of porcine circovirus associated with postweaning multisystemic wasting syndrome in pigs. *J Virol.* 1998;72:5262-7. <https://doi.org/10.1128/JVI.72.6.5262-67>
33. Lv QZ, Guo KK, Zhang YM. Current understanding of genomic DNA of porcine circovirus type 2. *Virus Genes.* 2014;49:1-10. <https://doi.org/10.1007/s11262-014-1099-z>
34. Li D, Wang J, Xu S, Cai S, Ao C, Fang L, et al. Identification and functional analysis of the novel ORF6 protein of porcine circovirus type 2 in vitro. *Vet Res Commun.* 2018;42:1-10. <https://doi.org/10.1007/s11259-017-9702-0>
35. Gu J, Zhang Y, Lian X, Sun H, Wang J, Liu W, et al. Functional analysis of the interferon-stimulated response element of porcine circovirus type 2 and its role during viral replication in vitro and in vivo. *Virol J.* 2012;9:152. <https://doi.org/10.1186/1743->

36. Nawagitgul P, Morozov I, Bolin SR, Harms PA, Sorden SD, Paul PS. Open reading frame 2 of porcine circovirus type 2 encodes a major capsid protein. *J Gen Virol.* 2000;81:2281-7. <https://doi.org/10.1099/0022-1317-81-9-2281>
37. Lin WL, Chien MS, Wu PC, Lai CL, Huang C. The porcine circovirus type 2 nonstructural protein ORF3 induces apoptosis in porcine peripheral blood mononuclear cells. *Open Virol J.* 2011;5:148-53. <https://doi.org/10.2174/1874357901105010148>
38. He J, Cao J, Zhou N, Jin Y, Wu J, Zhou J. Identification and functional analysis of the novel ORF4 protein encoded by porcine circovirus type 2. *J Virol.* 2013;87:1420-9. <https://doi.org/10.1128/JVI.01443-12>
39. Rakibuzzaman AG, Ramamoorthy S. Comparative immunopathogenesis and biology of recently discovered porcine circoviruses. *Transbound Emerg Dis.* 2021;68:2957-68. <https://doi.org/10.1111/tbed.14244>
40. Bolin SR, Stoffregen WC, Nayar GP, Hamel AL. Postweaning multisystemic wasting syndrome induced after experimental inoculation of cesarean-derived, colostrum-deprived piglets with type 2 porcine circovirus. *J Vet Diagn Invest.* 2001;13:185-94. <https://doi.org/10.1177/104063870101300301>
41. Shibata I, Okuda Y, Yazawa S, Ono M, Sasaki T, Itagaki M, et al. PCR detection of porcine circovirus type 2 DNA in whole blood, serum, oropharyngeal swab, nasal swab, and feces from experimentally infected pigs and field cases. *J Vet Med Sci.* 2003;65:405-8. <https://doi.org/10.1292/jvms.65.405>
42. Segalés J, Calsamiglia M, Olvera A, Sibila M, Badiella L, Domingo M. Quantification of porcine circovirus type 2 (PCV2) DNA in serum and tonsillar, nasal, tracheo-bronchial, urinary and faecal swabs of pigs with and without postweaning multisystemic wasting syndrome (PMWS). *Vet Microbiol.* 2005;111:223-9. <https://doi.org/10.1016/j.vetmic.2005.10.008>
43. McIntosh KA, Harding JCS, Parker S, Krakowka S, Allan G, Ellis JA. Quantitative polymerase chain reaction for porcine circovirus-2 in swine feces in a porcine circovirus disease-affected commercial herd and a nonaffected commercial herd. *Can Vet J.* 2008;49:1189-94.
44. Madec F, Rose N, Grasland B, Cariolet R, Jestin A. Post-weaning multisystemic wasting syndrome and other PCV2-related problems in pigs: a 12-year experience. *Transbound Emerg Dis.* 2008;55:273-83. <https://doi.org/10.1111/j.1865-1682.2008.01035.x>
45. Patterson AR, Ramamoorthy S, Madson DM, Meng XJ, Halbur PG, Opriessnig T. Shedding and infection dynamics of porcine circovirus type 2 (PCV2) after experimental infection. *Vet Microbiol.* 2011;149:91-8. <https://doi.org/10.1016/j.vetmic.2010.10.020>
46. Rose N, Opriessnig T, Grasland B, Jestin A. Epidemiology and transmission of porcine circovirus type 2 (PCV2). *Virus Res.* 2012;164:78-89.

47. Segales J, Allan GM, Domingo M. Porcine circovirus diseases. *Anim. Health Res Rev.* 2005;6:119–42. <https://doi.org/10.1079/ahr2005106>
48. Mateusen B, Maes DG, Van Soom A, Lefebvre D, Nauwynck HJ. Effect of a porcine circovirus type 2 infection on embryos during early pregnancy. *Theriogenology.* 2007;68:896–901. <https://doi.org/10.1016/j.theriogenology.2007.07.014>
49. Sanchez Jr RE, Meerts P, Nauwynck HJ, Pensaert MB. Change of porcine circovirus 2 target cells in pigs during development from fetal to early postnatal life. *Vet Microbiol.* 2003;95:15–25. [https://doi.org/10.1016/s0378-1135\(03\)00120-2](https://doi.org/10.1016/s0378-1135(03)00120-2)
50. Saha D, Lefebvre D, Van Doorselaere J, Atanasova K, Barbé F, Geldhof M, et al. Pathological and virological outcome in mid-gestational porcine foetuses after experimental inoculation with PCV2a and PCV2b. *Vet Microbiol.* 2010;145:62–8. <https://doi.org/10.1016/j.vetmic.2010.03.017>
51. Pensaert MB, Sanchez Jr RM, Ladekjær-Mikkelsen AS, Allan GM, Nauwynck HJ, et al. Viremia and effect of fetal infection with porcine viruses with special reference to porcine circovirus 2 infection. *Vet Microbiol.* 2004;98:175–83. <https://doi.org/10.1016/j.vetmic.2003.10.011>
52. Sanchez Jr RE, Meerts P, Nauwynck HJ, Ellis JA, Pensaert MB. Characteristics of porcine circovirus 2 replication in lymphoid organs of pigs inoculated in late-gestation or postnatally and possible relation to clinical and pathological outcome of infection. *J Vet Diag Invest.* 2004;16:168–78. <https://doi.org/10.1177/104063870401600301>
53. Lefebvre DJ, Meerts P, Costers S, Misinzo G, Barbé F, Van Reeth K, et al. 2008. Increased porcine circovirus type 2 replication in porcine leukocytes in vitro and in vivo by concanavalin A stimulation. *Vet Microbiol.* 2008;132:74–86. <https://doi.org/10.1016/j.vetmic.2008.05.004>
54. Ramamoorthy S, Meng XJ. Porcine circoviruses: a minuscule yet mammoth paradox. *Anim Health Res Rev.* 2009;10:1–20. <https://doi.org/10.1017/S1466252308001461>
55. Darwich L, Mateu E. Immunology of porcine circovirus type 2 (PCV2). *Virus Res.* 2012;164:61–7. <https://doi.org/10.1016/j.virusres.2011.12.003>
56. Darwich L, Segalés J, Mateu E. Pathogenesis of postweaning multisystemic wasting syndrome caused by porcine circovirus 2: an immune riddle. *Arch Virol.* 2004;149:857–74. <https://doi.org/10.1007/s00705-003-0280-9>
57. Meerts P, Misinzo G, McNeilly F, Nauwynck HJ. Replication kinetics of different porcine circovirus 2 strains in PK-15 cells, fetal cardiomyocytes and macrophages. *Arch Virol.* 2005;150:427–41. <https://doi.org/10.1007/s00705-004-0444-2>
58. Hirai T, Nunoya T, Ihara T, Saitoh T, Shibuya K, Nakamura K. Infectivity of porcine circovirus 1 and circovirus 2 in primary porcine hepatocyte and kidney cell cultures. *J*

59. Jensen TK, Vigre H, Svensmark B, Bille-Hansen V. Distinction between porcine circovirus type 2 enteritis and porcine proliferative enteropathy caused by *Lawsonia intracellularis*. J Comp Pathol. 2006;135:176-82. <https://doi.org/10.1016/j.jcpa.2006.08.006>
60. Yu S, Opriessnig T, Kitikoon P, Nilubol D, Halbur PG, Thacker E. Porcine circovirus type 2 (PCV2) distribution and replication in tissues and immune cells in early infected pigs. Vet Immunol Immunopathol. 2007;115:262-72. <https://doi.org/10.1016/j.vetimm.2006.11.006>
61. Steiner E, Balmelli C, Herrmann B, Summerfield A, McCullough K. Porcine circovirus type 2 displays pluripotency in cell targeting. Virology. 2008;378:311-22. <https://doi.org/10.1016/j.virol.2008.06.009>
62. Weingartl HM, Sabara M, Pasick J, Van Moorlehem E, Babiuk L. Continuous porcine cell lines developed from alveolar macrophages: partial characterization and virus susceptibility. J Virol Methods. 2002;104:203-16. [https://doi.org/10.1016/s0166-0934\(02\)00085-x](https://doi.org/10.1016/s0166-0934(02)00085-x)
63. Dulac GC, Afshar A. Porcine circovirus antigens in PK-15 cell line (ATCC CCL-33) and evidence of antibodies to circovirus in Canadian pigs. Can J Vet Res. 1989;53:431-3.
64. Fenaux M, Halbur PG, Haqshenas G, Royer R, Thomas P, Nawagitgul P, et al. Cloned genomic DNA of type 2 porcine circovirus is infectious when injected directly into the liver and lymph nodes of pigs: characterization of clinical disease, virus distribution, and pathologic lesions. J Virol. 2002;76:541-51. <https://doi.org/10.1128/jvi.76.2.541-551.2002>
65. Misinzo G, Meerts P, Bublot M, Mast J, Weingartl HM, Nauwynck HJ. Binding and entry characteristics of porcine circovirus 2 in cells of the porcine monocytic line 3D4/31. J Gen Virol. 2005;86:2057-68. <https://doi.org/10.1099/vir.0.80652-0>
66. Misinzo G, Delputte PL, Meerts P, Lefebvre DJ, Nauwynck HJ. Porcine circovirus 2 uses heparan sulfate and chondroitin sulfate B glycosaminoglycans as receptors for its attachment to host cells. J Virol. 2006;80:3487-94. <https://doi.org/10.1128/JVI.80.7.3487-3494.2006>
67. Khayat R, Brunn N, Speir JA, Hardham JM, Ankenbauer RG, Schneemann A, et al. The 2.3-angstrom structure of porcine circovirus 2. J Virol. 2011;85:7856-62. <https://doi.org/10.1128/JVI.00737-11>
68. Misinzo G, Delputte P, Nauwynck HJ. Inhibition of endosomal-lysosomal system acidification enhances porcine circovirus 2 infection of porcine epithelial cells. J Virol. 2008;82:1128-35. <https://doi.org/10.1128/JVI.01229-07>
69. Cheung AK. Porcine circovirus: transcription and DNA replication. Virus Res. 2012;164:46-53 <https://doi.org/10.1016/j.virusres.2011.10.012>

70. Segalés J, Pastor J, Cuenca R, Domingo M. Haematological parameters in postweaning multisystemic wasting syndrome-affected pigs. *Vet Rec.* 2000;146:675-6. <https://doi.org/10.1136/vr.146.23.675>
71. Segalés J, Alonso F, Rosell C, Pastor J, Chianini F, Campos E, et al. Changes in peripheral blood leukocyte populations in pigs with natural postweaning multisystemic wasting syndrome (PMWS). *Vet Immunol Immunopathol.* 2001 Aug;81:37-44. [https://doi.org/10.1016/s0165-2427\(01\)00326-9](https://doi.org/10.1016/s0165-2427(01)00326-9)
72. Darwich L, Segalés J, Domingo M, Mateu E. Changes in CD4⁺, CD8⁺, CD4⁺ CD8⁺, and immunoglobulin M-positive peripheral blood mononuclear cells of postweaning multisystemic wasting syndrome-affected pigs and age-matched uninfected wasted and healthy pigs correlate with lesions and porcine circovirus type 2 load in lymphoid tissues. *Clin Diagn Lab Immunol.* 2002;9:236-42. <https://doi.org/10.1128/cdli.9.2.236-242.2002>
73. Sorden SD. Update on porcine circovirus and postweaning multisystemic wasting syndrome (PMWS). *Swine Health Prod.* 2000;8:133-6.
74. Quintana J, Segalés J, Rosell C, Calsamiglia M, Rodríguez-Arrioja GM, Chianini F, et al. Clinical and pathological observations on pigs with postweaning multisystemic wasting syndrome. *Vet Rec.* 2001;149:357-61. <https://doi.org/10.1136/vr.149.12.357>
75. Chianini F, Majó N, Segalés J, Domínguez J, Domingo M. Immunohistochemical characterisation of PCV2 associate lesions in lymphoid and non-lymphoid tissues of pigs with natural postweaning multisystemic wasting syndrome (PMWS). *Vet Immunol Immunopathol.* 2003;94:63-75. [https://doi.org/10.1016/s0165-2427\(03\)00079-5](https://doi.org/10.1016/s0165-2427(03)00079-5)
76. Carrasco L, Segalés J, Bautista MJ, Gómez-Villamandos JC, Rosell C, Ruiz-Villamor E, et al. Intestinal chlamydial infection concurrent with postweaning multisystemic wasting syndrome in pigs. *Vet Rec.* 2000;146:21-3. <https://doi.org/10.1136/vr.146.1.21>
77. Núñez A, McNeilly F, Perea A, Sánchez-Cordón PJ, Huerta B, Allan G, et al. Coinfection by *Cryptosporidium parvum* and porcine circovirus type 2 in weaned pigs. *J Vet Med B Infect Dis Vet Public Health.* 2003;50:255-8. <https://doi.org/10.1046/j.1439-0450.2003.00664.x>
78. Timmusk S, Fossum C, Berg M. Porcine circovirus type 2 replicase binds the capsid protein and an intermediate filament-like protein. *J Gen Virol.* 2006;87:3215-23. <https://doi.org/10.1099/vir.0.81785-0>
79. Finsterbusch T, Steinfeldt T, Doberstein K, Rödner C, Mankertz A. Interaction of the replication proteins and the capsid protein of porcine circovirus type 1 and 2 with host proteins. *Virology.* 2009;386:122-31. <https://doi.org/10.1016/j.virol.2008.12.039>
80. Fan LL, Yuan ZZ, Jiao ZJ, Zeng H, Yang YF, Tan ZP. Haploinsufficiency of syncoilin leads to hypertrophic cardiomyopathy. *Genes Dis.* 2022;9:1412-4. <https://doi.org/10.1016/j.gendis.2022.02.011>

81. Van Rossen E, Liu Z, Blijweert D, Eysackers N, Mannaerts I, Schroyen B, et al. Syncoilin is an intermediate filament protein in activated hepatic stellate cells. *Histochem Cell Biol.* 2014;141:85-99. <https://doi.org/10.1007/s00418-013-1142-5>
82. Parlakian A, Paulin D, Izmiryan A, Xue Z, Li Z. Intermediate filaments in peripheral nervous system: Their expression, dysfunction and diseases. *Rev Neurol (Paris).* 2016;172:607-13. <https://doi.org/10.1016/j.neurol.2016.07.015>
83. Liang JJ, Yu CY, Liao CL, Lin YL. Vimentin binding is critical for infection by the virulent strain of Japanese encephalitis virus. *Cell Microbiol.* 2011;13:1358-70. <https://doi.org/10.1111/j.1462-5822.2011.01624.x>
84. Roy S, Kapoor A, Zhu F, Mukhopadhyay R, Ghosh AK, Lee H, et al. Artemisinins target the intermediate filament protein vimentin for human cytomegalovirus inhibition. *J Biol Chem.* 2020;295:15013-28. <https://doi.org/10.1074/jbc.RA120.014116>
85. Ramos I, Stamatakis K, Oeste CL, Pérez-Sala D. Vimentin as a Multifaceted Player and Potential Therapeutic Target in Viral Infections. *Int J Mol Sci.* 2020 ;21:4675. <https://doi.org/10.3390/ijms21134675>
86. Hölzel M, Kohlhuber F, Schlosser I, Hölzel D, Lüscher B, Eick D. Myc/Max/Mad regulate the frequency but not the duration of productive cell cycles. *EMBO Rep.* 2001;2:1125-32. <https://doi.org/10.1093/embo-reports/kve251>
87. Dominguez-Sola D, Ying CY, Grandori C, Ruggiero L, Chen B, Li M, et al. Non-transcriptional control of DNA replication by c-Myc. *Nature.* 2007;448:445-51. <https://doi.org/10.1038/nature05953>
88. Dominguez-Sola D, Gautier J. MYC and the control of DNA replication. *Cold Spring Harb Perspect Med.* 2014;4:a014423. <https://doi.org/10.1101/cshperspect.a014423>
89. Neddermann P, Gallinari P, Lettieri T, Schmid D, Truong O, Hsuan JJ, et al. Cloning and expression of human G/T mismatch-specific thymine-DNA glycosylase. *J Biol Chem.* 1996;271(22):12767-74. <https://doi.org/10.1074/jbc.271.22.12767>
90. Tian XL, Kadaba R, You SA, Liu M, Timur AA, Yang L, et al. Identification of an angiogenic factor that when mutated causes susceptibility to Klippel-Trenaunay syndrome. *Nature.* 2004;427:640-5. <https://doi.org/10.1038/nature02320>
91. Yao HH, Wang BJ, Wu Y, Huang Q. High expression of angiogenic factor with G-patch and FHA domain 1 (AGGF1) predicts poor prognosis in gastric cancer. *Med Sci Monit.* 2017;23:1286-94. <https://doi.org/10.12659/msm.903248>
92. Wang W, Li GY, Zhu JY, Huang DB, Zhou HC, Zhong W, et al. Overexpression of AGGF1 is correlated with angiogenesis and poor prognosis of hepatocellular carcinoma. *Med Oncol.* 2015;32:131. <https://doi.org/10.1007/s12032-015-0574-2>
93. Adams DJ, van der Weyden L, Mayeda A, Stamm S, Morris BJ, Rasko JE. ZNF265--a novel spliceosomal protein able to induce alternative splicing. *J Cell Biol.* 2001;154:25-

32. <https://doi.org/10.1083/jcb.200010059>
94. Sako D , Chang XJ, Barone KM, Vachino G, White HM, Shaw G, et al. Expression cloning of a functional glycoprotein ligand for P-selectin. *Cell*. 1993;75:1179-86. [https://doi.org/10.1016/0092-8674\(93\)90327-m](https://doi.org/10.1016/0092-8674(93)90327-m)
95. McEver RP, Beckstead JH, Moore KL, Marshall-Carlson L, Bainton DF. GMP-140, a platelet alpha-granule membrane protein, is also synthesized by vascular endothelial cells and is localized in Weibel-Palade bodies. *J Clin Invest*. 1989;84:92-9. <https://doi.org/10.1172/JCI114175>
96. Kononchik J, Ireland J, Zou Z, Segura J, Holzapfel G, Chastain A, et al. HIV-1 targets L-selectin for adhesion and induces its shedding for viral release. *Nat Commun*. 2018;9:2825. <https://doi.org/10.1038/s41467-018-05197-2>
97. Ghebrehwet B, Lim BL, Kumar R, Feng X, Peerschke EI. gC1q-R/p33, a member of a new class of multifunctional and multicompartmental cellular proteins, is involved in inflammation and infection. *Immunol Rev*. 2001;180:65-77. <https://doi.org/10.1034/j.1600-065x.2001.1800106.x>
98. Dembitzer FR, Kinoshita Y, Burstein D, Phelps RG, Beasley MB, Garcia R, et al. gC1qR expression in normal and pathologic human tissues: differential expression in tissues of epithelial and mesenchymal origin. *J Histochem Cytochem*. 2012;60:467-74. <https://doi.org/10.1369/0022155412440882>
99. Peerschke EI, Reid KB, Ghebrehwet B. Platelet activation by C1q results in the induction of α IIb/ β 3 integrins (GPIIb-IIIa) and the expression of P-selectin and procoagulant activity. *J Exp Med*. 1993;178:579-87. <https://doi.org/10.1084/jem.178.2.579>
100. Ghebrehwet B, Kew RR, Gruber BL, Marchese MJ, Peerschke EI, Reid KB. Murine mast cells express two types of C1q receptors that are involved in the induction of chemotaxis and chemokinesis. *J Immunol*. 1995;155:2614-9.
101. Guan EN, Burgess WH, Robinson SL, Goodman EB, McTigue KJ, Tenner AJ. Phagocytic cell molecules that bind the collagen-like region of C1q. Involvement in the C1q-mediated enhancement of phagocytosis. *J Biol Chem*. 1991;266: 20345-55.
102. Goodman EB, Tenner AJ. Signal transduction mechanisms of C1q-mediated superoxide production. Evidence for the involvement of temporally distinct staurosporine-insensitive and sensitive pathways. *J Immunol*. 1992;148:3920-8.
103. Choi CY, Oh HN, Jun Lee S, Chun T. ORF2 protein of porcine circovirus type 2 promotes phagocytic activity of porcine macrophages by inhibiting proteasomal degradation of complement component 1, q subcomponent binding protein (C1QBP) through physical interaction. *J Gen Virol*. 2015;96:3294-301. <https://doi.org/10.1099/jgv.0.000282>
104. Du Q, Huang Y, Wang T, Zhang X, Chen Y, Cui B, et al. Porcine circovirus type 2

- activates PI3K/Akt and p38 MAPK pathways to promote interleukin-10 production in macrophages via Cap interaction of gC1qR. *Oncotarget*. 2016;7:17492-507. <https://doi.org/10.18632/oncotarget.7362>
105. Liu Q, Liang C, Zhou L. Structural and functional analysis of the Hsp70/Hsp40 chaperone system. *Protein Sci*. 2020;29:378-90. <https://doi.org/10.1002/pro.3725>
106. Chaari A. Molecular chaperones biochemistry and role in neurodegenerative diseases. *Int J Biol Macromol*. 2019;131:396-411. <https://doi.org/10.1016/j.ijbiomac.2019.02.148>
107. Liu J, Bai J, Zhang L, Jiang Z, Wang X, Li Y, et al. Hsp70 positively regulates porcine circovirus type 2 replication in vitro. *Virology*. 2013;447:52-62. <https://doi.org/10.1016/j.virol.2013.08.025>
108. Böhne A, Darras A, D'Cotta H, Baroiller JF, Galiana-Arnoux D, Volff JN. The vertebrate makorin ubiquitin ligase gene family has been shaped by large-scale duplication and retroposition from an ancestral gonad-specific, maternal-effect gene. *BMC Genomics*. 2010;11:721. <https://doi.org/10.1186/1471-2164-11-721>
109. Kim JH, Park SM, Kang MR, Oh SY, Lee TH, Muller MT, et al. Ubiquitin ligase MKRN1 modulates telomere length homeostasis through a proteolysis of hTERT. *Genes Dev*. 2005;19:776–81. <https://doi.org/10.1101/gad.1289405>
110. Lee EW, Lee MS, Camus S, Ghim J, Yang MR, Oh W, et al. Differential regulation of p53 and p21 by MKRN1 E3 ligase controls cell cycle arrest and apoptosis. *EMBO J*. 2009;28:2100-13. <https://doi.org/10.1038/emboj.2009.164>
111. Ito T, Bulger M, Kobayashi M, Kadonaga JT. Drosophila NAP-1 is a core histone chaperone that functions in ATP-facilitated assembly of regularly spaced nucleosomal arrays. *Mol Cell Biol*. 1996;16:3112-24. <https://doi.org/10.1128/MCB.16.6.3112>
112. Moshkin YM, Kan TW, Goodfellow H, Bezstarosti K, Maeda RK, Pilyugin M, et al. Histone chaperones ASF1 and NAP1 differentially modulate removal of active histone marks by LID-RPD3 complexes during NOTCH silencing. *Mol Cell*. 2009;35:782-93. <https://doi.org/10.1016/j.molcel.2009.07.020>
113. Borer RA, Lehner CF, Eppenberger HM, Nigg EA. Major nucleolar proteins shuttle between nucleus and cytoplasm. *Cell*. 1989;56:379-90. [https://doi.org/10.1016/0092-8674\(89\)90241-9](https://doi.org/10.1016/0092-8674(89)90241-9)
114. Grisendi S, Mecucci C, Falini B, Pandolfi PP. Nucleophosmin and cancer. *Nat Rev Cancer*. 2006;6:493-505. <https://doi.org/10.1038/nrc1885>
115. Chaudhry P, Singh M, Parent S, Asselin E. Prostate apoptosis response 4 (Par-4), a novel substrate of caspase-3 during apoptosis activation. *Mol Cell Biol*. 2012;32:826-39. <https://doi.org/10.1128/MCB.06321-11>
116. Zhou J, Li H, Yu T, Li J, Dong W, Ojha NK, et al. Protein interactions network of porcine circovirus type 2 capsid with host proteins. *Front Microbiol*. 2020;11:1129.

117. Gustafson EA, Wessel GM. DEAD-box helicases: posttranslational regulation and function. *Biochem Biophys Res Commun.* 2010;395:1-6. <https://doi.org/10.1016/j.bbrc.2010.02.172>
118. Lu WT, Wilczynska A, Smith E, Bushell M. The diverse roles of the eIF4A family: you are the company you keep. *Biochem Soc Trans.* 2014;42:166-72. <https://doi.org/10.1042/BST20130161>
119. Mo L, Meng L, Huang Z, Yi L, Yang N, Li G. An analysis of the role of HnRNP C dysregulation in cancers. *Biomark Res.* 2022;10:19. <https://doi.org/10.1186/s40364-022-00366-4>
120. Deane R, Schäfer W, Zimmermann HP, Mueller L, Görlich D, Prehn S, et al. Ran-binding protein 5 (RanBP5) is related to the nuclear transport factor importin-beta but interacts differently with RanBP1. *Mol Cell Biol.* 1997;17:5087-96. <https://doi.org/10.1128/MCB.17.9.5087>
121. Zhou W, Gross KM, Kuperwasser C. Molecular regulation of Snai2 in development and disease. *J Cell Sci.* 2019;132:jcs235127. <https://doi.org/10.1242/jcs.235127>
122. Davis III AE, Mejia P, Lu F. Biological activities of C1 inhibitor. *Mol Immunol.* 2008;45:4057-63. <https://doi.org/10.1016/j.molimm.2008.06.028>
123. Haller O, Kochs G. Interferon-induced mx proteins: dynamin-like GTPases with antiviral activity. *Traffic.* 2002;3:710-7. <https://doi.org/10.1034/j.1600-0854.2002.31003.x>
124. Liu J, Chen I, Kwang J. Characterization of a previously unidentified viral protein in porcine circovirus type 2-infected cells and its role in virus-induced apoptosis. *J Virol.* 2005;79:8262-74. <https://doi.org/10.1128/JVI.79.13.8262-8274.2005>
125. Mankertz A, Caliskan R, Hattermann K, Hillenbrand B, Kurzendoerfer P, Bettina M, et al. Molecular biology of porcine circovirus: analyses of gene expression and viral replication. *Vet Microbiol.* 2004;98:81-8. <https://doi.org/10.1016/j.vetmic.2003.10.014>
126. Teras M, Viisileht E, Pahtma-Hall M, Rump A, Paalme V, Pata P, et al. Porcine circovirus type 2 ORF3 protein induces apoptosis in melanoma cells. *BMC Cancer.* 2018;18:1237. <https://doi.org/10.1186/s12885-018-5090-2>
127. Liu J, Zhu Y, Chen I, Lau J, He F, Lau A, et al. The ORF3 protein of porcine circovirus type 2 interacts with porcine ubiquitin E3 ligase Pirh2 and facilitates p53 expression in viral infection. *J Virol.* 2007;81:9560-7. <https://doi.org/10.1128/JVI.00681-07>
128. Karuppannan AK, Liu S, Jia Q, Selvaraj M, Kwang J. Porcine circovirus type 2 ORF3 protein competes with p53 in binding to Pirh2 and mediates the deregulation of p53 homeostasis. *Virology.* 2010;398:1–11. <https://doi.org/10.1016/j.virol.2009.11.028>

129. Liu J, Chen I, Du Q, Chua H, Kwang J. The ORF3 protein of porcine circovirus type 2 is involved in viral pathogenesis in vivo. *J Virol.* 2006;80:5065-73. <https://doi.org/10.1128/JVI.80.10.5065-5073.2006>
130. Karuppannan AK, Jong MH, Lee SH, Zhu Y, Selvaraj M, Lau J, et al. Attenuation of porcine circovirus 2 in SPF piglets by abrogation of ORF3 function. *Virology.* 2009;383:338-47. <https://doi.org/10.1016/j.virol.2008.10.024>
131. Juhan NM, LeRoith T, Opriessnig T, Meng XJ. The open reading frame 3 (ORF3) of porcine circovirus type 2 (PCV2) is dispensable for virus infection but evidence of reduced pathogenicity is limited in pigs infected by an ORF3-null PCV2 mutant. *Virus Res.* 2010;147:60-6. <https://doi.org/10.1016/j.virusres.2009.10.007>
132. Timmusk S, Merlot E, Lövgren T, Järvekülg L, Berg M, Fossum C. Regulator of G protein signalling 16 is a target for a porcine circovirus type 2 protein. *J Gen Virol.* 2009;90:2425-36. <https://doi.org/10.1099/vir.0.008896-0>
133. Choi CY, Rho SB, Kim HS, Han J, Bae J, Lee SJ, et al. The ORF3 protein of porcine circovirus type 2 promotes secretion of IL-6 and IL-8 in porcine epithelial cells by facilitating proteasomal degradation of regulator of G protein signalling 16 through physical interaction. *J Gen Virol.* 2015;96:1098-108. <https://doi.org/10.1099/vir.0.000046>
134. Baldwin Jr AS. The NF- κ B and I κ B proteins: new discoveries and insights. *Annu Rev Immunol.* 1996;14:649-681. <https://doi.org/10.1146/annurev.immunol.14.1.649>
135. Vasilatos SN, Katz TA, Oesterreich S, Wan Y, Davidson NE, Huang Y. Crosstalk between lysine-specific demethylase 1 (LSD1) and histone deacetylases mediates antineoplastic efficacy of HDAC inhibitors in human breast cancer cells. *Carcinogenesis.* 2013;34:1196-207. <https://doi.org/10.1093/carcin/bgt033>
136. He J, Cao J, Zhou N, Jin Y, Wu J, Zhou J. Identification and functional analysis of the novel ORF4 protein encoded by porcine circovirus type 2. *J Virol.* 2013;87:1420-29. <https://doi.org/10.1128/JVI.01443-12>
137. Gao Z, Dong Q, Jiang Y, Opriessnig T, Wang J, Quan Y, et al. ORF4-protein deficient PCV2 mutants enhance virus-induced apoptosis and show differential expression of mRNAs in vitro. *Virus Res.* 2014;183:56-62. <https://doi.org/10.1016/j.virusres.2014.01.024>
138. Lv Q, Guo K, Wang T, Zhang C, Zhang Y. Porcine circovirus type 2 ORF4 protein binds heavy chain ferritin. *J Biosci.* 2015;40:477-85. <https://doi.org/10.1007/s12038-015-9551-3>
139. Pham CG, Bubici C, Zazzeroni F, Papa S, Jones J, Alvarez K, et al. Ferritin heavy chain upregulation by NF- κ B inhibits TNF α -induced apoptosis by suppressing reactive oxygen species. *Cell.* 2004;119:529-42. <https://doi.org/10.1016/j.cell.2004.10.017>
140. Lin C, Gu J, Wang H, Zhou J, Li J, Wang S, et al. Caspase-dependent apoptosis

- induction via viral protein ORF4 of porcine circovirus 2 binding to mitochondrial adenine nucleotide translocase 3. *J Virol.* 2018;92:e00238-18. <https://doi.org/10.1128/JVI.00238-18>
141. Rizzuto R, Nakase H, Darras B, Francke U, Fabrizi GM, et al. A gene specifying subunit VIII of human cytochrome c oxidase is localized to chromosome 11 and is expressed in both muscle and non-muscle tissues. *J Biol Chem.* 1989;264:10595-600.
 142. Hallmann K, Kudin AP, Zsurka G, Kornblum C, Reimann J, Stüve B, et al. Loss of the smallest subunit of cytochrome c oxidase, COX8A, causes Leigh-like syndrome and epilepsy. *Brain.* 2016;139:338-45. <https://doi.org/10.1093/brain/awv357>
 143. Naetar N, Ferraioli S, Foisner R. Lamins in the nuclear interior-life outside the lamina. *J Cell Sci.* 2017;130:2087-96. <https://doi.org/10.1242/jcs.203430>
 144. Lee MS, Lin YS, Deng YF, Hsu WT, Shen CC, Cheng YH, et al. Modulation of alternative splicing by expression of small nuclear ribonucleoprotein polypeptide N. *FEBS J.* 2014;281:5194-207. <https://doi.org/10.1111/febs.13059>
 145. Lv Q, Guo K, Xu H, Wang T, Zhang Y. Identification of Putative ORF5 Protein of Porcine circovirus type 2 and functional analysis of GFP-fused ORF5 protein. *PLoS One.* 2015;10:e0127859. <https://doi.org/10.1371/journal.pone.0127859>
 146. Ouyang Y, Xu L, Lv J, Hou Y, Fan Z, Xu P, et al. Porcine circovirus type 2 ORF5 protein induces endoplasmic reticulum stress and unfolded protein response in porcine alveolar macrophages. *Arch Virol.* 2019;164:1323-34. <https://doi.org/10.1007/s00705-019-04185-1>
 147. Lv J, Jiang Y, Feng Q, Fan Z, Sun Y, Xu P, et al. Porcine circovirus type 2 ORF5 protein induces autophagy to promote viral replication via the PERK-eIF2 α -ATF4 and mTOR-ERK1/2-AMPK signaling pathways in PK-15 Cells. *Front Microbiol.* 2020;11:320. <https://doi.org/10.3389/fmicb.2020.00320>
 148. Chen T, Tu S, Ding L, Jin M, Chen H, Zhou H. The role of autophagy in viral infections. *J Biomed Sci.* 2023;30:5. <https://doi.org/10.1186/s12929-023-00899-2>
 149. Choi CY, Choi YC, Park IB, Lee CH, Kang SJ, Chun T. The ORF5 protein of porcine circovirus type 2 enhances viral replication by dampening type I interferon expression in porcine epithelial cells. *Vet Microbiol.* 2018;226:50-58. <https://doi.org/10.1016/j.vetmic.2018.10.005>
 150. Kang SJ, Park IB, Chun T. Open reading frame 5 protein of porcine circovirus type 2 induces RNF128 (GRAIL) which inhibits mRNA transcription of interferon- β in porcine epithelial cells. *Res Vet Sci.* 2021;140:79-82. <https://doi.org/10.1016/j.rvsc.2021.08.002>
 151. Guo K, Xu L, Wu M, Hou Y, Jiang Y, Lv J, et al. A host factor GPNMB restricts porcine circovirus Type 2 (PCV2) replication and interacts with PCV2 ORF5 protein. *Front Microbiol.* 2019;9:3295. <https://doi.org/10.3389/fmicb.2018.03295>

152. Guo K, Zhang X, Hou Y, Liu J, Feng Q, Wang K, et al. A novel PCV2 ORF5-interacting host factor YWHAB inhibits virus replication and alleviates PCV2-induced cellular response. *Vet Microbiol.* 2020;251:108893. <https://doi.org/10.1016/j.vetmic.2020.108893>
153. Lazaratos AM, Annis MG, Siegel PM. GPNMB: a potent inducer of immunosuppression in cancer. *Oncogene.* 2022;41:4573-90. <https://doi.org/10.1038/s41388-022-02443-2>
154. Leal MF, Ribeiro HF, Rey JA, Pinto GR, Smith MC, Moreira-Nunes CA, et al. YWHAE silencing induces cell proliferation, invasion and migration through the up-regulation of CDC25B and MYC in gastric cancer cells: new insights about YWHAE role in the tumor development and metastasis process. *Oncotarget.* 2016;7:85393-410. <https://doi.org/10.18632/oncotarget.13381>

ACCEPTED

Figure Legends




Figure 1

Fig. 1. Schematic diagram of PCV2 genome. The circular genome of PCV2 is enlarged with each linearized open reading frame (ORF) [32-39]. Each ORF is illustrated as different colors of lines with arrow indicating the transcriptional orientation of each ORF. Predicted molecular weight of each ORF-encoded protein is denoted with the unit of kDa. ORFs 1, 2, and 4-encoded proteins contain putative glycosylation sites indicated by an asterisk. An arrow of stem loop structures marks the replication origin and used as a starting point for numbering of nucleotide sequences.




Fig. 2. PCV2 ORF3-mediated apoptosis and inflammatory responses. (A) PCV2 ORF3-mediated apoptosis [127, 128]. In the steady state, Pirh2, an E3 ubiquitin ligase, can downregulate p53 expression via ubiquitin-dependent proteasomal degradation. During PCV2 infection, ORF3 can interfere with Pirh2 function by binding to Pirh2. As a result,

apoptosis is induced by p53 target gene expression via increased half-life of p53. (B) PCV2 ORF3-mediated inflammatory responses [133-135]. In the steady state, RGS16 can block nuclear entry of NF κ B. During PCV2 infection, ORF3 can cause proteasomal degradation of RGS16 by increasing ubiquitination. Subsequently, NF κ B can migrate into the nucleus and initiate mRNA transcription of IL-6 and IL-8.

ACCEPTED




Figure 3

Fig. 3. PCV2 ORF4 facilitates host cell apoptosis by inducing mitochondrial damage. During PCV2 infection, PCV2 ORF4 can directly bind to ANT3 through its *N*-terminus amphipathic α -helix region. By assistance of ANT3, mitochondria anchored PCV2 ORF4 can damage mitochondria. Subsequently, cytochrome c is released from mitochondria which triggers caspase pathway and leads to host cell apoptosis [140].




Figure 4

Fig. 4. Mechanism of PCV2 ORF5-mediated autophagy induction. During PCV2 infection, PCV2 ORF5 can cause ER stress. Subsequently, PERK-eIF2α-ATF4 pathway can induce autophagosome formation [147]. ER stress can also lead to activation of AMPK pathway which can enhance autophagy by inhibiting phosphorylation of mTOR [147].

Table 1. Interactions of host cellular proteins with the ORFs of PCV2

PCV2 ORF	Host cell protein	Function of host cell protein	Biological meaning between specific ORF and host cell protein	Reference
ORF1 (Rep)	Syncoilin	Intermediate filament protein involved in transport processes	Transports viral proteins to subcellular organelles in host cells	[78]
	c-myc	Transcription factor which promotes cell cycle progress and DNA replication	Initiates viral DNA replication	[78]
	TDG	Thymidine DNA glycosylase associated in G/T mismatch repair	Participates in viral DNA replication	[79]
	VG5Q	Angiogenic factor which regulates vascular development	unknown	[79]
	ZNF265	Subunit of pre-mRNA spliceosome required for alternative splicing	Affects transcription and viral mRNA splicing	[79]
ORF2 (Cap)	P-selectin	Cell adhesion molecule involved in an interaction between endothelial cells and leukocytes	Facilitates viral attachment and entry into host cells	[78]
	C1qBP	Complement factor which binds with the globular heads of C1q	Mediates phagocytosis and production of proinflammatory cytokines	[78,79,103,104]
	Hsp40	Chaperone which delivers unfolded and misfolded proteins to Hsp70	Assists Hsp70 to enhance progeny virus production	[79]
	Hsp70	Chaperone involved in protein folding	Increases progeny virus production by elevating PCV2 genome replication	[107]
	MKRN1	RNA-binding E3 ubiquitin ligase which modulates cell cycle or senescence	unknown	[79]
	NAP1	Histone chaperone which participates in nucleosome assembly	unknown	[79]
	NPM1	Nucleolar protein shuttling between nucleus and cytoplasm	unknown	[79,116]
	Par-4	Caspase-3 substrate	Promotes apoptosis in host cells	[79]
	DDX21	RNA helicase involved in ribosomal RNA processing	unknown	[116]
	EIF4A2	RNA helicase which is a component of eIF4F complex	unknown	[116]
	hnRNPC	RNA-binding protein which regulates RNA homeostasis	unknown	[116]
	IC1	Protease inhibitor which has anti-inflammatory effects	Participates in local inflammation	[116]
	IPO5	Nuclear transport receptor which belongs to importin beta family	unknown	[116]
	MX2	IFN-induced GTPase interfering with viral capsid transport	Blocks the progeny virus production	[116]
	SNAI2	Transcription factor upregulating cell differentiation and migration	unknown	[116]

ORF3	Pirh2	E3 ubiquitin ligase involved in proteasomal degradation of p53	Causes apoptosis by increasing half-life of p53	[127,128]
	DDE-like transposase	Transposase catalyzing the movement of DNA segments	unknown	[78]
	RGS16	G protein signaling inhibitor which suppresses NFκB translocation	Promotes inflammatory responses by degradation of RGS16	[78,132,133]
ORF4	FHC	Ferroxidase which removes reactive oxygen intermediates	Rescues host cells from apoptosis	[138]
	ANT3	Translocase that mediates ADP/ATP exchange in mitochondrial matrix and cytosol	Leads to apoptosis by mitochondrial damage	[140]
	COX8A	Component of COX involved in the hydrogen transfer in mitochondria	unknown	[138]
	SNRPN	Polypeptide associated with pre-mRNA splicing	unknown	[138]
	Lamin C	Type V intermediate filament protein which is a major component of the nuclear lamina	unknown	[138]
ORF5	CYP1A1	Member of the cytochrome P450 which functions as monooxygenases	unknown	[145]
	GPNMB	Type I transmembrane glycoprotein associated with cell proliferation and anti-inflammatory responses	Inhibits viral replication	[145, 151]
	SRSF3	Pre-mRNA splicing factor which promotes exon-inclusion during alternative splicing	unknown	[145]
	YWHAB	Adaptor protein which facilitates cell proliferation	Prevents PCV2-induced autophagy and apoptosis	[145, 152]
	ZNF511	Transcription factor interacting with RNA polymerase II	unknown	[145]