# JAST (Journal of Animal Science and Technology) TITLE PAGE Upload this completed form to website with submission

ARTICLE INFORMATION	Fill in information in each box below
Article Type	Research article
Article Title (within 20 words without abbreviations)	Effects of dietary xylanase supplementation on growth performance and gut health of weaned pigs fed a high non-starch polysaccharides diet
Running Title (within 10 words)	Dietary xylanase in a high non-starch polysaccharides diet
Author	Jinmu Ahn1#, Jin Ho Cho2#, Hyeun Bum Kim3#, Jinuk Nam1, Yonggu Kang1, Ikcheol Shin1, Yoon Tack Jang4, Min Ah Park4, Yujin Kim4, Seungeun Lee4, Jun-ok Mun4, Minho Song1*, Hyunjin Kyoung1*
	<ul> <li># These authors contributed equally to this work as the first author</li> <li>* Corresponding authors</li> </ul>
Affiliation	<ol> <li>Division of Animal and Dairy Science, Chungnam National University, Daejeon 34134, Republic of Korea</li> <li>Department of Animal Science, Chungbuk National University, Cheongju 28644, Republic of Korea</li> <li>Department of Animal Resources Science, Dankook University, Cheonan 31116, Republic of Korea</li> <li>Application Center, CJ Blossom Park, Suwon 16495, Republic of Korea</li> </ol>
ORCID (for more information, please visit https://orcid.org)	Jinmu Ahn (https://orcid.org/0009-0005-1490-2974) Jin Ho Cho (https://orcid.org/0000-0001-7151-0778) Hyeun Bum Kim (https://orcid.org/0009-0003-1366-6090) Jinuk Nam (https://orcid.org/0009-0003-8696-5697) Yonggu Kang (https://orcid.org/0009-0008-3593-2656) Ikcheol Shin (https://orcid.org/0009-0003-0072-5626) Yoon Tack Jang (https://orcid.org/0000-0002-3296-4491) Min Ah Park (https://orcid.org/0000-0002-4993-6374) Yujin Kim (https://orcid.org/0009-0006-9632-641X) Seungeun Lee (https://orcid.org/0009-0001-7149-9270) Jun-ok Mun(https://orcid.org/0000-0002-2118-2833) Minho Song (https://orcid.org/0000-0002-4515-5212) Hyunjin Kyoung (https://orcid.org/0000-0001-5742-5374) No potential conflict of interest relevant to this article was reported.
<b>Funding sources</b> State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available.	This research was supported by the High Value-Added Food Technology Development Program of the Korean Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry, and Fisheries (IPET), the Ministry for Food, Agriculture, Forestry and Fisheries of the Republic of Korea (321037-05-3- HD030) and the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (RS-2023-00271355).
Acknowledgements	We acknowledge supports from Application Center, CJ Blossom Park, Suwon 16495, Republic of Korea.
Availability of data and material	Upon reasonable request, the datasets of this study can be available from the corresponding author.
<b>Authors' contributions</b> Please specify the authors' role using this form.	Conceptualization: Ahn J, Cho JH, Kim HB, Song M, Kyoung H. Data curation: Ahn J, Cho JH, Kim HB, Song M, Kyoung H. Formal analysis: Ahn J, Nam J, Kang Y, Shin I. Methodology: Ahn J, Song M, Kyoung H. Software: Ahn J, Kyoung H. Validation: Cho J, Kim HB, Song M, Kyoung H. Investigation: Jang YT, Park MA, Kim Y, Lee S, Mun J.

	Writing - original draft: Ahn J, Cho JH, Kim HB. Writing - review & editing: Ahn J, Cho J, Kim HB, Nam J, Kang Y, Shin I, Jang YT, Park MA, Kim Y, Lee S, Mun J, Song M, Kyoung H.
Ethics approval and consent to participate	The experimental protocol for this study was reviewed and approved by the Institutional Animal Care and Use Committee of Chungnam National University, Daejeon, Republic of Korea (approval #: 202112-CNU-182)

## CORRESPONDING AUTHOR CONTACT INFORMATION

For the corresponding author (responsible for correspondence, proofreading, and reprints)	Fill in information in each box below
First name, middle initial, last name	Minho Song Hyunjin Kyoung
Email address – this is where your proofs will be sent	mhsong@cnu.ac.kr hjkyoung@cnu.ac.kr
Secondary Email address	
Address	Division of Animal and Dairy Science, Chungnam National University, Daejeon 34134, Republic of Korea Division of Animal and Dairy Science, Chungnam National University, Daejeon 34134, Republic of Korea
Cell phone number	
Office phone number	+82-42-821-5776 +82-42-821-7857
Fax number	

#### 8 Abstract

9 Non-starch polysaccharides (NSPs) in cereal grains can negatively affect the growth performance of early weaned 10 pigs. Weaned pigs cannot digest NSPs due to a lack of endogenous enzymes. Feeds containing high levels of NSPs 11 can decrease nutrient digestibility by increasing digesta viscosity and modulating the gut environment. Dietary 12 xylanase (XYL) is used to increase nutrient utilization by degrading NSPs containing anti-nutritional factors. 13 Therefore, this study was conducted to evaluate the effects of XYL on the high NSPs diet on growth performance, 14 frequency of diarrhea, blood profiles, systemic immune responses, digesta characteristics, nutrient digestibility, 15 and intestinal health parameters of weaned pigs. XYL improved (p < 0.05) the average daily gain (ADG) and gain 16 to feed ratio from day 1 to 7 and day 1 to 14 compared with the control group (CON). Additionally, pigs fed XYL 17 tended to have a higher (p = 0.098) ADG from day 8 to 14 than those fed CON. Pigs fed XYL tended to have a 18 lower (p = 0.093) number of white blood cells on day 28 than those fed CON. The XYL group tended to increase 19 (p = 0.088) digesta pH in the duodenum, but decreased digesta pH in the jejunum (p = 0.069) and cecum (p < 0.069)20 0.05) on day 28 compared with the CON. Pigs fed XYL had higher (p < 0.05) apparent total tract digestibility 21 (ATTD) and apparent ileal digestibility of dry matter on day 28 than those fed CON. Additionally, the XYL group 22 tended to improve ATTD of energy (p = 0.083) and crude protein (p = 0.082) compared with the CON. Dietary 23 XYL decreased concentrations of tumor necrosis factor- $\alpha$  (p < 0.05), immunoglobulin G (p = 0.066), 24 malondialdehyde (p = 0.070) in jejunal mucosa compared with CON. In conclusion, supplementation of high 25 NSPs diet with XYL enhanced the growth performance of weaned pigs by enhancing nutrient digestibility through 26 the modulation of the intestinal environment.

27 Keywords: Non-starch polysaccharides, Nutrients digestibility, Xylanase, Gut health, Weaned pigs.

# Introduction

29 Among the various factors affecting the swine industry, such as feed, disease, management, and 30 environmental conditions, feed cost is regarded as the largest expense [1]. Additionally, cereal grain costs have 31 continued to increase over the past 50 years. Moreover, the quantity of cereal grain available in swine diets is 32 limited due to increasing consumer demand for meat following population growth and rising industrial demand, 33 such as bioethanol and biodiesel production [2,3]. Therefore, swine nutritionists have been seeking substitutes for 34 conventional feed ingredients to reduce feed costs. Grain co-products such as corn distiller's dried grains with 35 solubles (DDGS), wheat bran, and wheat middlings can be used as low-cost alternative ingredients [4,5]. However, 36 these co-products have high concentrations of non-starch polysaccharides (NSPs) [6,7]. NSPs in feed encapsulate 37 other nutrients that are hydrolyzed by endogenous enzymes in the intestine, thereby hindering their utilization 38 [6,8].

39 Early weaned pigs are exposed to environmental, nutritional, physiological, and immunological changes 40 during this period [9-11]. In particular, weaned pigs consume plant-based feedstuffs containing anti-nutrient 41 factors, such as NSPs [12]. However, pigs have an immature digestive tract, which results in poor nutrient 42 utilization [13]. NSPs are primarily composed of cellulose, hemicellulose, and pectin. They constitute a notable 43 portion of the plant cell wall and more than 90% of its structural strength [14]. NSPs are classified as soluble 44 and insoluble NSPs based on their physicochemical properties. Excessive NSP inclusion in feed leads to negative 45 effects on the gastrointestinal tract and causes morphological changes [15]. Soluble NSPs have a high water-46 holding capacity and can increase the bulk and viscosity of the digesta, affecting the passage rate. In contrast, 47 insoluble NSPs suppress the enzymatic activity of nutrients in the digesta, thereby delaying transit time and 48 disrupting intestinal motility [10,12]. These functions reduce nutrient absorption and lead to intestinal disorders 49 due to local inflammation, which leads to diarrhea and pathogenic infections [16,17]. Xylanase (XYL) enzyme 50 degrade NSPs such as xylan and arabinoxylan structures in cereal grains [12], which can improve growth 51 performance and nutrient digestibility. Additionally, XYL can release metabolites from indigest nutrients through 52 hydrolysis action, which not only modulates the gut environment and body immune status [10,18], but also the 53 released metabolites can be utilized for energy utilization in the body [19]. However, the effects of dietary XYL 54 on growth and health of pigs vary depending on the dietary grain composition [10,12,20]. Moreover, 55 supplementation of dietary XYL in high level NSPs diets can be expected to have positive effects in pigs [21]. 56 Therefore, the objective of this study was to evaluate the effect of XYL supplementation on growth performance,

57 frequency of diarrhea, blood profiles, systemic immune responses, digesta characteristics, nutrients digestibility,58 and intestinal health parameters of weaned pigs fed high NSPs diets.

59

60

# Materials and methods

### **61** Experimental design, animals, and diets

62 The protocols for this animal study were approved by the Institutional Animal Care and Use Committee
63 of Chungnam National University, Daejeon, Republic of Korea and followed the guidelines and regulations for
64 animal use (approval# 202112-CNU-182).

65 A total of 60 newly weaned pigs [(Landrace  $\times$  Yorkshire)  $\times$  Duroc; 8.04  $\pm$  0.99 kg of average initial body weight (BW); 4 weeks of age] were assigned to 2 dietary treatments (5 pigs per pen; 6 replicate pens per 66 67 dietary treatment) using a randomized complete block design (block = initial BW). Control (CON) pigs were fed 68 a high NSPs diet, and the other pigs were fed a CON diet with 0.03% XYL for 28 days. The experimental diet 69 was formulated to meet or exceed the nutrient requirements of the weaned pigs, as estimated by the National Research Council [22] (Table 1). The experimental diet was designed with high NSPs content using corn DDGS 70 71 and wheat. XYL product was obtained from a commercial company (CJ Blossom Park, Suwon, Republic of 72 Korea). During the last week of the study, chromium oxide (Daejung Chemicals & Metals Co. Ltd., Siheung-si, 73 Gyeonggi-do, Republic of Korea) was added at a concentration of 3 g/kg to the dietary treatments as an indicator 74 of nutrient digestibility [23]. All pigs had ad libitum access to feed and water and were housed in pens of equal 75 size  $(2 \text{ m} \times 2 \text{ m})$  with automatically controlled temperature, humidity, and lighting during the experimental period.

76

#### 77 Data and sample collection

Feed intake and pigs' BW in each pen were recorded on day 1, 7, 14, and 28 to calculate growth performance parameters, including average daily gain (ADG), average daily feed intake (ADFI), and gain to feed ratio (G:F). The frequency of diarrhea in pigs was recorded during the first 2 weeks after weaning by visual observation with a score ranging from 1 to 5 (1 = dry feces, 2 = normal feces, 3 = slightly mild feces, 4 = mild diarrhea, and 5 = watery severe diarrhea) and was calculated by counting the number of days with a pen average

83 diarrhea score of 4 or higher [24]. Blood samples were collected from one randomly selected pig in each pen on 84 days 1, 7, 14, and 28 using a 10 mL vacutainer tube with or without ethylenediaminetetraacetic acid (EDTA) to 85 yield whole blood and serum, respectively [25]. Serum samples were obtained after centrifugation of non-EDTA tubes at  $3,000 \times \text{g}$  for 15 min at 4 °C. The supernatant after centrifugation was collected and stored at -80°C until 86 87 analysis of immune responses [26]. Fecal samples were collected from one randomly selected pig per pen by rectal 88 stimulation for 3 days following a four day adaptation period and stored at -20 °C for subsequent analysis of 89 apparent total tract digestibility (ATTD) [12]. On the last day of the study, two randomly selected pigs per pen 90 were anesthetized with 2 mL of suxamethonium chloride (Succicholine; Ilsung Pharm. Co. Ltd., Seoul, Republic 91 of Korea). Immediately after injection, the pigs were euthanized by exposure to  $CO_2$  gas. Ileal digesta samples 92 were collected into 50 mL tubes from a site 30 cm proximal to the ileocecal junction and stored at -20 °C until 93 apparent ileal digestibility (AID) analysis [27]. To measure the viscosity and pH of digesta, samples were collected 94 from the stomach, duodenum, jejunum, ileum, and cecum. The samples for viscosity analysis were placed into 50 95 mL tubes and stored at -20 °C, while digesta pH was measured immediately after collection using a digital pH 96 meter (Accumet; Fisher Scientific, Hampton, NH, USA) [26]. Mid-jejunal segments were collected, rinsed with 97 distilled water, and fixed in 50 mL conical tubes with 10% neutral buffered formalin solution for 98 histomorphological measurements. Mucosal samples were scraped from the remaining mid-jejunum, placed into 99 2 mL microtubes, and stored at -80 °C freezer to determine the mucosal immune responses and oxidative stress 100 indicators.

101

#### **102** Blood profiles and systemic immune responses

103 Whole blood samples in EDTA tubes were analyzed by an automated hematology analyzer (scil Vet abc 104 hematology analyzer, scil animal care company, F-67120 Altorf, France) for evaluating the number of white blood 105 cells (WBC), platelet, red blood cells, mean corpuscular volume, hemoglobin, mean corpuscular, hematocrit, and 106 mean corpuscular concentration. Serum samples were used to measure tumor necrosis factor- alpha (TNF-a; R&D 107 System Inc., Minneapolis, MN, USA), transforming growth factor- beta1 (R&D System Inc., Minneapolis, MN, 108 USA), immunoglobulin G (IgG; Bethyl Laboratories Inc., Waltham, MA, USA), IgA (Bethyl Laboratories Inc., 109 Waltham, MA, USA), interleukin-10 (IL-10; R&D System Inc., Minneapolis, MN, USA), and C-reactive protein 110 (CRP; Aviva Systems Biology Inc., San Diego, CA, USA) using porcine-specific enzyme-linked immunosorbent 111 assay (ELISA) kits following the provided manufacturer protocols. Absorbance was measured at 450 nm using a

- microplate reader (Epoch microplate spectrophotometer, BioTek Instruments Inc., Winooski, VT, USA) and
  software (Gen5 Data Analysis Software, BioTek Instruments Inc., Winooski, VT, USA), and the concentrations
- **114** were calculated based on the standard curve from each ELISA kit.
- 115

#### 116 Viscosity of digesta

117 The procedure for determining the digesta viscosity was adapted from a previous study [15]. The 118 viscosity of the digesta in the stomach, jejunum, ileum, and cecum was measured using a viscometer (Model DV-119 II Version 2.0; Brookfield Engineering Laboratories Inc., Stoughton, MA, USA). The stored sample tubes were 120 centrifuged at  $1,000 \times g$  for 10 min to obtain the liquid phase. After centrifugation, the liquid phase was transferred 121 to a 2 mL microtube to second centrifuge at  $1,000 \times g$  for 10 min. The supernatant was obtained and transferred 122 to a 2 mL microtube for further analysis. Before measuring 0.5 mL of digesta supernatant, the viscometer was set 123 to 25°C. The viscometer results were calculated as the average between 45.0/s and 22.5/s shear rates and recorded 124 as millipascal-seconds.

125

#### **126** Nutrient digestibility

Diets, ileal digesta, and fecal samples were dried using air-forced drying oven at 65 °C for 72 h. All 127 128 samples were ground to powder using a grinder (80350, Hamiltonbeach Inc, Virginia, USA) for AID and ATTD 129 analysis. The bomb calorimeter (Parr 1261EA Bomb Calorimeter, Parr Instrument CO, Moline, IL, USA) was 130 used for measuring energy [28]. Dry matter (DM; method 930.15), crude protein method (CP; method 988.05), 131 and crude fiber (CF; method 962.09) were analyzed based on Association of Official Analytical Chemists [29]. 132 NSPs were analyzed based on a previous report [30]. The chromium concentration was analyzed using an 133 absorption spectrophotometer (Hitachi Z-5000 Absorption Spectrophotometer, Hitachi High-Technologies Co, 134 Tokyo, Japan). The calculation methods for AID and ATTD have been described in a previous study [27,31].

135

#### **136** Histomorphological analysis

Histomorphological analyses were performed as previously described [12,32]. The two sections from
the mid-jejunum were dehydrated, embedded in paraffin wax, sectioned to 5 μm and stained using hematoxylin
and eosin. The villus height to crypt depth ratio (VH:CD) was measured from stained slides using an Olympus

140 CX31 microscope (Lumenera Corporation, Ottawa, Canada) equipped with an Infinity 2-2 digital CCD camera. 141 Fifteen well-oriented, intact villi and their associated crypt depths were measured on each slide. The length was 142 measured from the top of the villi to the villus crypt junction, and the crypt depth was measured from the villus 143 crypt junction to the bottom of the crypt. The images for counting Ki-67 positive cells in the crypt were cropped 144 into 15 intact images from each slide, and the ImageJS software was used to calculate the percentage of Ki-67 145 positive cells to total cells in the crypt.

146

147 Intestinal immune responses and oxidative stress indicators

Mucosal samples were weighed, suspended in 1 mL of phosphate-buffered saline, homogenized on ice 148 149 using a tissue homogenizer (Tissuemiser; Thermo Fisher Scientific Inc., Waltham, MA USA), and centrifuged at 150 14,000 × g at 4 °C for 3 min. After centrifugation, the supernatant was collected and stored at -80 °C until further 151 analysis [10]. Jejunal mucosal immune responses [TNF-a (R&D System Inc., Minneapolis, MN, USA), IgG 152 (Bethyl Laboratories Inc., Waltham, MA, USA), IgA (Bethyl Laboratories Inc., Waltham, MA, USA), IL-6 (R&D 153 System Inc., Minneapolis, MN, USA), and IL-8 (R&D System Inc., Minneapolis, MN, USA)] and oxidative stress [malondialdehyde (MDA; Cell Biolabs, San Diego, CA, USA), protein carbonyl (PC; Cell Biolabs, San Diego, 154 155 CA, USA), and endotoxin (Aviva Systems Biology Inc., San Diego, CA, USA)] were determined using ELISA 156 kits following the manufacturer's instruction. The absorbance was measured using a microplate reader (Epoch 157 microplate spectrophotometer, BioTek Instruments Inc., Winooski, VT, USA) and software (Gen5 Data Analysis 158 Software, BioTek Instruments Inc., Winooski, VT, USA). All concentrations were calculated based on standard 159 curve generated from the concentration and absorbance of each standard.

160

#### **161** Statistical analyses

All data, except for the frequency of diarrhea, were analyzed using the GLM procedure in SAS (SAS
Inst. Inc., Cary, NC, USA), using a randomized complete block design (block = initial BW). The experimental
unit used was a pen. Statistical models for growth performance, digesta characteristics, nutrient digestibility,
jejunal health parameters, blood profiles, and systemic immune responses included effects of dietary treatments
as the main effects and BW as a covariate. Chi-square test was used to determine the frequency of diarrhea.

167	Statistical significance and tendency between dietary treatments were considered at $p < 0.05$ and $0.05 \le p < 0.10$ ,
168	respectively.
169	
170	Results
171	Growth performance and frequency of diarrhea
172	XYL increased ( $p < 0.05$ ) ADG and G:F from day 1 to 7 and from day 1 to 14 compared with the CON
173	(Table 2). Additionally, pigs fed XYL tended to have a higher ( $p = 0.098$ ) ADG from day 8 to 14 than those fed
174	CON. However, there was no difference on ADFI of weaned pigs during overall experimental period between the
175	treatments. There was no difference in frequency of diarrhea between CON and XYL.
176	
177	Blood profiles and systemic immune responses
178	Pigs fed XYL tended to have lower ( $p = 0.093$ ) WBC counts on day 28 than those fed the CON (Table
179	3). However, no differences were observed in serum immune responses of weaned pigs between the CON and
180	XYL (Table 4).
181	
182	Digesta characteristics and nutrient digestibility
183	Pigs fed XYL tended to have a higher ( $p = 0.088$ ) digesta pH in the duodenum than those fed the CON
184	(Table 5). In contrast, the XYL group digesta pH decreased in the jejunum ( $p = 0.069$ ) and cecum ( $p < 0.05$ ).
185	However, no differences were observed in the digesta viscosity of the stomach, duodenum, jejunum, ileum, and
186	cecum between the dietary treatments. Dietary XYL increased the AID and ATTD of DM ( $p < 0.05$ ) and ATTD of
187	energy ( $p = 0.083$ ) and CP ( $p = 0.082$ ) on day 28 compared with the CON (Table 6).
188	
189	Intestinal histomorphology, immune responses, and oxidative stress indicators
190	No differences were found in VH:CD and percentage of Ki-67 positive cells in the jejunum between the
191	dietary treatments (Table 7). However, pigs fed XYL had lower ( $p < 0.05$ ) concentrations of TNF- $\alpha$ and IgG in

**192** the jejunal mucosa than those fed CON. In addition, dietary XYL tended to decrease (p = 0.070) MDA level in **193** the jejunal mucosa compared with CON.

194

195

# Discussion

196 In this study, we demonstrated that supplemental dietary XYL in weaned pigs fed a high NSPs diet 197 improved growth performance and nutrient digestibility, modulated digesta pH in the gut, and reduced local 198 immune responses and oxidative stress indicator. These positive effects of dietary XYL may be attributed to the 199 increased nutrient utilization efficiency and the regulation of the gut environment through the enzymatic breakdown of NSPs in the grain cell walls.

The high NSPs diets used in this study included corn DDGS and wheat, which contained approximately 201 202 3.1 and 2.4% soluble NSPs and 25.2 and 9.0% insoluble NSPs, respectively [33-35]. The negative effects of 203 DDGS inclusion in swine diets on growth performance and nutrient digestibility reported in previous studies could 204 be explained by the increased NSPs levels in the feed, which reduce nutrients availability and gut functions 205 [17,35,36]. Additionally, wheat is a relatively viscous grain compared to corn, and wheat-based diets have a higher 206 total NSP content than corn-based diets [13,37]. NSPs are not degraded by the endogenous digestive enzymes of 207 pigs, which limit their nutrient utilization [38]. Dietary XYL, an exogenous enzyme, has been used to break down 208 the structural bonds of NSP, thereby modulating digesta characteristics and enhancing nutrient utilization in the 209 feed [7]. However, the effects of dietary XYL on the growth performance of pigs have been inconsistent [15,20]. 210 In the present study, the addition of dietary XYL improved ADG and G:F in pigs for the first 2 weeks after weaning. 211 The differences in the effects of XYL on growth performance may be attributed to the differences in NSPs content 212 in the diet, and the effects of XYL were relatively pronounced when supplemented with high NSP diet [18,20]. 213 The high NSP diet used in this study is considered sufficient as a substrate for enzymatic XYL activity, thereby 214 improving nutrient utilization and resulting in beneficial performance in weaned pigs. Thus, our results showed 215 that although supplemental XYL did not influence the frequency of diarrhea for the first 2 weeks after weaning, 216 the growth performance indicated a result indicates positive effect of dietary XYL during the critical period after 217 weaning.

218 The viscosity of digesta in the small intestine is related to the structure and molecular weight of the 219 polysaccharides, which can have a greater influence on viscosity than the type of linkage or sugar composition of 220 the polysaccharide [13]. The high digesta viscosity in the digestive tract can disturb nutrient digestibility and 221 absorption and can also increase the level of oxidative stress and inflammatory responses, which results in damage to the intestinal histomorphology [12]. Previous studies have demonstrated the functional effects of supplemental 222 223 XYL on the digestive tract of weaned pigs fed a high NSPs diet [10,12]. However, the effect of dietary XYL on 224 digesta viscosity can varies depending on the presence of primary cereal grains. Several studies have evaluated 225 the effects of supplemental XYL in corn-soybean meal or corn-soybean meal-corn DDGS-based diets on intestinal 226 digesta viscosity [10,20,39] and have also assessed the absence of interactions between dietary XYL and corn 227 DDGS [18]. In this study, we observed that the addition of dietary XYL to corn-soybean meal-corn DDGS-wheat-228 based diets containing high levels of NSPs did not affect the viscosity of gastrointestinal digesta in weaned pigs. 229 Cereal co-products generally do not elevate digesta viscosity to a similar extent as conventional cereal grains 230 because of their higher levels of insoluble NSPs [13,33]. Nevertheless, corn DDGS and wheat in the basal diets 231 of this study not only had higher levels of soluble NSPs than corn and soybean meal but also contained more 232 insoluble NSPs than soluble NSPs [34,40]. The lack of a substantial effect on digesta viscosity despite XYL 233 supplementation may be attributed to the high content of insoluble NSPs in the experimental diets, which may not 234 have been readily degraded by exogenous XYL. Previous studies have shown that the porcine digestive tract digestibility of insoluble NSP following XYL addition is lower than that of soluble NSP [41,42]. Unlike soluble 235 236 NSPs, which contribute to digesta viscosity, insoluble NSPs act as physical barriers to digestive enzymes in the 237 gastrointestinal tract [43], thereby impeding nutrient utilization. Therefore, high levels of NSPs in the diet may 238 provide sufficient substrates for XYL activity, but the impact on digesta viscosity is limited due to high levels of 239 insoluble NSPs in the diets.

We found that dietary XYL modulated intestinal digesta pH in weaned pigs. Soluble NSPs are primarily
degraded in the proximal intestine [44], and the observed increase in duodenal digesta pH following XYL addition
may be associated with the accelerated degradation of NSPs. This change enhances the hydrolysis of NSPs,
making nutrients more accessible to endogenous digestive enzymes. In contrast, the lower digesta pH in the
jejunum and cecum of XYL supplemented pigs may indicate the fermentation of NSPs and/or their degradation
products. In general, the water-holding capacity of fibers affects fermentability, soluble NSPs are more
fermentable than insoluble NSPs [39] supplemental XYL indirectly provides fermentable xylooligosaccharides

(XOS) via hydrolysis of the xylan backbone [45]. Fermentation of NSPs end-products, such as XOS, by
commensal gut microbiota leads to the production of short-chain fatty acids (SCFAs), which lower the pH of the
gut [46,47]. In addition, SCFAs produced by gut microbiota play crucial roles in gut health by modulating barrier
function and immune responses [47]. Therefore, it is suggested that the pH of the gut digesta modified by XYL
supplementation may contribute not only to improved accessibility to nutrients but also to changes in the gut
environment of weaned pigs.

253 Our findings on nutrient digestibility further support the beneficial effects of XYL supplementation. 254 Supplemental XYL did not affect NSP digestibility in weaned pigs but improved the AID of DM and ATTD of 255 DM, energy, and CP. The proposed mode of action of supplemental XYL to enhance nutrient digestibility is to 256 degrade the main chains of NSPs in the diet, thereby increasing the accessibility of nutrients to endogenous 257 digestive enzymes [48]. This hydrolysis converts the digesta compounds into smaller molecules that can be 258 absorbed more efficiently into the intestine. However, the improvement in nutrient digestibility did not support 259 the improvement in growth performance, which would require consideration of nutrient absorption. Furthermore, 260 further studies are needed to evaluate the dosage of diePtary XYL at different growth stages to effectively feed 261 animals for the long-term translation of growth enhancement through nutrients in feed. Although improvements in nutrient digestibility did not lead to improved growth performance of weaned pigs, findings of this study 262 263 suggests that the addition of XYL to high NSPs diets can enhance nutrient availability rather than directly 264 improving NSP digestion by mitigating the NSP-induced physical barrier to digestion. Additionally, the marked 265 improvement in ATTD, relative to AID, suggests that the large intestinal microbiota may contribute to nutrient 266 digestibility through fermentation, as supported by the digesta pH in the hindgut. Further studies are needed to 267 investigate the effect of dietary XYL on different diet compositions and its effects on the proximal to distal gut 268 may provide its potential to enhance growth by improving nutrient utilization in weaned pigs.

The inclusion of DDGS in the feed can cause oxidative stress in pigs, which may modulate their immune responses [49,50]. In addition, highly viscous soluble NSP can increase oxidative stress and inflammatory responses associated with intestinal enterotoxigenic *Escherichia coli* proliferation [51–53], and a positive correlation with post-weaning colibacillosis has been reported [54,55]. One of the final products of lipid and protein peroxidation are considered oxidative stress products such as MDA and PC [10], which can destroy the intestinal environment. Excessive levels of reactive oxygen species can damage cellular components such as the cell membrane, DNA, and proteins, leading to chronic inflammation [56]. Thus, oxidative stress can induce pro276 inflammatory cytokines in intestinal immune cells and mucosal epithelial cells to modulate immune responses in 277 the intestine [10,57]. In the present study, mucosal TNF-α, IgG, and MDA were reduced following dietary XYL 278 addition, indicating that XYL appears to have anti-inflammatory and anti-oxidant effects in weaned pigs. In 279 addition, the decrease in jejunal mucosal IgG levels following XYL supplementation suggests that the local 280 immune response was modulated. As high levels of NSP in feed can act as anti-nutritional factors [58], effective 281 NSPs degradation and fermentation by dietary XYL may be a result of a reduction in the antigenic components 282 reaching the intestinal immune system. Furthermore, the reduced WBC count in XYL-supplemented pigs suggests 283 a potential systemic immunomodulatory effect, because the number of WBC can be used as an indicator of 284 systemic inflammatory responses. However, as no effects were found on the levels of systemic inflammatory 285 cytokines, the impact of dietary XYL on systemic immune responses requires further investigation.

286

287

# Conclusion

288 Our findings suggest that supplementation of high NSP diet with dietary XYL including corn DDGS 289 and wheat, can improve the early growth performance and nutrient digestibility of weaned pigs. Dietary xylanase 290 may play an important role in enhancing gut health by degrading and fermenting complex NSPs, thereby 291 regulating nutrient utilization and the biochemical environment in the gut. In addition, the reduction in 292 inflammatory markers and oxidative stress indicators suggests that dietary xylanase helps maintain intestinal 293 integrity and functions, demonstrating its potential as a feed additive to improve the growth and health of weaned 294 pigs. Further studies are needed to evaluate the effects of dietary xylanase on the growth performance, intestinal 295 health, and local and systemic immune responses in weaned pigs fed in various cereal grain-based diets.

296

297

# Funding

298 This research was supported by the High Value-Added Food Technology Development Program of the
299 Korean Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry, and Fisheries (IPET),
300 the Ministry for Food, Agriculture, Forestry and Fisheries of the Republic of Korea (321037-05-3-HD030) and
301 the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the
302 Ministry of Education (RS-2023-00271355).

303	
304	Acknowledgements
305	We acknowledge supports from CJ CheilJedang, Seoul, Republic of Korea

# References

- Maniraguha V, Hong JS, Yu M, Oketch EO, Yi YJ, Yun H, et al. Feeding dietary non-starch polysaccharides supplemented with xylanase could improve the performance of broilers. J Anim Sci Technol. 2023;66:999.
   https://doi.org/10.5187/jast.2023.e114
- 310 2. Lindberg JE. Fiber effects in nutrition and gut health in pigs. J Anim Sci Biotechnol. 2014;5:1–7.
  311 https://doi.org/10.1186/2049-1891-5-15
- 312 3. Rosegrant MW. Biofuels and grain prices: Impacts and policy responses. Testimony for the US Senate Committee on Homeland Security and Governmental Affairs. Washington, DC, USA. 2008.
- 314 4. Zhang Z, Tun HM, Li R, Gonzalez BJM, Keenes HC, Nyachoti CM, et al. Impact of xylanases on gut microbiota of growing pigs fed corn- or wheat-based diets. Anim Nutr. 2018;4:339–50. https://doi.org/10.1016/j.aninu.2018.06.007
- 317 Ndou SP, Kiarie E, Agyekum AK, Heo JM, Romero LF, Arent S, et al. Comparative efficacy of xylanases on 5. 318 growth performance and digestibility in growing pigs fed wheat and wheat bran- or corn and corn DDGS-319 supplemented with phytase. Feed Sci Technol. 2015;209:230-9. based diets Anim 320 https://doi.org/10.1016/j.anifeedsci.2015.08.011
- Betry AL, Patience JF. Xylanase supplementation in corn-based swine diets: A review with emphasis on potential mechanisms of action. J Anim Sci. 2020;98:skaa318. https://doi.org/10.1093/jas/skaa318
- 323 7. Zijlstra RT, Owusu-Asiedu A, Simmins PH. Future of NSP-degrading enzymes to improve nutrient utilization
   324 of co-products and gut health in pigs. Livest Sci. 2010;134:255–7.
   325 https://doi.org/10.1016/j.livsci.2010.07.017
- Nortey TN, Patience JF, Sands JS, Zijlstra RT. Xylanase supplementation improves energy digestibility of wheat by-products in grower pigs. Livest Sci. 2007;109:96–9. https://doi.org/10.1016/j.livsci.2007.01.092
- 328 9. Lu H, Yan H, Masey O'Neill HM, Bradley C, Bedford MR, Wilcock P, et al. Effect of timing of postweaning xylanase supplementation on growth performance, nutrient digestibility, and fecal microbial composition in weanling pigs. Can J Anim Sci. 2020;100:27–36. https://doi.org/10.1139/cjas-2019-0021
- 10. Duarte ME, Zhou FX, Dutra WM, Kim SW. Dietary supplementation of xylanase and protease on growth performance, digesta viscosity, nutrient digestibility, immune and oxidative stress status, and gut health of newly weaned pigs. Anim Nutr. 2019;5:351–8. https://doi.org/10.1016/j.aninu.2019.04.005
- 11. Lee J, Oh S, Kim M. Response to environmental enrichment of weanling pigs on growth, behaviour and welfare after weaning. J Anim Sci Technol. 2024;66:1246. https://doi.org/10.5187/jast.2023.e128
- 336 12. Moita VHC, Duarte ME, Kim SW. Functional roles of xylanase enhancing intestinal health and growth performance of nursery pigs by reducing the digesta viscosity and modulating the mucosa-Associated microbiota in the jejunum. J Anim Sci. 2022;100:skac116. https://doi.org/10.1093/jas/skac116

- Baker JT, Duarte ME, Holanda DM, Kim SW. Friend or foe? Impacts of dietary xylans, xylooligosaccharides, and xylanases on intestinal health and growth performance of monogastric animals. Animals. 2021;11:609. https://doi.org/10.3390/ani11030609
- 342 14. Kerr BJ, Shurson GC. Strategies to improve fiber utilization in swine. J Anim Sci Biotechnol. 2013;4:1–12.
   343 https://doi.org/10.1186/2049-1891-4-11
- 344 15. Baker JT, Duarte ME, Kim SW. Effects of dietary xylanase supplementation on growth performance, intestinal health, and immune response of nursery pigs fed diets with reduced metabolizable energy. J Anim Sci. 2024;102:skae026. https://doi.org/10.1093/jas/skae026
- 347 16. Urriola PE, Stein HH. Effects of distillers dried grains with solubles on amino acid, energy, and fiber digestibility and on hindgut fermentation of dietary fiber in a corn-soybean meal diet fed to growing pigs. J Anim Sci. 2010;88:1454–62. https://doi.org/10.2527/jas.2009-2162
- 350 17. Zheng L, Duarte ME, Sevarolli Loftus A, Kim SW. Intestinal health of pigs upon weaning: challenges and nutritional intervention. Front Vet Sci. 2021;8:628258. https://doi.org/10.3389/fvets.2021.628258
- 18. Chen H, Zhang S, Kim SW. Effects of supplemental xylanase on health of the small intestine in nursery pigs fed diets with corn distillers' dried grains with solubles. J Anim Sci. 2020;98:skaa185.
  https://doi.org/10.1093/jas/skaa185
- 355 19. Cheng YC, Kim SW. Use of microorganisms as nutritional and functional feedstuffs for nursery pigs and broilers. Animals. 2022;12:3141. https://doi.org/10.3390/ani12223141
- 357 20. He X, Yu B, He J, Huang Z, Mao X, Zheng P, et al. Effects of xylanase on growth performance, nutrients digestibility and intestinal health in weaned piglets. Livest Sci. 2020;233:103940.
  359 https://doi.org/10.1016/j.livsci.2020.103940
- 360 21. Adeola O, Cowieson AJ. Board-invited review: Opportunities and challenges in using exogenous enzymes
   361 to improve nonruminant animal production. J Anim Sci. 2011;89:3189–218. https://doi.org/10.2527/jas.2010-3715
- **363** 22. NRC. Nutrient Requirements of Swine: Eleventh Revised Edition. The National Academies Press. 2012.
- Park S, Lee J, Yun W, Hong S, Oh H, Song D, et al. Supplementation of protease and different nutrient density diets in growing-finishing pigs. J Anim Sci Technol. 2024;62:326. https://doi.org/10.5187/jast.2023.e55
- 366 24. Kang J, Cho J, Jang K, Kim J, Kim S, Mun D, et al. Citation: Effects of dietary enzyme cocktail on diarrhea and immune responses of weaned pigs Effects of dietary enzyme cocktail on diarrhea and immune responses of weaned pigs. Korean J Agric Sci. 2017;44:525–30. https://doi.org/10.7744/kjoas.20170058
- 369 25. Cho HA, Song MH, Lee J hwan, Oh HJ, Chang SY, An JW, et al. Effects of phytogenic feed additives in growing and finishing pigs under different stocking density. J Anim Sci Technol. 2024;66:981.
  371 https://doi.org/10.5187/jast.2023.e124

- 372 26. Duarte ME, Sparks C, Kim SW. Modulation of jejunal mucosa-associated microbiota in relation to intestinal health and nutrient digestibility in pigs by supplementation of β-glucanase to corn-soybean meal-based diets with xylanase. J Anim Sci. 2021;99:skab190. https://doi.org/10.1093/jas/skab190
- 375 27. Saha DC, Gilbreath RL. A modified chromic oxide indicator ratio technique for accvrate determin ation of nutrient digestibility. 1993;73:1001–4. https://doi.org/10.4141/cjas93-104
- 28. Park S, Choe J, Cho JH, Jang KB, Kyoung H, Park K Il, et al. Determination of optimal energy system and level for growing pigs. J Anim Sci Technol. 2024;66:514. https://doi.org/10.5187/jast.2023.e63
- **379** 29. AOAC International. Official Methods of Analysis. 17th ed. Arlington, MA: AOAC International. 2010.
- 380 30. Englyst HN, Quigley ME, Hudson GJ. Determination of dietary fibre as non-starch polysaccharides with gas–liquid chromatographic, high-performance liquid chromatographic or spectrophotometric measurement of constituent sugars. Analyst. 1994;119:1497–509. https://doi.org/10.1039/AN9941901497
- 383 31. Hossain MM, Hwang HS, Pang M, Choi MK, Kim IH. Effect of dietary Achyranthes japonica extract on growth performance of growing pigs and absorption rate of quercetin in blood. J Anim Sci Technol. 2024;66:103. https://doi.org/10.5187/jast.2023.e23
- 386 32. Kyoung H, Lee JJ, Cho JH, Choe J, Kang J, Lee H, et al. Dietary glutamic acid modulates immune responses and gut health of weaned pigs. Animals. 2021;11:504. https://doi.org/10.3390/ani11020504
- 388 33. Choct M. Feed non-starch polysaccharides for monogastric animals: Classification and function. Anim Prod
   389 Sci. 2015;55:1360–6. https://doi.org/10.1071/AN15276
- 34. Pedersen MB, Dalsgaard S, Knudsen KEB, Yu S, Lærke HN. Compositional profile and variation of distillers
   dried grains with solubles from various origins with focus on non-starch polysaccharides. Anim Feed Sci
   Technol. 2014;197:130–41. https://doi.org/10.1016/j.anifeedsci.2014.07.011
- 393 35. Jang JC, Zeng Z, Urriola PE, Shurson GC. Effects of feeding corn distillers dried grains with solubles diets
   394 without or with supplemental enzymes on growth performance of pigs: A meta-analysis. Transl Anim Sci. 2021;5:txab029. https://doi.org/10.1093/tas/txab029
- 396 36. Ayoade DI, Kiarie E, Slominski BA, Nyachoti CM. Growth and physiological responses of growing pigs to wheat-corn distillers dried grains with solubles. J Anim Physiol Anim Nutr. 2014;98:569–77.
  398 https://doi.org/10.1111/jpn.12117
- 37. Nguyen HT, Bedford MR, Wu SB, Morgan NK. Dietary soluble non-starch polysaccharide level influences performance, nutrient utilisation and disappearance of non-starch polysaccharides in broiler chickens. Animals. 2022;12:547. https://doi.org/10.3390/ani12050547
- 402 38. Chen Y, Shen D, Zhang L, Zhong R, Liu Z, Liu L, et al. Supplementation of non-starch polysaccharide enzymes cocktail in a corn-miscellaneous meal diet improves nutrient digestibility and reduces carbon dioxide emissions in finishing pigs. Animals. 2020;10:232. https://doi.org/10.3390/ani10020232

- 405 39. Passos AA, Cardoso Moita VH, Kim SW. Individual or combinational use of phytase, protease, and xylanase
  406 for the impacts on total tract digestibility of corn, soybean meal, and distillers dried grains with soluble fed
  407 to pigs. Anim Biosci. 2023;36:1869. https://doi.org/10.5713/ab.23.0212
- 408 40. Gutierrez NA, Serão NVL, Kerr BJ, Zijlstra RT, Patience JF. Relationships among dietary fiber components and the digestibility of energy, dietary fiber, and amino acids and energy content of nine corn coproducts fed to growing pigs. 2014;92:4505–17. https://doi.org/10.2527/jas.2013-7265
- 41. Abelilla JJ, Stein HH. Degradation of dietary fiber in the stomach, small intestine, and large intestine of growing pigs fed corn- or wheat-based diets without or with microbial xylanase. J. Anim. Sci. 2018;97:338–52. https://doi.org/10.1093/jas/sky403
- 414 42. Bartelt J, Jadamus A, Wiese F, Swiech E, Buraczewska L, Simon O. Apparent precaecal digestibility of nutrients and level of endogenous nitrogen in digesta of the small intestine of growing pigs as affected by various digesta viscosities. Arch Anim Nutr. 2002;56:93–107. https://doi.org/10.1080/00039420214182
- 43. Morgan N, Bhuiyan MM, Hopcroft R. Non-starch polysaccharide degradation in the gastrointestinal tract of broiler chickens fed commercial-type diets supplemented with either a single dose of xylanase, a double dose of xylanase, or a cocktail of non-starch polysaccharide-degrading enzymes. Poult Sci. 2022;101:101846. https://doi.org/10.1016/j.psj.2022.101846
- 42. Wellock IJ, Houdijk JGM, Kyriazakis I. Effect of dietary non-starch polysaccharide solubility and inclusion level on gut health and the risk of post weaning enteric disorders in newly weaned piglets. Livest Sci. 2007;108:186–9. https://doi.org/10.1016/j.livsci.2007.01.050
- 424 45. Sutton TA, O'Neill HVM, Bedford MR, McDermott K, Miller HM. Effect of xylanase and xylooligosaccharide supplementation on growth performance and faecal bacterial community composition in growing pigs. Anim Feed Sci Technol. 2021;274:114822. https://doi.org/10.1016/j.anifeedsci.2021.114822
- 427 46. Sun F, Li H, Sun Z, Liu L, Zhang X, Zhao J. Effect of arabinoxylan and xylo-oligosaccharide on growth performance and intestinal barrier function in weaned piglets. Animals. 2023;13:964. https://doi.org/10.3390/ani13060964
- 430 47. Akhtar M, Chen Y, Ma Z, Zhang X, Shi D, Khan JA, et al. Gut microbiota-derived short chain fatty acids are potential mediators in gut inflammation. Anim Nutr. 2022;8:350–60. https://doi.org/10.1016/j.aninu.2021.11.005
- 48. Singh AK, Mishra B, Bedford MR, Jha R. Effects of supplemental xylanase and xylooligosaccharides on production performance and gut health variables of broiler chickens. J Anim Sci Biotechnol. 2021;12:1–15. https://doi.org/10.1186/s40104-021-00617-8
- 436
  49. Weber TE, Kerr BJ. Effect of dietary distillers dried grains with solubles on indicators of oxidative stress and immune function in growing pigs. Livest Sci. 2011;142:85–91. https://doi.org/10.1016/j.livsci.2011.06.025
- 438 50. Li G, Wang X, Lin M, Lu Z, Yao W. Effects of corn DDGS in combination with compound enzymes on growth performance, carcass fat quality, and plasma and tissue redox homeostasis of growing-finishing pigs. Livest Sci. 2012;149:46–52. https://doi.org/10.1016/j.livsci.2012.06.019

- 441 51. Kim JC, Hansen CF, Mullan BP, Pluske JR. Nutrition and pathology of weaner pigs: Nutritional strategies to support barrier function in the gastrointestinal tract. Anim Feed Sci Technol. 2012;173:3–16. https://doi.org/10.1016/j.anifeedsci.2011.12.022
- 444 52. Tiwari UP, Chen H, Kim SW, Jha R. Supplemental effect of xylanase and mannanase on nutrient digestibility and gut health of nursery pigs studied using both in vivo and in vitro models. Anim Feed Sci Technol. 2018;245:77–90. https://doi.org/10.1016/j.anifeedsci.2018.07.002
- 53. Duarte ME, Tyus J, Kim SW. Synbiotic effects of enzyme and probiotics on intestinal health and growth of newly weaned pigs challenged with Enterotoxigenic F18+ Escherichia coli. Front Vet Sci. 2020;7:573. https://doi.org/10.3389/fvets.2020.00573
- 450 54. Wellock IJ, Fortomaris PD, Houdijk JGM, Wiseman J, Kyriazakis I. The consequences of non-starch polysaccharide solubility and inclusion level on the health and performance of weaned pigs challenged with enterotoxigenic Escherichia coli. Br J Nutr. 2008;99:520–30. https://doi.org/10.1017/S0007114507819167
- 453 55. Rhouma M, Fairbrother JM, Beaudry F, Letellier A. Post weaning diarrhea in pigs: Risk factors and non-colistin-based control strategies. Acta Vet Scand. 2017;59:1–19. https://doi.org/10.1186/s13028-017-0299-7
- 455 56. Hussain T, Tan B, Yin Y, Blachier F, Tossou MCB, Rahu N. Oxidative stress and inflammation: What polyphenols can do for us? Oxid Med Cell Longev. 2016:7432797. https://doi.org/10.1155/2016/7432797
- 457 57. Xue X, Falcon DM. The role of immune cells and cytokines in intestinal wound healing. Int J Mol Sci. 2019;20:6097. https://doi.org/10.3390/ijms20236097
- 459 58. Park S, Li W, St-Pierre B, Wang Q, Woyengo TA. Growth performance, nutrient digestibility, and fecal microbial composition of weaned pigs fed multi-enzyme supplemented diets. J Anim Sci. 2020;98:skaa306. https://doi.org/10.1093/jas/skaa306
- 462

464

**466** Table 1. Composition of experimental diets (as-fed basis)

Item	Control
Ingredient, %	
Corn	12.50
Corn distillers dried grains with soluble	30.00
Soybean meal, 44%	14.00
Wheat	30.00
Whey permeate	5.00
Spray dried porcine plasma	0.60
Fish meal	1.00
Soybean oil	3.30
Limestone	1.30
Dicalcium phosphate	0.75
Iodized salt	0.20
Vitamin-mineral premix <sup>1</sup>	0.35
Lysine-HCl	0.71
DL-Methionine	0.10
L-Threonine	0.17
L-Tryptophan	0.02
Total	100
Calculated energy and nutrient contents	
Dry matter, %	88.44
Metabolizable energy, kcal/kg	3398.69
Crude protein, %	21.93
SID <sup>2</sup> lysine, %	1.23
SID <sup>2</sup> methionine, %	0.40
SID <sup>2</sup> cysteine + methionine, %	0.69
SID <sup>2</sup> threonine, %	0.73
SID <sup>2</sup> tryptophan, %	0.20
Calcium, %	0.78
Total phosphorus, %	0.60
STTD <sup>3</sup> phosphorous, %	0.38
ATTD <sup>4</sup> phosphorous, %	0.33
Non-starch polysaccharides, %	15.47

 $<sup>\</sup>textbf{468} \qquad \text{K3, 3 mg; D-pantothenic acid, 15 mg; nicotinic acid, 40 mg; choline, 400 mg; and vitamin B_{12}, 12 \, \mu\text{g}; Fe, 90 \, \text{mg}}$ 

- 469 from iron sulfate; Cu, 8.8 mg from copper sulfate; Zn, 100 mg from zinc oxide; Mn, 54 mg from manganese oxide;
- **470** I, 0.35 mg from potassium iodide; Se, 0.30 mg from sodium selenite.
- 471 <sup>2</sup>SID, standardized ileal digestible.

- **472** <sup>3</sup>STTD, standardized total tract digestible.
- **473** <sup>4</sup>ATTD, apparent total tract digestible.

**474** Table 2. Effects of dietary xylanase on growth performance of weaned pigs<sup>1</sup>

Item <sup>2</sup>	CON	XYL	SEM	<i>p</i> -value
Day 1 to 7				-
Initial BW, kg	8.04	8.05	0.44	0.985
Final BW, kg	8.94	9.40	0.46	0.497
ADG, g/d	128.57	192.86	17.12	0.026
ADFI, g/d	225.52	245.90	12.84	0.288
G:F, g/g	0.574	0.785	0.056	0.014
Day 8 to 14				
Initial BW, kg	8.94	9.40	0.46	0.497
Final BW, kg	10.86	11.76	0.53	0.255
ADG, g/d	274.29	337.14	24.87	0.098
ADFI, g/d	397.19	434.67	30.71	0.408
G:F, g/g	0.688	0.776	0.064	0.306
Day 1 to 14				
Initial BW, kg	8.04	8.05	0.44	0.985
Final BW, kg	10.86	11.76	0.53	0.255
ADG, g/d	201.43	265.00	15.25	0.014
ADFI, g/d	311.36	340.29	19.96	0.330
G:F, g/g	0.647	0.779	0.045	0.043
Day 15 to 28				
Initial BW, kg	10.86	11.76	0.53	0.255
Final BW, kg	16.82	18.22	0.93	0.311
ADG, g/d	425.71	461.43	32.08	0.455
ADFI, g/d	801.29	857.00	47.65	0.428
G:F, g/g	0.532	0.539	0.020	0.644
Day 1 to 28				
Initial BW, kg	8.04	8.05	0.44	0.985
Final BW, kg	16.82	18.22	0.93	0.311
ADG, g/d	313.57	363.21	19.33	0.100
ADFI, g/d	556.32	598.64	31.80	0.369
G:F, g/g	0.564	0.607	0.020	0.105
Frequency of diarrhea, %	20.24	14.29	-	0.391

<sup>475</sup> 

477 ADG, average daily gain; ADFI, average daily feed intake; G:F, gain to feed ratio; Frequency of diarrhea for the

<sup>&</sup>lt;sup>1</sup>Each value is the mean value of 6 replicates (5 pigs/pen).

**<sup>476</sup>** <sup>2</sup>CON, high non-starch polysaccharides diet; XYL, CON + 0.03% dietary xylanase; BW, body weight;

**<sup>478</sup>** first two weeks after weaning = (number of diarrhea with score higher than 4 / number of pen days)  $\times 100$ .

Table 3. Effects of dietary xylanase on blood profiles of weaned pigs<sup>1</sup> 479

Item <sup>2</sup>	CON	XYL	SEM	<i>p</i> -value
White blood cell, ×10 <sup>3</sup> /µL				
Day 1	13.95	14.65	2.67	0.859
Day 7	19.70	18.50	1.53	0.592
Day 14	24.95	22.95	2.55	0.591
Day 28	25.45	21.42	1.54	0.093
Platelet, ×10 <sup>3</sup> / μL				
Day 1	426.25	480.00	48.54	0.463
Day 7	430.00	514.50	122.13	0.652
Day 14	719.00	637.33	108.32	0.591
Day 28	485.00	515.17	515.17	0.612
Hematocrit, %				
Day 1	32.15	31.18	1.81	0.716
Day 7	34.98	28.65	3.06	0.175
Day 14	29.33	28.60	2.08	0.808
Day 28	30.68	30.32	1.05	0.810
Red blood cell, ×10 <sup>6</sup> /µL				
Dav 1	5.65	5.53	0.30	0.783
Day 7	5.53	7.34	0.99	0.226
Day 14	4.68	5.63	0.46	0.178
Dav 28	6.11	5.95	0.23	0.627
MCV, fL				
Day 1	56.75	56.50	0.91	0.852
Day 7	49.67	50.50	2.64	0.828
Day 14	48.50	50.83	2.33	0.494
Day 28	50.33	51.00	1.29	0.722
Hemoglobin, g/dL				
Day 1	10.95	10.70	0.59	0.775
Day 7	9.65	12.78	1.98	0.289
Day 14	7.88	9.85	0.88	0.143
Day 28	10.50	10.35	0.35	0.770
MCH. pg		10.00	0.00	01110
Day 1	19.33	19.45	0.51	0.868
Day 7	16.95	17.17	0.82	0.855
Day 14	16.85	17.47	0.80	0.597
Day 28	17.27	17.38	0.47	0.864
Hematocrit, %		- /	,	
Day 1	32.15	31.18	1.81	0.716
Day 7	34.98	28.65	3.06	0.175
Day 14	29.33	28.60	2.08	0.808
Dav 28	30.68	30.32	1.05	0.810
MCHC, g/dL	20.00			
Day 1	34.03	34.50	0.52	0.544
Day 7	34.28	34.12	0.45	0.796
Dav 14	34.88	34.32	0.50	0.443
D = 20	24.20	24.12	0.20	0.772

480

Each value is the mean value of 6 replicates (1 pig/pen).

<sup>2</sup>CON, high non-starch polysaccharides diet; XYL, CON + 0.03% dietary xylanase; MCV, mean 481 482 corpuscular volume; MCH, mean corpuscular hemoglobin; MCHC, mean corpuscular hemoglobin concentration.

Item <sup>2</sup>	CON	XYL	SEM	<i>p</i> -value
Day 1				-
TNF-α, pg/mL	150.24	148.24	1.33	0.398
TGF-β1, pg/mL	2124.29	1888.61	100.22	0.172
IL-10, pg/mL	48.49	37.26	4.43	0.215
CRP, ng/mL	18.44	18.88	0.82	0.740
IgG, mg/mL	20.02	20.12	13.79	0.996
IgA, mg/mL	0.15	0.17	0.06	0.831
Day 28				
TNF-α, pg/mL	130.61	96.29	19.81	0.249
TGF-β1, pg/mL	2014.28	1840.31	167.33	0.479
IL-10, pg/mL	74.00	145.17	55.92	0.389
CRP, ng/mL	114.28	108.16	5.75	0.469
IgG, mg/mL	22.29	13.17	4.09	0.146
IgA, mg/mL	0.20	0.26	0.06	0.493

**483** Table 4. Effects of dietary xylanase on systemic immune responses of weaned pigs<sup>1</sup>

<sup>1</sup>Each value is the mean value of 6 replicates (1 pig/pen).

485 <sup>2</sup>CON, high non-starch polysaccharides diet; XYL, CON + 0.03% dietary xylanase; TNF- $\alpha$ , tumor

486 necrosis factor-α; IgG, immunoglobulin G; IgA, immunoglobulin A; IL-10, interleukin-10; CRP, C-reactive

**487** protein; TGF- $\beta$ 1, transforming growth factor- $\beta$ 1.

/

Item <sup>2</sup>	CON	XYL	SEM	<i>p</i> -value
Digesta pH				
Stomach	3.42	3.66	0.32	0.600
Duodenum	6.20	6.43	0.10	0.088
Jejunum	6.25	6.07	0.07	0.069
Ileum	7.00	7.15	0.13	0.419
Caecum	6.07	5.91	0.05	0.048
Digesta viscosity				
Stomach	1.15	1.05	0.05	0.183
Duodenum	1.16	1.23	0.05	0.278
Jejunum	1.17	1.12	0.04	0.304
Ileum	1.50	1.55	0.08	0.663
Caecum	1.88	1.88	0.15	0.985

**488** Table 5. Effects of dietary xylanase on pH and viscosity of gut digesta of weaned pigs<sup>1</sup>

489

<sup>1</sup>Each value is the mean value of 6 replicates (2 pigs/pen).

490

<sup>2</sup>CON, high non-starch polysaccharides diet; XYL, CON + 0.03% dietary xylanase.

/

**491** Table 6. Effects of dietary xylanase on nutrient digestibility of weaned pigs<sup>1</sup>

Item <sup>2</sup>	CON	XYL	SEM	<i>p</i> -value
Apparent ileal digestibility, %				
Dry matter	73.16	75.41	0.52	0.013
Energy	70.36	71.84	1.62	0.531
Crude protein	67.56	69.54	2.07	0.514
Crude fiber	40.79	41.59	2.95	0.852
Non-starch polysaccharides	38.38	43.09	2.17	0.156
Apparent total tract digestibility, %				
Dry matter	81.95	84.43	0.54	0.008
Energy	82.96	86.33	1.24	0.083
Crude protein	79.79	81.32	0.56	0.082
Crude fiber	51.32	52.52	5.30	0.876
Non-starch polysaccharides	51.69	52.05	3.12	0.936

<sup>492</sup> 

<sup>1</sup>Each value is the mean value of 6 replicates (1 pig/pen).

493

<sup>2</sup>CON, high non-starch polysaccharides diet, XYL, CON + 0.03% dietary xylanase.

/

**494** Table 7. Effects of dietary xylanase on intestinal health parameters of weaned pigs<sup>1</sup>

Item <sup>2</sup>	CON	XYL	SEM	<i>p</i> -value
Histomorphology				
VH:CD, μm/μm	3.45	3.47	0.22	0.952
Ki-67 positive, %	35.28	34.32	1.21	0.575
Mucosal immune responses				
TNF-α, pg/mg	0.56	0.14	0.05	< 0.001
IL-6, pg/mg	0.49	0.62	0.11	0.404
IL-8, ng/mg	0.58	0.57	0.09	0.927
IgG, μg/mg	7.02	4.41	0.97	0.066
IgA, μg/mg	1.51	2.54	0.45	0.110
Mucosal oxidative stress				
MDA, µM/mg	0.34	0.21	0.05	0.070
PC, nmol/mg	0.57	0.50	0.05	0.341
Endotoxin, EU/mL	94.90	60.28	17.50	0.167

495 <sup>1</sup>Each value is the mean value of 6 replicates (2 pigs/pen).

**496** <sup>2</sup>CON, high non-starch polysaccharides diet; XYL, CON + 0.03% dietary xylanase; VH:CD, villus

**497** height to crypt depth ratio; TNF-α, tumor necrosis factor-α; IgG, immunoglobulin G; IgA, immunoglobulin A;

498 IL-6, interleukin-6; IL-8, interleukin-8; MDA, malondialdehyde; PC, protein carbonyl.