

JAST (Journal of Animal Science and Technology) TITLE PAGE

Upload this completed form to website with submission

ARTICLE INFORMATION	Fill in information in each box below
Article Type	Research article
Article Title (within 20 words without abbreviations)	Estimation of standardized ileal digestible calcium and phosphorus requirements of broiler chickens from 10 to 21 days of age
Running Title (within 10 words)	Standardized ileal digestible calcium and phosphorus requirements of broilers
Author	Chae Won Lee ¹ , Changsu Kong ^{1,2,3*}
Affiliation	1 Research Institute for Innovative Animal Science, Kyungpook National University, Sangju 37224, Republic of Korea 2 Department of Animal Science, Kyungpook National University, Sangju 37224, Republic of Korea 3 Department of Animal Science and Biotechnology, Kyungpook National University, Sangju 37224, Republic of Korea
ORCID (for more information, please visit https://orcid.org)	Chae Won Lee (https://orcid.org/0000-0002-8281-5478) Changsu Kong (https://orcid.org/0000-0002-3876-6488)
Competing interests	No potential conflict of interest relevant to this article was reported.
Funding sources State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available.	This work was financially supported by Rural Development Administration, Korea (Project No. RS-2022-RD010370)
Acknowledgements	Not applicable.
Availability of data and material	Upon reasonable request, the datasets of this study can be available from the corresponding author.
Authors' contributions Please specify the authors' role using this form.	Conceptualization: Kong C. Data curation: Kong C. Formal analysis: Lee CW, Kong C. Methodology: Kong C. Software: Lee CW, Kong C. Validation: Lee CW, Kong C. Investigation: Lee CW, Kong C. Writing - original draft: Lee CW, Kong C. Writing - review & editing: Lee CW, Kong C.
Ethics approval and consent to participate	The protocols for the present study were reviewed and approved by the Institutional Animal Care and Use Committee of Kyungpook National University (KNU 2023-0226).

CORRESPONDING AUTHOR CONTACT INFORMATION

For the corresponding author (responsible for correspondence, proofreading, and reprints)	Fill in information in each box below
First name, middle initial, last name	Changsu Kong
Email address – this is where your proofs will be sent	changsupkong@knu.ac.kr
Secondary Email address	
Address	2559, Gyeongsang-daero, Sangju-si, Gyeongsangbuk-do, 37224, Korea

Cell phone number	+82-10-3805-4776
Office phone number	+82-54-530-1225
Fax number	+82-54-530-1229

6
7

ACCEPTED

(Unstructured) Abstract (up to 350 words)

The current study aimed to estimate the standardized ileal digestible (SID) calcium (Ca) and phosphorus (P) requirements of broiler chickens based on response surface methodology (RSM). Nine experimental diets were formulated with different SID Ca and P concentrations (2.80 and 5.50, 3.44 and 4.44, 3.44 and 6.56, 5.00 and 7.00, 5.00 and 5.50, 5.00 and 4.00, 6.56 and 4.44, 6.56 and 6.56, and 7.20 and 5.50 g/kg, respectively). A total of 480 10-day-old Ross 308 male broilers were weighed and randomly assigned to nine treatments based on body weight. Each treatment had five replicates, except for the central treatment (5.00 g/kg Ca and 5.50 g/kg P), which had eight replicates with 10 birds per pen. On day 21, body weight and feed leftovers were recorded to calculate body weight gain (BWG) and the gain-to-feed ratio (G:F). Left tibia bones were also collected for compositional analysis and bone mineral density (BMD) assessment. Response surface analysis revealed significant quadratic models for all criteria. The maximum BWG was estimated at 5.13 and 5.74 g/kg for SID Ca and P, respectively. The maximum G:F was observed when 6.41 and 7.00 g/kg of SID Ca and P were used, respectively. Multi-objective optimization analysis demonstrated that 6.02 g/kg of SID Ca and 6.61 g/kg of SID P were required to achieve both optimal BWG and G:F. Furthermore, the ideal SID Ca and P concentrations for optimal tibia ash, tibia Ca, tibia P, and BMD were estimated at 7.20 and 7.00 g/kg, 5.75 and 5.87 g/kg, 7.20 and 7.00 g/kg, and 7.20 and 6.96 g/kg, respectively. Multi-objective optimization indicated that 6.50 and 6.83 g/kg of SID Ca and P, respectively, are required to achieve optimal growth performance and bone mineralization. This study's findings suggest that RSM is a feasible and effective approach to determining the optimal SID Ca and P requirements of broiler chickens, as it efficiently evaluates multiple factors while considering several response criteria.

Keywords (3 to 6):

Broiler, Calcium, Phosphorus, Requirement, Standardized ileal digestible, Response surface methodology

Introduction

Calcium (Ca) and phosphorus (P) are the most abundant macro-minerals deposited in bones, playing important roles in nutrient metabolism and bone mineralization [1, 2]. Ca, which acts as a cation in the digestive tract, increases intestinal pH. Additionally, it reduces the digestibility of both Ca and P by binding to phytate and forming a calcium–phytate complex [3-5]. Since Ca and P concentrations influence each other in terms of digestion and absorption, their interaction should be considered when seeking to enhance growth performance and promote the efficient utilization of minerals. Recent studies have investigated the interactive effects of Ca and P on their availability in broilers by simultaneously adjusting their dietary levels in experimental diets [6-8].

In general, poultry diet is formulated based on the total Ca and available or non-phytate P (NPP) concentrations. “Available nutrients” refer to the actual amount of dietary nutrients that are digested and absorbed by an animal, with the commonly used NPP content representing the available P content in diet [9]. However, this does not account for the fact that NPP is not fully utilized, and some of the phytate P is also utilized [10]. Therefore, as digestibility has become a common criterion for evaluating the nutrient availability of poultry feed ingredients, there has been a shift toward using digestible Ca and P bases for diet formulation, rather than total Ca and available P or NPP bases [11, 12]. Several studies have measured the standardized ileal digestibility values of Ca and P, which have been corrected for endogenous losses from apparent ileal digestibility [13-15].

Response surface methodology (RSM) is a mathematical and statistical method used to analyze the interactive effects of multiple factors on response variables and determine the optimal factor settings for the best response [16, 17]. RSM helps evaluate multiple parameters and their interactions using quantitative data and has been used in various experiments, including studies on broilers, to determine nutrient requirements in animal nutrition [18-20]. Growth performance and bone mineralization have been evaluated as response variables owing to their sensitivity to dietary Ca and P concentrations [21, 22]. While certain studies have assessed dietary Ca and P requirements using RSM based on total Ca and NPP levels, to the best of our knowledge, no study has simultaneously estimated standardized ileal digestible (SID) Ca and P requirements. Therefore, this study aimed to estimate SID Ca and P requirements using RSM by measuring growth performance and bone mineralization in broiler chickens fed diets with varying SID Ca and P levels.

Materials and Methods

58 **Animal ethics**

59 The Institutional Animal Care and Use Committee of Kyungpook National University, Republic of Korea,
60 reviewed and approved all experimental procedures (approval number: KNU 2023-0226).

61

62 **Animals, management, and experimental design**

63 A total of 480 10-day-old Ross 308 male broiler chickens were randomly allocated to nine dietary treatments
64 using a five-level, two-factor central composite design (CCD). Each treatment was replicated five times, except for the
65 central treatment (run no. 5), which had eight replicates), and each replicate comprised 10 birds. From d 10 to 21, birds
66 were allowed *ad libitum* access to water and experimental diets for 11 days. House temperature was maintained at 34°C
67 on the first day and gradually reduced to 25°C by 21 days of age at a rate of 3°C per week. All birds were housed in 1 m
68 × 1 m floor pens under a continuous 24-hour lighting program. Lighting intensity was maintained at 30 lux throughout
69 the experiment.

70

71 **Dietary treatments**

72 SID Ca and P concentrations in feed ingredients were obtained from previous research in the same laboratory
73 [23]. A corn–soybean meal basal diet was formulated to meet or exceed nutrient specifications of broilers [24], except
74 those for Ca and P (Table 1). All experimental diets were isoenergetic and isonitrogenous. The basal diet was adjusted
75 with appropriate amounts of cornstarch, soybean oil, limestone, monocalcium phosphate, and monosodium phosphate to
76 formulate nine experimental diets containing varying SID Ca and SID P concentrations, as indicated in Table 1: Diet 1:
77 2.80 and 5.50 g/kg, Diet 2: 3.44 and 4.44 g/kg, Diet 3: 3.44 and 6.56 g/kg, Diet 4: 5.00 and 7.00 g/kg, Diet 5 (central
78 treatment): 5.00 and 5.50 g/kg, Diet 6: 5.00 and 4.00 g/kg, Diet 7: 6.56 and 4.44 g/kg, Diet 8: 6.56 and 6.56 g/kg, Diet
79 9: 7.20 and 5.50 g/kg.

80

81 **Measurement and chemical analyses**

82 On day 10 and 21, all chickens were individually weighed, and feed intake was recorded to determine the mean
83 body weight, body weight gain (BWG), and the gain-to-feed ratio (G:F). Mortality was monitored daily and used to
84 adjust the feed intake values. At the end of the experiment (21 days of age), two birds with body weights representing

the median from each pen euthanized by CO₂ asphyxiation for left tibias collection. The tibias were placed in labeled plastic bags and stored in a freezer at -20°C until further analysis. After separating the meat and soft tissues from each tibia, the tibias were defatted by soaking in ethyl ether for 48 h and subsequently dried to a constant weight using a drying oven at 105°C for 24 h. Bone mineral density (BMD) was determined using dual-energy X-ray absorptiometry (InAlyzer; Medikors Inc., South Korea). Thereafter, the samples were ashed in a muffle furnace at 550°C for 12 h to determine tibial ash, total Ca, and total P contents. All experimental diets were ground using a mill grinder (CT 293 Cyclotec™; Foss Ltd., Denmark) through a 1.0-mm screen. The total Ca and P concentrations in the experimental diets and tibia were analyzed using inductively coupled plasma-optical emission spectrometry (Optima 8300; PerkinElmer, USA) [25].

Statistical Analyses

Experimental data were analyzed using the MIXED procedure of SAS 9.4 (SAS Inst. Inc., Cary, NC, USA). The statistical model included dietary treatment as the fixed variable and the replications as random variables. Statistical significance was set at $p < 0.05$. Where dietary effects were significant, the Tukey–Kramer method was used to compare mean values among treatments.

In this experiment, a two-factor, five-level CCD for SID Ca (2.8, 3.4, 5.0, 6.6, and 7.2 g/kg) and SID P (4.0, 4.4, 5.5, 6.6, and 7.0 g/kg) was used. Each pen was set as an experimental unit. Experimental data were fitted to the following second-order polynomial regression equation:

$$Y = \beta_0 + \beta_i x_i + \beta_j x_j + \beta_{ii} x_i^2 + \beta_{jj} x_j^2 + \beta_{ij} x_i x_j + \varepsilon$$

where Y is the response of interest; β_0 denotes the intercept; and β_i , β_j , β_{ii} , β_{jj} , and β_{ij} are the coefficients estimated by the model. Furthermore, x_i and x_j represent SID Ca and SID P, respectively, and ε is the residual associated with the experiment. The experimental data were analyzed using Design-Expert® software (version 13; Stat-Ease Inc., Minneapolis, USA) to optimize requirements. R² indicates the extent to which each independent variable is interpreted concerning changes in the dependent variable and is used to assess how closely the predicted values align with the actual values [26]. The discrepancy between the predicted R² and the adjusted R² values was examined to confirm the regression fit, ensuring it was < 0.2. The desirability function and importance were used to evaluate the effectiveness of each factor in determining the SID Ca and P requirements of chickens. Desirability scores range from 0 to 1, and the value closest to 1 was selected as the optimal point. According to Design-Expert software, importance

values range from 1 to 5, with a score of 5 assigned to growth performance and that of 3 to bone mineralization. However, while tibial ash and mineral contents exhibited a quadratic response, the “lack of fit” was significant. Therefore, an importance level of 1 was assigned to tibial ash and mineral content for multi-objective optimization.

Results

During the experimental period, the mortality rate was 0.62%, and it was not influenced by the experimental diets. The total Ca and P contents of the experimental diets were analyzed on an as-fed basis, as expected (Table 1). Increasing supplementation of SID Ca and P exerted a quadratic effect ($p < 0.05$) on growth performance (Table 2) and bone mineralization (Table 3), indicating that the regression model was adequate to determine requirements. The lack-of-fit test of the RSM model was not significant for BWG, G:F, and BMD. The difference between the predicted R^2 and the adjusted R^2 values was < 0.2 , indicating that the quadratic model adequately fit the data for all response criteria.

Birds fed Diet 5 exhibited the greatest BWG, and Diets 2, 4, 5, and 8 yielded the highest G:F values (Table 2). As shown in Table 3, bone mineralization also displayed differences among treatments, although the values were comparable. The tibia ash, Ca, and P contents were highest in Diets 5 and 9, Diets 5 and 9, and Diets 5 and 8, respectively, while BMD peaked with Diet 8 (Table 3). Single-objective optimization indicated that the estimated optimal values of SID Ca and P for maximal BWG and G:F were 5.13 and 5.74 g/kg and 6.41 and 7.00 g/kg, respectively (Table 4). The ideal SID Ca and P concentrations for tibia ash, tibia Ca, tibia P, and BMD were 7.20 and 7.00 g/kg, 5.75 and 5.87 g/kg, 7.20 and 7.00 g/kg, and 7.20 and 6.96 g/kg, respectively (Table 4). Multi-objective optimization demonstrated that 6.02 g/kg of SID Ca and 6.61 g/kg of SID P were required to optimize growth performance, while 7.13 g/kg of SID Ca and 6.80 g/kg of SID P were required to maximize bone mineralization (Table 4). The most desirable concentrations of SID Ca and P for all responses were 6.50 and 6.83 g/kg, respectively (Table 4).

Discussion

Ca and P homeostasis is determined by the interaction between these minerals, and since their respective concentrations influence their utilization, their interrelationships serve an important role in defining mineral requirements [27]. Therefore, when estimating Ca or P requirements, formulating experimental diets that simultaneously consider both factors rather than setting only one nutrient as an independent variable is more accurate. Several factorial experiments have been conducted to examine the interaction between Ca and P by concurrently

considering their levels [8, 28, 29]. Establishing SID Ca and P levels at five levels each, as in the current study, would typically require 25 treatments under a 5×5 full factorial arrangement. However, applying a CCD, a more efficient alternative, reduces the number of dietary treatments required in poultry research to only nine [17]. Specifically, the CCD is designed to generate maximal information regarding parameters from a minimal number of treatments by identifying three distinct points: a central point as well as factorial and axial points, which are equidistantly positioned from the central point [30, 31]. Moreover, as a design applied in RSM, the CCD incorporates five levels for independent factors, facilitating the statistical optimization of variables that influence responses [32].

Based on a literature review, SID Ca and P requirements have been estimated to be 5.0–5.5 g/kg. Therefore, in this study, the dietary SID Ca and P concentrations of the experimental diets were adjusted accordingly to ensure that the estimated values were centered within the CCD framework. Subsequently, single-objective optimization identified 5.13 and 5.74 g/kg as the estimated optimal SID Ca and P levels for maximal BWG, respectively, within the expected range. Since feed intake did not exhibit a significant difference among treatments, it was excluded from the multi-objective optimization analysis. Multi-objective optimization revealed that 6.02 g/kg of SID Ca and 6.61 g/kg of SID P were required to maximize BWG and G:F, while 7.13 g/kg of SID Ca and 6.80 g/kg of SID P were required to maximize tibia ash, tibia Ca, tibia P, and BMD. Meanwhile, the SID Ca requirement based on tibia ash, tibia P, and BMD was determined to be 7.2 g/kg, equating to the highest SID Ca concentration incorporated into the experimental diets. This finding was unanticipated, considering that the total dietary Ca and P levels exceeded the recommended Aviagen specifications. The comparatively high SID Ca and P requirements generated based on bone mineralization probably emanated from its sensitivity to Ca and P concentrations. Tibia bone development in broilers is commonly evaluated using parameters reflecting bone mineralization [8, 28]. Research has consistently demonstrated that broilers require higher levels of NPP when bone mineralization serves as the evaluation criterion compared to when growth performance is used [33]. Approximately 99% of Ca and 80% of P are stored in the bones, while the remaining 20% of P participates in various metabolic processes, playing essential roles in growth, cellular and membrane function, energy metabolism, and acid–base balance [34]. Low plasma Ca concentrations stimulate parathyroid hormone release, triggering Ca and P release from bones to maintain homeostasis [34]. Similarly, low plasma P concentrations cause bone resorption, providing the P required for bird maintenance and growth [35]. Considering this physiological regulation, the dietary Ca and P consumed by broilers primarily accumulate in the bones, and as bones serve as a storage reservoir for Ca and P, they may release these minerals when the amounts required for maintenance and growth become insufficient. In this study, broilers were fed a pre-starter diet that met nutrient recommendations up to day 10,

170 ensuring the sufficient storage of Ca and P in the bones. Consequently, even when experimental diets with lower Ca and
171 P levels than those recommended by Aviagen were administered, the birds' stored mineral reserves potentially
172 compensated for the deficiency, allowing them to maintain essential physiological functions. The findings of current
173 study are consistent with those of Walk et al. [36], who reported no effect of graded SID Ca levels (0.16–0.56%) on
174 feed intake or BWG, whereas tibial ash content responded quadratically to these levels. Additionally, our results align
175 with the findings of Walk et al. [21], who observed no linear or quadratic effects on BWG in birds fed a diet with SID
176 Ca levels of 0.2–0.6%. However, they found significant differences in tibial ash concentration depending on the SID Ca
177 level. In broilers, Fallah et al. [7]] also observed higher total Ca and NPP requirements for bone mineralization than for
178 performance traits, emphasizing the role of Ca and P in regulating bone mineralization and growth performance. These
179 findings suggest that tibial mineralization, as a response criterion, may be a more sensitive measure of Ca and P
180 requirements than growth performance. Moreover, future studies should consider higher maximum dietary SID Ca and
181 P levels to obtain more accurate estimations.

182 The current study found the optimal SID Ca-to-SID P ratio to be 0.95, based on all analyzed criteria. The diets
183 with similar ratios, listed in order, were Diets 5 (5 g/kg Ca), 8 (6.56 g/kg Ca), and 2 (3.44 g/kg Ca). Diet 5 yielded the
184 greatest BWG, while Diets 5 and 8 yielded the highest tibial ash content. Moreover, the tibia Ca and P contents were
185 highest in Diets 5 and 9 and Diets 5 and 8, respectively, while BMD peaked with Diet 8. These results indicate that the
186 experimental diets with SID Ca-to-SID P ratios approximating 0.95 might have contributed to improvements in growth
187 performance and bone mineralization. David et al. [37] estimated the SID Ca requirement for maximum weight gain in
188 broilers aged 24 days to be 3.05 g/kg, while that for maximum tibial ash content was 3.69 g/kg, with a recommended
189 SID P concentration of 3.5 g/kg. These values indicated SID Ca-to-SID P ratios of 1.05 and 0.87, respectively. The
190 relatively high SID Ca requirements observed in the current study possibly reflect differences in the evaluated SID Ca
191 and P concentration ranges. While David et al. [37] examined SID Ca levels of 1.8–4.55 g/kg, the current study used
192 wider ranges: 2.8–7.2 g/kg for SID Ca and 4.4–7.0 g/kg for SID P. However, the SID Ca-to-SID P ratio of 0.95
193 obtained in the current study remained within the range (1.05–0.87) suggested by David et al. [37], suggesting partial
194 alignment between the findings of the two experiments. Furthermore, their study demonstrated that SID Ca
195 requirements potentially increase with increasing dietary P levels, a trend consistent with the results of the current study
196 [37].

197 In the current study, Diet 7, which had the highest SID Ca-to-SID P ratio, resulted in the lowest BWG and G:F
198 values. Gautier et al. [38] demonstrated that when the NPP content was fixed and Ca content increased to amplify the

Ca ratio, growth performance decreased with increasing Ca levels. However, when the Ca-to-NPP ratio was fixed at 2:1, and both dietary Ca and P levels had been elevated, growth performance was not affected by dietary treatments [38]. These results are consistent with those of Fallah et al. [7], who reported that increasing dietary Ca levels, particularly in low-NPP diets, negatively affected the performance of broiler chickens. High dietary Ca levels have been reported to decrease growth performance in broiler chickens, and this decrease can be alleviated by increasing the dietary NPP level [39, 40]. The reduction in growth performance observed with high dietary Ca levels possibly resulted from the effects of Ca circulation processes aimed at maintaining homeostasis in the body. Once Ca is digested, it is absorbed into the bloodstream and subsequently deposited in the bones [41]. High plasma Ca levels stimulate the secretion of calcitonin, which inhibits osteoclast activity and promotes the renal excretion of both Ca and P [42, 43]. In another study, the maintenance of a balanced Ca:NPP ratio in a low-P diet did not affect serum P levels, suggesting that an imbalanced Ca:NPP ratio potentially influences fluctuations in serum P levels [44]. Additionally, high Ca levels raise the pH of the gastrointestinal tract, reducing the soluble fraction of other nutrients and consequently decreasing their availability and absorption [45, 46]. Li et al. [47] reported that at an NPP level of 1.9 g/kg, increasing dietary Ca levels from 6.5 to 9.5 g/kg decreased apparent ileal P digestibility. Furthermore, phytate can bind proteins to form protein-phytate complexes, which depress protein utilization [3, 48]. These adverse effects suggest that an imbalance of dietary Ca and P may diminish growth performance. In a study by Noruzi et al. [49], reducing dietary Ca and available P by up to 30% while maintaining the same ratio did not affect performance. Moreover, since broilers can adapt to diets with low dietary P and Ca levels by increasing digestibility and absorption [50], considering both the absolute concentrations and Ca-to-P ratio may provide a more accurate estimation of Ca and P requirements.

The optimal conditions for the multi-objective optimization of growth performance and bone mineralization in broilers from day 10 to 21 were achieved at SID Ca and P levels of 6.50 and 6.83 g/kg, respectively, at an SID Ca-to-SID P ratio of 0.95. Based on growth performance, the optimal levels of SID Ca (6.02 g/kg) and SID P (6.61 g/kg) estimated in this study correspond to approximately 7.9–8.3 g/kg of total Ca and 3.7–4.2 g/kg of NPP. These values are lower than those recommended by NRC [51] for both Ca and NPP. Compared to the Aviagen guidelines, NPP levels were similar, but Ca levels were estimated to be slightly higher. Since optimal bone development requires higher levels of Ca and P than those needed for growth performance alone, nutrient requirement recommendations based solely on performance outcomes may not adequately reflect the levels necessary for proper skeletal health and development. Additionally, although the diets had identical SID values, differences in total Ca and NPP concentrations resulting from the variation in ingredient composition limit the direct comparison of the present results with existing feeding

228 guidelines. The results also suggest that further research is required to establish the precise requirements by varying SID
229 Ca and P levels while maintaining a fixed SID Ca-to-SID P ratio. The findings of the current study also suggest that
230 using RSM to determine the optimal SID Ca and P requirements for broiler chickens is a feasible and effective approach,
231 as it enables the simultaneous evaluation of multiple factors while considering response criteria to optimize nutrient
232 requirements efficiently.

233

234

Acknowledgments

235 This work was carried out with the support of "Cooperative Research Program for Agriculture Science and
236 Technology Development (Project No. RS-2022-RD010370)" Rural Development Administration, Republic of Korea.

ACCEPTED

References (Vancouver or NLM style)

- 238 1. Takeda E, Yamamoto H, Yamanaka-Okumura H, Taketani Y. Dietary phosphorus in bone health and quality of
239 life. *Nutr Rev.* 2012;70:311-21. 10.1111/j.1753-4887.2012.00473.x.
- 240 2. David LS, Anwar MN, Abdollahi MR, Bedford MR, Ravindran V. Calcium nutrition of broilers: current
241 perspectives and challenges. *Animals (Basel).* 2023;13:1590. 10.3390/ani13101590.
- 242 3. Selle PH, Ravindran V, Caldwell A, Bryden WL. Phytate and phytase: consequences for protein utilisation.
243 *Nutr Res Rev.* 2000;13:255-78. 10.1079/095442200108729098.
- 244 4. Bradbury EJ, Wilkinson SJ, Cronin GM, Thomson PC, Bedford MR, Cowieson AJ. Nutritional geometry of
245 calcium and phosphorus nutrition in broiler chicks. Growth performance, skeletal health and intake arrays.
246 *Animal.* 2014;8:1071-9. 10.1017/S1751731114001037.
- 247 5. Sung JY, Emmert BJ, Karcher DM, Walk CL, Adeola O. Phosphorus equivalency of exogenous phytase
248 relative to phosphorus in monosodium phosphate in broiler chickens. *Poult Sci.* 2024;103:103248.
249 10.1016/j.psj.2023.103248.
- 250 6. Liu JB, Chen DW, Adeola O. Phosphorus digestibility response of broiler chickens to dietary calcium-to-
251 phosphorus ratios. *Poult Sci.* 2013;92:1572-8. 10.3382/ps.2012-02758.
- 252 7. Fallah H, Karimi A, Sadeghi A, Behroozi-Khazaei N. Modelling and optimizing of calcium and non-phytate
253 phosphorus requirements of male broiler chickens from 1 to 21 days of age using response surface
254 methodology. *Animal.* 2020;14:1598-609. 10.1017/S1751731120000452.
- 255 8. Wu J, Ma X, Liao X, Song C, Li S, Zhang L, et al. Dietary calcium and nonphosphate phosphorus interaction
256 influences tibiotarsus development and related gene expression of broilers from 1 to 21 days of age. *Poult Sci.*
257 2023;102:102851. 10.1016/j.psj.2023.102851.
- 258 9. Rodehutscord M, WPSA WGNNotEFoBo. Determination of phosphorus availability in poultry. *World's Poult*
259 *Sci J.* 2013;69:687-98. 10.1017/s0043933913000688.
- 260 10. Leske K, Coon C. The development of feedstuff retainable phosphorus values for broilers. *Poult Sci.*
261 2002;81:1681-93. 10.1093/ps/81.11.1681.
- 262 11. Adedokun SA, Pescatore AJ, Ford MJ, Ao T, Jacob JP. Investigating the effect of dietary calcium levels on
263 ileal endogenous amino acid losses and standardized ileal amino acid digestibility in broilers and laying hens.
264 *Poult Sci.* 2018;97:131-9. 10.3382/ps/pex271.
- 265 12. David LS, Abdollahi MR, Bedford MR, Ravindran V. Requirement of digestible calcium at different dietary
266 concentrations of digestible phosphorus for broiler chickens. 1. Broiler starters (d 1 to 10 post-hatch). *Poult Sci.*
267 2021;100:101439. 10.1016/j.psj.2021.101439.

- 268 13. Li W, Angel R, Plumstead PW, Enting H. Effects of limestone particle size, phytate, calcium source, and
269 phytase on standardized ileal calcium and phosphorus digestibility in broilers. *Poult Sci.* 2021;100:900-9.
270 10.1016/j.psj.2020.10.075.
- 271 14. An SH, Sung JY, Kong C. Ileal digestibility and total tract retention of phosphorus in inorganic phosphates fed
272 to broiler chickens using the direct method. *Animals (Basel).* 2020;10:2167. 10.3390/ani10112167.
- 273 15. Dilelis F, de Freitas LW, Quaresma DV, Machado NDB, Reis TL, Souza CS, et al. Standardized ileal
274 phosphorus digestibility of meat and bone meal and poultry byproduct meal for broilers. *Rev Bras Zootecn.*
275 2021;50:e20200086. ARTN e2020008610.37496/rbz5020200086.
- 276 16. Karimifard S, Alavi Moghaddam MR. Application of response surface methodology in physicochemical
277 removal of dyes from wastewater: A critical review. *Sci Total Environ.* 2018;640-641:772-97.
278 10.1016/j.scitotenv.2018.05.355.
- 279 17. Mehri M. Optimization of response surface and neural network models in conjugation with desirability
280 function for estimation of nutritional needs of methionine, lysine, and threonine in broiler chickens. *Poult Sci.*
281 2014;93:1862-7. 10.3382/ps.2013-03689.
- 282 18. Ahmadi H, Golian A. Response surface and neural network models for performance of broiler chicks fed diets
283 varying in digestible protein and critical amino acids from 11 to 17 days of age. *Poult Sci.* 2011;90:2085-96.
284 10.3382/ps.2011-01367.
- 285 19. Mehri M, Davarpanah AA, Mirzaei HR. Estimation of ideal ratios of methionine and threonine to lysine in
286 starting broiler chicks using response surface methodology. *Poult Sci.* 2012;91:771-7. 10.3382/ps.2011-01818.
- 287 20. Yolmeh M, Habibi Najafi MB, Farhoosh R. Optimisation of ultrasound-assisted extraction of natural pigment
288 from annatto seeds by response surface methodology (RSM). *Food Chem.* 2014;155:319-24.
289 10.1016/j.foodchem.2014.01.059.
- 290 21. Walk CL, Wang Z, Wang S, Wu J, Sorbara JOB, Zhang J. Determination of the standardized ileal digestible
291 calcium requirement of male Arbor Acres Plus broilers from hatch to day 10 post-hatch. *Poult Sci.*
292 2021;100:101364. 10.1016/j.psj.2021.101364.
- 293 22. Bai S, Yang Y, Ma X, Liao X, Wang R, Zhang L, et al. Dietary calcium requirements of broilers fed a
294 conventional corn-soybean meal diet from 1 to 21 days of age. *J Anim Sci Biotechnol.* 2022;13:11.
295 10.1186/s40104-021-00652-5.
- 296 23. Lee CW, Kong C. Standardized ileal digestibility of calcium and phosphorus in feed ingredients for 21-day-old
297 broilers. *Animals (Basel).* 2024;14:2603. 10.3390/ani14172603.
- 298 24. Aviagen. Ross 308: Broiler nutrition specifications. Aviagen Inc., Huntsville, AL. 2022.
- 299 25. MAFRA [Ministry of Agriculture FaRA. Statistical yearbook of agriculture, food and rural affairs. 2022.
300 2022Report No.: 11-1543000-000261-10.

- 301 26. Parhi SK, Patro SK. Prediction of compressive strength of geopolymers using a hybrid ensemble of
302 grey wolf optimized machine learning estimators. *J Build Eng.* 2023;71:106521. ARTN
303 10652110.1016/j.jobbe.2023.106521.
- 304 27. Li X, Zhang D, Bryden W. Calcium and phosphorus metabolism and nutrition of poultry: are current diets
305 formulated in excess? *Anim Prod Sci.* 2017;57:2304-10. doi.org/10.1071/AN17389.
- 306 28. Li T, Xing G, Shao Y, Zhang L, Li S, Lu L, et al. Dietary calcium or phosphorus deficiency impairs the bone
307 development by regulating related calcium or phosphorus metabolic utilization parameters of broilers. *Poult Sci.*
308 2020;99:3207-14. 10.1016/j.psj.2020.01.028.
- 309 29. Wang YB, Wang WW, Fan QL, Ye JL, Zhang S, Jiang SQ. Effects and interaction of dietary calcium and non-
310 phytate phosphorus for slow-growing yellow-feathered broilers during the starter phase. *Animal.*
311 2021;15:100201. 10.1016/j.animal.2021.100201.
- 312 30. Yolmeh M, Jafari SM. Applications of Response Surface Methodology in the Food Industry Processes. *Food*
313 *Bioprocess Tech.* 2017;10:413-33. 10.1007/s11947-016-1855-2.
- 314 31. Boukroufa M, Boutekedjiret C, Petigny L, Rakotomanomana N, Chemat F. Bio-refinery of orange peels waste:
315 a new concept based on integrated green and solvent free extraction processes using ultrasound and microwave
316 techniques to obtain essential oil, polyphenols and pectin. *Ultrason Sonochem.* 2015;24:72-9.
317 10.1016/j.ultsonch.2014.11.015.
- 318 32. Beg S, Rahman Z. Central composite designs and their applications in pharmaceutical product development
319 2021.
- 320 33. Liu SB, Liao XD, Lu L, Li SF, Wang L, Zhang LY, et al. Dietary non-phytate phosphorus requirement of
321 broilers fed a conventional corn-soybean meal diet from 1 to 21 d of age. *Poult Sci.* 2017;96:151-9.
322 10.3382/ps/pew212.
- 323 34. Proszkowiec-Weglarz M, Angel R. Calcium and phosphorus metabolism in broilers: Effect of homeostatic
324 mechanism on calcium and phosphorus digestibility. *J Appl Poultry Res.* 2013;22:609-27. 10.3382/japr.2012-
325 00743.
- 326 35. Nari N, Ghasemi HA. Growth performance, nutrient digestibility, bone mineralization, and hormone profile in
327 broilers fed with phosphorus-deficient diets supplemented with butyric acid and *Saccharomyces boulardii*.
328 *Poult Sci.* 2020;99:926-35. 10.1016/j.psj.2019.10.046.
- 329 36. Walk CL, Wang Z, Wang S, Sorbara JOB, Zhang J. Determination of the standardized ileal digestible calcium
330 requirement of male Arbor Acres Plus broilers from day 11 to 24 post-hatch. *Poult Sci.* 2022;101:101836.
331 10.1016/j.psj.2022.101836.
- 332 37. David L, Abdollahi M, Bedford M, Ravindran V. Requirement of digestible calcium at different dietary
333 concentrations of digestible phosphorus for broiler chickens. 2. Broiler growers (d 11 to 24 post-hatch). *Poult*
334 *Sci.* 2022;101:102135. doi.org/10.1016/j.psj.2022.102135.

- 335 38. Gautier AE, Walk CL, Dilger RN. Influence of dietary calcium concentrations and the calcium-to-non-phytate
336 phosphorus ratio on growth performance, bone characteristics, and digestibility in broilers. *Poult Sci.*
337 2017;96:2795-803. 10.3382/ps/pex096.
- 338 39. Sebastian S, Touchburn SP, Chavez ER, Lague PC. Efficacy of supplemental microbial phytase at different
339 dietary calcium levels on growth performance and mineral utilization of broiler chickens. *Poult Sci.*
340 1996;75:1516-23. 10.3382/ps.0751516.
- 341 40. Rao SVR, Raju MVLN, Reddy MR, Pavani P. Interaction between dietary calcium and non-phytate phosphorus
342 levels on growth, bone mineralization and mineral excretion in commercial broilers. *Anim Feed Sci Tech.*
343 2006;131:133-48. 10.1016/j.anifeedsci.2006.02.011.
- 344 41. Zhang QQ, Chang C, Chu Q, Wang HH, Zhang J, Yan ZX, et al. Dietary calcium and non-phytate phosphorus
345 levels affect the performance, serum biochemical indices, and lipid metabolism in growing pullets. *Poult Sci.*
346 2023;102:102354. 10.1016/j.psj.2022.102354.
- 347 42. Pondel M. Calcitonin and calcitonin receptors: bone and beyond. *Int J Exp Pathol.* 2000;81:405-22.
348 10.1046/j.1365-2613.2000.00176.x.
- 349 43. Matuszewski A, Lukasiewicz M, Niemiec J. Calcium and phosphorus and their nanoparticle forms in poultry
350 nutrition. *World's Poult Sci J.* 2020;76:328-45. 10.1080/00439339.2020.1746221.
- 351 44. Xu L, Li N, Farnell YZ, Wan XL, Yang HM, Zhong XQ, et al. Effect of feeding a high calcium: phosphorus
352 ratio, phosphorous deficient diet on hypophosphatemic rickets onset in broilers. *Agriculture-Basel.*
353 2021;11:955. ARTN 95510.3390/agriculture11100955.
- 354 45. Selle PH, Cowieson AJ, Ravindran V. Consequences of calcium interactions with phytate and phytase for
355 poultry and pigs. *Livest Sci.* 2009;124:126-41. 10.1016/j.livsci.2009.01.006.
- 356 46. Tancharoenrat P, Ravindran V. Influence of tallow and calcium concentrations on the performance and energy
357 and nutrient utilization in broiler starters. *Poult Sci.* 2014;93:1453-62. 10.3382/ps.2013-03817.
- 358 47. Li W, Angel R, Kim SW, Jiménez-Moreno E, Proszkowiec-Weglarz M, Plumstead PW. Impacts of age and
359 calcium on Phytase efficacy in broiler chickens. *Anim Feed Sci Tech.* 2018;238:9-17.
360 10.1016/j.anifeedsci.2018.01.021.
- 361 48. Humer E, Schwarz C, Schedle K. Phytate in pig and poultry nutrition. *J Anim Physiol Anim Nutr (Berl).*
362 2015;99:605-25. 10.1111/jpn.12258.
- 363 49. Noruzi H, Hassanabadi A, Golian A, Aziz-Aliabadi F. Effects of dietary calcium and phosphorus restrictions
364 on growth performance, intestinal morphology, nutrient retention, and tibia characteristics in broiler chickens.
365 *Br Poult Sci.* 2023;64:231-41. 10.1080/00071668.2022.2136510.
- 366 50. Yan F, Angel R, Ashwell C, Mitchell A, Christman M. Evaluation of the broiler's ability to adapt to an early
367 moderate deficiency of phosphorus and calcium. *Poult Sci.* 2005;84:1232-41. 10.1093/ps/84.8.1232.

368 51. NRC(National Research Council). Nutrient requirements of poultry, 9th. rev. ed.: Natl. Acad. Press,
369 Washington, DC; 1994.

ACCEPTED

Tables and Figures

Table 1. Ingredient and chemical compositions of experimental diets on an as-fed basis.

Ingredient, g/kg	Dietary treatments								
	1	2	3	4	5	6	7	8	9
Corn	560.0	560.0	560.0	560.0	560.0	560.0	560.0	560.0	560.0
Soybean meal	330.0	330.0	330.0	330.0	330.0	330.0	330.0	330.0	330.0
Cornstarch	61.0	60.4	51.0	40.5	43.0	45.5	29.3	25.8	21.0
Soybean oil	14.9	15.2	19.5	24.4	23.2	22.1	29.6	31.2	33.4
Sodium bicarbonate	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Limestone	-	5.5	-	2.8	9.3	15.8	22.2	13.0	21.1
Monocalcium phosphate	11.1	8.6	15.3	22.0	14.2	6.3	8.6	19.7	14.2
Monosodium phosphate	2.7	-	3.9	-	-	-	-	-	-
Sodium chloride	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0
Vitamin premix ¹	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Mineral premix ²	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Choline chloride	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0
L-Arg	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
L-Ile	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7
L-Lys-HCl	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4
L-Met	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6
L-Cys	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6
L-Thr	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3
L-Val	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1

Calculated value

AME, kcal/kg	3000.0	3000.0	3000.0	3000.0	3000.0	3000.0	3000.0	3000.0	3000.0
Crude protein	203.3	203.3	203.3	203.3	203.3	203.3	203.3	203.3	203.3
Total Ca	3.3	4.8	4.1	6.3	7.2	8.1	10.7	9.5	11.4
Total P	6.3	5.2	7.5	8.0	6.4	4.7	5.2	7.5	6.4
Non-phytate P	3.8	2.7	5.0	5.5	3.8	2.2	2.7	5.0	3.8
SID Ca ³	2.80	3.44	3.44	5.00	5.00	5.00	6.55	6.56	7.20
SID P ³	5.50	4.44	6.56	7.00	5.50	4.00	4.44	6.56	5.50
SID Ca: SID P ratio	0.51	0.77	0.52	0.71	0.91	1.25	1.48	1.00	1.31
Analyzed value									
Total Ca	3.6	5.5	4.4	6.4	7.5	7.2	11.4	9.8	11.7
Total P	6.5	5.5	7.6	8.0	6.7	4.7	5.7	8.0	6.7

AME=apparent metabolizable energy; Ca=calcium; P=phosphorus; SID=standardized ileal digestible.

¹Supplies the following quantities per kilogram of diet: vitamin A, 18,000 IU; vitamin D₃, 6,000 IU; vitamin E, 75 mg; Vitamin K₃, 5 mg; Thiamin, 5 mg; Riboflavin, 13 mg; Nicotinic acid, 90 mg; Pantothenic acid, 30 mg; Pyridoxine, 6.8 mg; cobalamin, 0.03 mg; Folic acid, 3.3 mg; Biotin 0.33 mg.

²Supplies the following quantities per kilogram of diet: Mn, 180 mg; Zn, 165 mg; Fe, 75 mg; Co, 24 mg; I, 1.9 mg; Se, 0.5 mg.

³Based on SID values determined in a previous digestibility study conducted in the same laboratory [23].

Table 2. Growth performance of broiler chickens fed diets varying in concentrations of standardized ileal digestible (SID) calcium (Ca) and phosphorous (P) from d 10 to 21.

Input variables, g/kg			Output variables			
Run no. (n ¹)	SID Ca	SID P	BWG (g)	SEM	G:F (g/g)	SEM
1 (5)	2.80	5.50	495.8 ^{bcd}	11.40	0.72 ^{ab}	0.008
2 (5)	3.44	4.44	542.2 ^{ab}	11.40	0.75 ^a	0.008
3 (5)	3.44	6.56	483.0 ^{cd}	11.40	0.72 ^{ab}	0.008
4 (5)	5.00	7.00	527.4 ^{abc}	11.40	0.75 ^a	0.008
5 (8)	5.00	4.00	545.3 ^a	9.01	0.75 ^a	0.006
6 (5)	5.00	5.50	503.0 ^{abcd}	11.40	0.73 ^{ab}	0.008
7 (5)	6.55	4.44	456.0 ^d	11.40	0.70 ^b	0.008
8 (5)	6.56	6.56	521.4 ^{abc}	11.40	0.75 ^a	0.008
9 (4)	7.20	5.50	516.5 ^{abc}	12.74	0.73 ^{ab}	0.009
Adjusted R ²			0.5202		0.5117	
Predicted R ²			0.3679		0.3520	
<i>p</i> values						
Diet			<.0001		0.0003	
Quadratic source						
Sequential			<0.001		<0.001	
Lack of fit			0.078		0.718	

BWG=body weight gain; G:F=gain-to-feed ratio; SEM=standard error of the mean

The average initial body weight was 301 ± 31.3 g.

¹Number of observations (pens) per treatment after excluding outliers.

Table 3. Bone mineralization of broiler chickens fed diets varying in concentrations of standardized ileal digestible (SID) calcium (Ca) and phosphorous (P) from d 10 to 21.

Run no. (n ¹)	Input variables, g/kg		Output variables							
	SID Ca	SID P	Tibia ash (%)	SEM	Tibia Ca (%)	SEM	Tibia P (%)	SEM	BMD (g/cm ²)	SEM
1 (5)	2.8	5.5	45.26 ^d	0.635	16.40 ^c	0.426	9.27 ^{cd}	0.202	0.105 ^d	0.0025
2 (5)	3.4	4.4	48.20 ^{bc}	0.635	17.95 ^{abc}	0.426	9.98 ^{abc}	0.202	0.114 ^{cd}	0.0025
3 (5)	3.4	6.6	46.97 ^{cd}	0.635	17.47 ^{abc}	0.426	9.62 ^{bcd}	0.202	0.111 ^{cd}	0.0025
4 (5)	5.0	4.0	50.76 ^{ab}	0.635	18.47 ^{ab}	0.426	10.31 ^{ab}	0.202	0.131 ^b	0.0025
5 (8)	5.0	5.5	51.09 ^a	0.505	19.06 ^a	0.336	10.42 ^a	0.169	0.134 ^{ab}	0.0020
6 (5)	5.0	7.0	45.77 ^{cd}	0.635	16.83 ^{bc}	0.426	9.00 ^d	0.202	0.115 ^{cd}	0.0025
7 (5)	6.6	4.4	46.39 ^{cd}	0.635	17.63 ^{abc}	0.426	9.05 ^d	0.202	0.119 ^c	0.0025
8 (5)	6.6	6.6	51.12 ^a	0.635	18.35 ^{ab}	0.426	10.54 ^a	0.202	0.144 ^a	0.0025
9 (5)	7.2	5.5	51.16 ^a	0.635	18.94 ^a	0.426	10.35 ^{ab}	0.202	0.138 ^{ab}	0.0025
Adjusted R ²			0.3803		0.3475		0.5424		0.8302	
Predicted R ²			0.3127		0.1878		0.4304		0.7889	
<i>p</i> values										
Diet			<.0001		0.0001		<.0001		<.0001	
Quadratic source										
Sequential			<0.001		0.006		0.004		<0.001	
Lack of fit			0.003		0.029		0.038		0.672	

SEM=standard error of the mean; BMD= bone mineral density.

¹Number of observations (pens) per treatment

Table 4. The single-objective and multi-objective optimization of performance of broilers fed experimental diets from day 10 to 21¹.

Item	Input variables, g/kg		Predicted output at optimal point	Desirability
	SID Ca	SID P		
BWG	5.13	5.74	Maximum = 545.5 (g)	0.78
G:F	6.41	7.00	Maximum = 0.76 (g/g)	0.78
Tibia ash	7.20	7.00	Maximum = 52.52 (%)	0.97
Tibia Ca	5.75	5.87	Maximum = 19.24 (%)	0.69
Tibia P	7.20	7.00	Maximum = 10.85 (%)	0.90
BMD	7.20	6.96	Maximum = 0.147 (g/cm ²)	0.95
Growth performance*	6.02	6.61	Maximum BWG = 541.1 (g) Maximum G:F = 0.75 (g/g) Maximum ash = 52.20 (%)	0.75
Bone mineralization**	7.13	6.80	Maximum Ca = 18.47 (%) Maximum P = 10.83 (%) Maximum BMD = 0.147 (g/cm ²) Maximum BWG = 537.9 (g) Maximum G:F = 0.76 (g/g) Maximum ash = 51.70 (%)	0.86
Total***	6.50	6.83	Maximum Ca = 18.69 (%) Maximum P = 10.79 (%) Maximum BMD = 0.145 (g/cm ²)	0.78

Ca=calcium; P=phosphorus; SID=standardized ileal digestible; BWG=body weight gain; G:F=gain-to-feed ratio; BMD=bone mineral density.

*Importance in determining SID Ca and P requirements according to Design-Expert software: BWG=5; G:F=5.

**Importance in determining SID Ca and P requirements according to Design-Expert software: Tibia ash=1; Tibia Ca=1; Tibia P=1; BMD=3.

***Importance in determining SID Ca and P requirements according to Design-Expert software: BWG=5; G:F=5; Tibia ash=1; Tibia Ca=1; Tibia P=1; BMD=3.