


Physicochemical properties of crust derived from dry-aged *Holstein* and *Hanwoo* loins

Jeong-Ah Lee and Hack-Youn Kim*

Department of Animal Resources Science, Kongju National University, Yesan 32439, Korea



Received: May 14, 2020
 Revised: Jul 13, 2020
 Accepted: Jul 27, 2020

*Corresponding author

Hack-Youn Kim
 Department of Animal Resources
 Science, Kongju National University,
 Yesan 32439, Korea.
 Tel: +82-41-330-1241
 E-mail: kimhy@kongju.ac.kr

Copyright © 2020 Korean Society of
 Animal Sciences and Technology.
 This is an Open Access article
 distributed under the terms of the
 Creative Commons Attribution
 Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted
 non-commercial use, distribution, and
 reproduction in any medium, provided
 the original work is properly cited.

ORCID

Jeong-Ah Lee
<https://orcid.org/0000-0003-3019-8321>
 Hack-Youn Kim
<https://orcid.org/0000-0001-5303-4595>

Competing interests

No potential conflict of interest relevant
 to this article was reported.

Funding sources

This research was supported by Korea
 Institute of Planning and Evaluation for
 Technology in Food, Agriculture and
 Forestry (IPET), funded by Ministry
 of Agriculture, Food and Rural Affairs
 (MAFRA: 119024-03).

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets
 of this study can be available from the
 corresponding author.

Abstract

This study evaluated the quality characteristics of crust derived from dry-aged *Holstein* and *Hanwoo* loins and their effects on food as additives. With respect to physicochemical properties, we examined the proximate composition, pH value, salinity, color, water and fat absorption, emulsifying capacity, and swelling yield. The protein and ash contents in the *Holstein* crust were significantly higher than those in the *Hanwoo* crust ($p < 0.0001$). The fat content in the *Hanwoo* crust was significantly higher than that in the *Holstein* crust ($p < 0.01$). The salinity, lightness, and yellowness of the *Hanwoo* crust were significantly lower than those of the *Holstein* crust ($p < 0.001$). Furthermore, the pH value and emulsifying capacity of the *Hanwoo* crust were significantly higher than those of the *Holstein* crust ($p < 0.001$). The fat absorption of the *Holstein* crust was significantly higher than that of the *Hanwoo* crust ($p < 0.001$). The swelling yield of the *Holstein* crust was significantly higher than that of the *Hanwoo* crust at pH 3 and 4 ($p < 0.001$), whereas the swelling yield of the *Hanwoo* crust was significantly higher than that of the *Holstein* crust at pH 7 ($p < 0.001$). Principal component analysis of dry-aged *Hanwoo*, *Holstein*, and non-aged *Holstein* showed different flavor patterns for each sample. Finally, the results showed that the crusts derived from dry-aged *Hanwoo* and *Holstein* loins were suitable flavor enhancers.

Keywords: Crust, Dry aging, *Holstein*, *Hanwoo*, Quality characteristics

INTRODUCTION

The aging of meat involves myofibrillar fragmentation of protein, and the methods of aging can be categorized as wet- and dry-aging. During the aging process of the meat, a change in pH in accordance with a corresponding action activates calpain and cathepsin, which are proteolytic enzymes in muscles and increase free amino acids [1–5]. Additionally, triglycerides produce free fatty acids, which enhance tenderness and flavor [6]. Recent studies evaluated that dry-aging, an aging method that involves controlling humidity and wind speed in the air to maximize the flavor and taste of meat, improves and enriches the flavor, taste, and tenderness of meats more than wet-aging. [7–9]. Dashdorj et al. [10] reported that dry-aged meat with a unique flavor was highly preferred by consumers. Choe and Kim [11] reported that this aging technique was being used to add market value to low-grade cattle beef.

Crust is produced by surface hardening during the dry-aging of meat, and thus, the crust has low utility as meat because of its hard texture [10]. As the duration of dry-aging increases, the quantity of

Authors' contributions

Conceptualization: Kim HY.
 Data curation: Kim HY.
 Formal analysis: Lee JA.
 Methodology: Kim HY.
 Software: Lee JA.
 Validation: Kim HY.
 Investigation: Lee JA.
 Writing - original draft: Lee JA.
 Writing - review & editing: Lee JA, Kim HY.

Ethics approval and consent to participate

This article does not require IRB/ACUC approval because there are no human and animal participants.

crust increases, which decreases the yield of dry-aged meat. In addition, dry-aged meats are distributed at high market prices because of the duration of dry-aging [1,2,8,12]. DeGeer et al. [13] reported a 34% trimming loss after beef loins were dry-aged for 28 days. Several studies have been conducted to reduce this trimming loss. Lee et al. [14] reported that the microbial composition of crust changed depending on wind speed and direction in addition to further changes in the physicochemical properties and composition of the flavor compounds. Park et al. [15] improved some functionalities, including the flavor and antioxidant properties, by applying crust with antioxidant and antihypertensive properties to beef patties. However, only a few studies on crust depending on bovine species and breed have been conducted, and product development using crust is limited. In particular, dry-aging has been performed mainly on beef for industrialization; however, crust derived from *Holstein* and *Hanwoo* has not been researched.

Hanwoo is a breed highly accepted by Koreans, unlike *Holstein* and imported meats, and it has a high degree of marbling, excellent flavor, and soft meat quality [16,17]. *Hanwoo* has a high tenderness sufficient for roasting, and it can be distributed as high-quality meat following dry-aging [18,19]. Furthermore, *Hanwoo* has significantly high distribution prices, which means that the development of meat products using crust derived from *Hanwoo* will promote the development of the *Hanwoo* industry [20]. Therefore, in this study we prepared loin (*musculus longissimus dorsi*) crusts from dry-aged *Hanwoo* and *Holstein* to compare their physicochemical quality characteristics as well as processing suitability.

MATERIALS AND METHODS

Materials and preparation of loin crust

In total, 12 pieces of beef loin (*M. longissimus dorsi*) were obtained from 6 carcasses (*Holstein*, Korea quality grade 2; *Hanwoo*, Korea quality grade 2) that were two days postmortem, and were divided into three sections of equal length and width. *Holstein* loin (*M. longissimus dorsi*, I home meat, Seoul, Korea) and *Hanwoo* loin (*M. longissimus dorsi*, Dawoo hanwoo, Chungnam, Korea) were refrigerated for 24 h and used for the tests. Dry aging occurred in a dry-aging refrigerator at 4 °C, 60%–70% relative humidity, and air velocity of 5 ± 3 m/s for 4 weeks. After aging, *Holstein* and *Hanwoo* loins were trimmed off by 30–70 mm from outside. Subsequently, the loin crusts were stored at –18 °C for 24 h (CA-H17DZ, LG, Seoul, Korea) to freeze-dry, and lyophilization was performed at –80 °C for 15 h using a freeze dryer (FDU-1110, Eyela, Tokyo, Japan). The crust was pulverized to a size of 15 mesh using a mixer (MQ5135, Braun, Kronberg im Taunus, Germany) and stored at 4 °C until the analyses.

Proximate composition

In compliance with the AOAC method [21], the protein content was measured via the Kjeldahl method, the fat content was measured via the Soxhlet method, the moisture content was measured via the drying oven method at 105 °C, and the ash content was measured via dry ashing at 550 °C.

Determination of salinity

To measure the salinity, we mixed 1 g of sample with 20 mL of distilled water with an Ultra Turrax homogenizer (HMZ-20DN, Pooglim Tech., Seoul, Korea) at 6,991×g for 1 min. The salinity was recorded using a salinity meter (SB-2000PRO, HMDigital, Seoul, Korea), and the measured value was calculated as a percentage by multiplying dilution ($\times 20$).

Determination of pH values

A mixture of loin crust and distilled water (1:4) was homogenized with an Ultra Turrax homogenizer (HMZ-20DN, Pooglim Tech) at 6,991×g for 1 min. The pH was determined using a pH-meter (Model S220, Mettler-Toledo, Greifensee, Switzerland).

Determination of instrumental color

We used a colorimeter (CR-10, Minolta, Tokyo, Japan) to measure the lightness (CIE L*), redness (CIE a*), and yellowness (CIE b*) of the samples. The colorimeter was calibrated with a white standard plate L* = 97.83, a* = -0.43, and b* = +1.98.

Determination of water absorption

The water absorption was measured via the method of Lin et al. [22], with some modifications. First, 1 g of sample and 10 mL distilled water were mixed using a vortex mixer (SVM-10, SciLab, Seoul, Korea) for 2 min, and the mixture was centrifuged at 15°C, 983×g for 20 min (Supra R22, Hanil, Gimpo, Korea). The supernatant was carefully decanted, and the weight of the precipitate was measured before drying (W₁). Then, the precipitate was dried for 24 h at 105°C in a drying oven (C-F03, Cheil, Gyeonggi-do, Korea), and the weight was measured after drying (W₂). The water absorption was calculated using the following equation:

$$\text{Water absorption (\%)} = \frac{W_1 - W_2}{W_1} \times 100$$

Determination of fat absorption

The fat absorption was determined via the method of Lin et al. [22], with some modifications. Based on the protein content in the sample, the total protein amount was calculated to be 1 g and the sample was used (W₁). The sample was mixed with 10 mL of soybean oil (V₁) using a vortex mixer (SVM-10, SciLab) for 30 s, and the mixture rested at room temperature (25°C) for 30 min. Then, the mixture was centrifuged at 25°C, 983 ×g for 20 min (Supra R22, Hanil), and the volume of the supernatant was measured (V₂). The fat absorption was calculated using the following equation.

$$\text{Fat absorption (\%)} = \frac{V_1 - V_2}{W_1} \times 100$$

Determination of emulsifying capacity

The emulsifying capacity was determined using the method of Yasumatsu et al. [23]. First, 7 g of the sample, 50 mL of distilled water, and 100 mL of soybean oil were mixed using the Ultra Turrax homogenizer (HMZ-20DN, Pooglin Tech.) at 983×g for 1 min. The prepared emulsion was allowed to stand in a graduated cylinder at room temperature (25°C) for approximately 1 h or more. The emulsifying capacity was measured based on the total amount of solution (B) and amount of mixture (A) using the following equation.

$$\text{Emulsifying capacity (\%)} = \frac{A}{B} \times 100$$

Determination of swelling yield

First, 10 g of the sample was mixed with 100 mL of Tris-HCl solution at pH 3, 4, and 7 using

a vortex mixer (SVM-10, SciLab) for 1 min. Then, the prepared mixture was allowed to stand at room temperature (25 °C) for 1 h. The supernatant was carefully decanted, and the weight of the precipitate was measured. The swelling yield was calculated using the following equation.

$$\text{Swelling yield (\%)} = \frac{\text{After swelling (g)}}{\text{Before swelling (g)}} \times 100$$

Electronic nose analysis

The loin crusts were individually placed in a 20 mL vial on a sample holder heated at 80 °C for 20 min. The headspace volatile compounds were injected into a gas chromatography-type electronic nose (HERACLES-2-E-NOSE, alpha-mos, Toulouse, France) equipped with dual columns (MXT-5 and 1701, Restek, Bellefonte, PA, USA) (length 10 m, inner diameter 180 µm, MXT-5: non-polarity, MXT-1701: slight polarity). The analysis conditions were set as follows: injection time of 20 min, volume of 2 mL, rate of 250 µL/s, temperature of 200 °C, and detector temperature of 260 °C. The principal component analysis (PCA) was integrated using the Alpha Soft program (Alphasoft, Alpha MOS, Toulouse, France).

Statistical analysis

The experimental results were assessed after a minimum of three repeated trials. Statistical analyses were performed using the General Linear Model procedure of the SAS program (2015, SAS Software for Window, Version 9.3, SAS Institute, Cary, NC, USA). *T*-tests were performed to compare each average of the treatments, and the significance was expressed as $p < 0.05$, $p < 0.01$, $p < 0.001$.

RESULTS AND DISCUSSION

Proximate composition, pH, salinity, and color

Table 1 shows the general components of the loin crusts of *Holstein* and *Hanwoo*. The moisture contents in *Hanwoo* and *Holstein* crusts showed no significant difference. The fat content in the *Hanwoo* crust was significantly higher than that in the *Holstein* crust ($p < 0.01$), whereas the protein and ash contents in the *Holstein* crust were significantly higher than those in the *Hanwoo* crust ($p < 0.001$). The moisture content in the crust during drying showed no significant difference because most of the free water was dried. Choe and Kim [11] reported that the 2nd grade *Holstein* loin contained 19.95% protein, 1.10% ash, and 5.97% fat, whereas the 2nd class *Hanwoo* loin contained 17% protein, 0.51% ash, and 23.71% fat [3]. The results suggested that the crust after dry-aging showed a similar tendency because the fat content in the *Holstein* loin was lower but the protein

Table 1. Proximate composition of crust derived from dry-aged *Holstein* and *Hanwoo* loin

Traits	Dry aging crust		SEM ¹⁾	Statistical analysis	
	<i>Holstein</i>	<i>Hanwoo</i>		<i>t</i> -value	<i>p</i> -value
Moisture content (%)	1.15	1.20	6.92	-0.25 ^{ns}	0.8123
Protein content (%)	62.82	44.06	3.99	12.05 ^{***}	0.0003
Fat content (%)	23.90	49.33	12.09	-6.34 ^{**}	0.0032
Ash content (%)	3.90	1.95	0.75	15.45 ^{***}	0.0001

All values are mean.

¹⁾n = 2.

^{**} $p < 0.01$, ^{***} $p < 0.001$.

ns, non-significant.

and ash contents were higher than those in *Hanwoo* loin of the same grade.

Table 2 shows the measurements of pH, salinity, and instrumental color of the *Holstein* and *Hanwoo* crusts. The pH of the *Hanwoo* crust was significantly higher than that of the *Holstein* crust ($p < 0.001$). Li et al. [7] reported that intramuscular fat content and pH showed a positive correlation in the same bovine species, which suggested that the pH of the *Hanwoo* crust with a higher fat content was higher than that of *Holstein*. The salinity of the *Holstein* crust was significantly higher than that of the *Hanwoo* crust ($p < 0.001$), this was related to free amino acids because they affect the salinity measurements. These results suggested that numerous free amino acids were released from *Holstein* with a higher protein content than that of *Hanwoo* [24]. Free amino acids hydrolyzed from proteins contain taste components such as inosine monophosphate (IMP) and guanosine-5'-monophosphate [25] and that the total amount of free amino acids increases as the aging is prolonged [26]. Among inorganic ionic components, Na, K, and P significantly enhance the umami taste along with free amino acids and IMPs [27]. Thus, free amino acids are more accumulated in the *Holstein* crust with high protein content than in the *Hanwoo* crust, and the salinity of the *Holstein* crust is high because it is typically measured based on the content of the total inorganic substance [28]. In addition, both *Hanwoo* and *Holstein* crust have a salinity of 0.8%–1.2%, which will not have a significant impact on salinity as food additives. With respect to the instrumental color of the *Hanwoo* and *Holstein* crusts, the lightness and yellowness of *Holstein* were significantly higher than those of *Hanwoo* ($p < 0.001$), and there was no significant difference in redness between the two treatment groups. Lee et al. [29] reported that the lightness, redness, and yellowness of *Hanwoo* were significantly low during the analysis of the physicochemical quality characteristics of *Holstein* and *Hanwoo*. Difference color between *Hanwoo* and *Holstein* is related to the moisture contents. In general, it is known to *Holstein* beef loin is higher moisture content than *Hanwoo* beef loin [11]. And high moisture content of beef loin related to the increase the lightness, redness, and yellowness [30]. Therefore, the moisture content of the *Holstein* crust was higher than *Hanwoo* crust, it was related to the higher lightness and yellowness than *Hanwoo* crust. The lightness and yellowness of the *Hanwoo* crust were significantly lower in this study, and no significant difference was found in the increase in redness due to dried moisture.

Water absorption, fat absorption, emulsifying capacity, and swelling yield

Table 3 shows the analytical results of water absorption, fat absorption, emulsifying capacity, and swelling yield of the *Hanwoo* and *Holstein* crusts. There was no significant difference in water absorption between the *Hanwoo* and *Holstein* treatment groups, whereas the fat absorption of the *Hol-*

Table 2. pH, salinity and color of crust derived from dry-aged *Holstein* and *Hanwoo* loin

Traits	Dry aging crust		SEM ¹⁾	Statistical analysis	
	<i>Holstein</i>	<i>Hanwoo</i>		<i>t</i> -value	<i>p</i> -value
pH	5.44	5.67	0.08	-22.30 ^{***}	<.0001
Salinity (%)	1.20	0.80	0.15	lnfty ^{***}	<.0001
Color					
CIE L*	54.26	37.82	6.13	77.50 ^{***}	<.0001
CIE a*	15.90	12.87	2.00	1.63 ^{ns}	0.1534
CIE b*	12.38	2.08	3.86	26.70 ^{***}	<.0001

All values are mean.

¹⁾n = 2.

^{***} $p < 0.001$.

ns, non-significant.

Table 3. Water absorption, fat absorption, emulsifying capacity and swelling yield of crust derived from dry-aged *Holstein* and *Hanwoo* loin

Traits	Dry aging crust		SEM ¹⁾	Statistical analysis	
	<i>Holstein</i>	<i>Hanwoo</i>		t-value	p-value
Water absorption (%)	2.96	2.26	0.39	1.83 ^{ns}	0.1420
Fat absorption (%)	2.85	0.62	0.86	40.49 ^{***}	<.0001
Emulsifying capacity (%)	17.20	54.10	14.46	-12.72 ^{***}	0.0002
Swelling yield					
pH 3	306.20	168.31	53.61	22.16 ^{***}	<.0001
pH 4	317.43	164.78	59.58	15.97 ^{***}	<.0001
pH 7	120.57	168.55	18.86	-11.55 ^{***}	0.0003

All values are mean.

¹⁾n = 2.

^{***}p < 0.001.

ns, non-significant.

stein crust was significantly higher than that of the *Hanwoo* crust ($p < 0.001$). Lee et al. [31] showed that the fat absorption of rice protein concentrate with low fat content was high when testing the dry frozen rice protein concentrate and isolated soy protein, which suggested that the fat absorption decreased as the fat content increased. Thus, it would be efficient to apply the *Holstein* crust with the high fat absorption capacity to products with high fat content and the *Hanwoo* crust with low fat absorption to low-fat food ingredients.

The emulsifying capacity of the *Hanwoo* crust was significantly higher than that of the *Holstein* crust ($p < 0.001$). Beef fat can be divided into neutral lipids and phospholipids. Phospholipids possess an emulsifying capacity to combine water and oil [32,33]. *Hanwoo* has a higher emulsifying capacity than *Holstein* because of its higher content of palmitic acid and stearic acid in phosphoglycerides and sphingolipids [34]. Thus, the *Hanwoo* crust can be utilized as a food additive for enhancing the binding power between moisture and oil [35].

The measurement of the swelling yield based on the pH must be preceded to increase the applicability of the powder crust that enhances flavor [36]. The larger the swelling yield of the crust, the less physicochemical change in the food processing, such as sauces, and the less drying loss [37,38]. The swelling yields of the *Hanwoo* and *Holstein* crusts were measured at pH values of 3, 4, and 7 for sauces. At pH 3 and 4, the *Holstein* crust showed significantly higher swelling yield than the *Hanwoo* crust ($p < 0.001$), whereas at pH 7, the *Hanwoo* crust showed a significantly higher swelling yield than the *Holstein* crust ($p < 0.001$). The swelling yield is a factor that highly affects the yield of liquid or semi-solid products [39], and it would be ideal to prepare food at a high pH corresponding to high swelling yield. The *Holstein* crust showed excellent swelling yield under pH 3 and 4; thus, it can be effectively applied to the sauce and vinegar product groups [40,41]. Conversely, the *Hanwoo* crust showed a high swelling yield under pH 7 (neutral); thus, it can be suitably applied to sauces and processed meat products [42].

Principal component analysis in electric nose

An electric nose analysis can be widely used to evaluate food quality and aging by classifying the characteristics that determine the overall flavor of food [43]. Fig. 1 shows the PCA for the *Holstein* and *Hanwoo* crusts using the electric nose technique. According to the PCA analysis, the contribution rate of the first principal component (PC1) value was 96.484%, and that of the second principal component (PC2) value was 3.441%. Because the contribution rate of PC1 was 90% or higher, the flavor was more discriminated by PC1 than PC2 [43]. The PC1 range of the non-aged *Holstein*,




Fig. 1. Principal component analysis of crust derived from dry-aged *Holstein* and *Hanwoo* loin.

Hanwoo, and *Holstein* crusts were approximately -75,000, -20,000, 95,000, respectively. The PC2 range of the *Hanwoo* crust was approximately 19,500, whereas those of the non-aged *Holstein* and *Holstein* crusts were between -5,000 and -15,000. Thus, in terms of PC1, which greatly contributed to flavor discrimination, the discrimination in flavor could be determined based on the difference between the treatment groups. The difference in the flavor of the *Hanwoo* and *Holstein* crusts depends on the breed in the same bovine species and site. Kang et al. [44] reported similar results on the difference between breeds at the same site of pork through electric nose analysis of flavor because various compositions of fatty acids and amino acids that depend on breed resulted in various aromatic patterns.

CONCLUSION

The present study analyzed the physicochemical properties of crust derived from dry-aged *Holstein* and *Hanwoo* loins as flavor enhancers. The emulsifying capacity of the *Hanwoo* crust was significantly higher than that of the *Holstein* crust. The fat absorption of the *Holstein* crust was higher than that of the *Hanwoo* crust. The dry-aged *Hanwoo* and *Holstein* samples showed notably different flavor patterns via PCA. Therefore, the addition of crust derived from dry-aged *Hanwoo* and *Holstein* loins would lead to enhanced flavor in food. And choose the type of beef crust consider to purpose of food flavor and characteristics.

REFERENCES

1. Campbell RE, Hunt MC, Levis P, Chambers E IV. Dry-aging effects on palatability of beef longissimus muscle. *J Food Sci.* 2001;66:196-9. <https://doi.org/10.1111/j.1365-2621.2001.tb11315.x>
2. Smith RD, Nicholson KL, Nicholson JDW, Harris KB, Miller RK, Griffin DB, et al. Dry versus wet aging of beef: retail cutting yields and consumer palatability evaluations of steaks from US Choice and US Select short loins. *Meat Sci.* 2008;79:631-9. <https://doi.org/10.1016/j.meatsci.2007.10.028>
3. Lee CW, Lee SH, Min Y, Lee S, Jo C, Jung S. Quality improvement of strip loin from Hanwoo with low quality grade by dry aging. *Korean J Food Nutr.* 2015;28:415-21. <https://doi.org/10.1007/s00003-015-0000-0>

- org/10.9799/ksfan.2015.28.3.415
4. Kim JH, Cho SH, Seong PN, Hah KH, Kim HK, Park BY, et al. Effect of ageing temperature and time on the meat quality of longissimus muscle from Hanwoo steer. *Korean J Food Sci Anim Resour.* 2007;27:171-8. <https://doi.org/10.5851/kosfa.2007.27.2.171>
 5. Laville E, Sayd T, Moraël M, Blinet S, Chambom C, Lepetit J, et al. Proteome changes during meat aging in tough and tender beef suggest the importance of apoptosis and protein solubility for beef aging and tenderization. *J Agric Food Chem.* 2009;57:10755-4. <https://doi.org/10.1021/jf901949r>
 6. Koutsidis G, Elmore JS, Oruna-Concha MJ, Campo MM, Wood JD, Mottram DS. Water-soluble precursors of beef flavour. Part II: effect of post-mortem conditioning. *Meat Sci.* 2008;79:270-7. <https://doi.org/10.1016/j.meatsci.2007.09.010>
 7. Li YX, Cabling MM, Kang HS, Kim TS, Yeom SC, Sohn YG, et al. Comparison and correlation analysis of different swine breeds meat quality. *Asian-Australas J Anim Sci.* 2013;26:905-10. <https://doi.org/10.5713/ajas.2012.12622>
 8. Laster MA, Smith RD, Nicholson KL, Nicholson JDW, Miller RK, Griffin DB, et al. Dry versus wet aging of beef: retail cutting yields and consumer sensory attribute evaluations of steaks from ribeyes, strip loins, and top sirloins from two quality grade groups. *Meat Sci.* 2008;80:795-804. <https://doi.org/10.1016/j.meatsci.2008.03.024>
 9. Smith AM, Harris KB, Griffin DB, Miller RK, Kerth CR, Savell JW. Retail yields and palatability evaluations of individual muscles from wet-aged and dry-aged beef ribeyes and top sirloin butts that were merchandised innovatively. *Meat Sci.* 2014;97:21-6. <https://doi.org/10.1016/j.meatsci.2013.12.013>
 10. Dashdorj D, Tripathi VK, Cho S, Kim Y, Hwang I. Dry aging of beef; review. *J Anim Sci Technol.* 2016;58:20. <https://doi.org/10.1186/s40781-016-0101-9>
 11. Choe JH, Kim HY. Effects of dry aging on physicochemical properties of beef cattle loins. *Korean J Food Sci Technol.* 2017;49:158-61. <https://doi.org/10.9721/KJFST.2017.49.2.158>
 12. Parrish FC Jr, Boles JA, Rust RE, Olson DG. Dry and wet aging effects on palatability attributes of beef loin and rib steaks from three quality grades. *J Food Sci.* 1991;56:601-3. <https://doi.org/10.1111/j.1365-2621.1991.tb05338.x>
 13. DeGeer SL, Hunt MC, Bratcher CL, Crozier-Dodson BA, Johnson DE, Stika JF. Effects of dry aging of bone-in and boneless strip loins using two aging processes for two aging times. *Meat Sci.* 2009;83:768-74. <https://doi.org/10.1016/j.meatsci.2009.08.017>
 14. Lee HJ, Yoon JW, Kim M, Oh H, Yoon Y, Jo C. Changes in microbial composition on the crust by different air flow velocities and their effect on sensory properties of dry-aged beef. *Meat Sci.* 2019;153:152-8. <https://doi.org/10.1016/j.meatsci.2019.03.019>
 15. Park B, Yong HI, Choe J, Jo C. Utilization of the crust from dry-aged beef to enhance flavor of beef patties. *Korean J Food Sci Anim Resour.* 2018;38:1019-28. <https://doi.org/10.5851/kosfa.2018.e35>
 16. Oh DY, Yeo JS, Lee JY. Major SNP identification for oleic acid and marbling score which are associated with Korean cattle. *J Korean Data Inf Sci Soc* 2014;25:1011-24. <https://doi.org/10.7465/jkdi.2014.25.5.1011>
 17. Ha Y, Hwang I, Choi J, Kim Y, Seol KH, Seo HW, et al. Comparison of meat quality traits for M. Longissimus thoracis, M. Glutaeusmedus, M. Semimembranosus from low-grade Hanwoo cow beef by dry aging and wet aging conditions. *Ann Anim Resour Sci.* 2019;30:121-32. <https://doi.org/10.12718/AARS.2019.30.3.121>
 18. Moon JH, Sung M, Kim JH, Kim BS, Kim Y. Quality factors of freshness and palatability of Hanwoo from their physicochemical and sensorial properties. *Korean J Food Sci Anim Resour.*

- 2013;33:796-805. <https://doi.org/10.5851/kosfa.2013.33.6.796>
19. Calvo JH, Iguácel LP, Kirinus JK, Serrano M, Ripoll G, Casasús I, et al. A new single nucleotide polymorphism in the calpastatin (CAST) gene associated with beef tenderness. *Meat Sci.* 2014;96:775-82. <https://doi.org/10.1016/j.meatsci.2013.10.003>
 20. Joo ST, Hwang YH, Frank D. Characteristics of Hanwoo cattle and health implications of consuming highly marbled Hanwoo beef. *Meat Sci.* 2017;132:45-51. <https://doi.org/10.1016/j.meatsci.2017.04.262>
 21. AOAC [Association of Official Analytical Chemists]. Official methods of analysis of AOAC International. 19th ed. Gathersburg, MD: AOAC International; 2012. p. 931.
 22. Lin MJY, Humbert ES, Sosulki FW. Certain functional properties of sunflower meal products. *J Food Sci.* 1974;39:368-70. <https://doi.org/10.1111/j.1365-2621.1974.tb02896.x>
 23. Yasumatsu K, Sawada K, Moritaka S, Misaki M, Toda J, Wada T, et al. Whipping and emulsifying properties of soybean products. *Agric Biol Chem.* 1972;36:719-27. <https://doi.org/10.1080/00021369.1972.10860321>
 24. Armstrong AD, Stave U. A study of plasma free amino acid levels. III. Variations during growth and aging. *Metaboilsm.* 1973;22:571-8. [https://doi.org/10.1016/0026-0495\(73\)90070-X](https://doi.org/10.1016/0026-0495(73)90070-X)
 25. Kim YHB, Kempa R, Samuelsson LM. Effects of dry-aging on meat quality attributes and metabolite profiles of beef loins. *Meat Sci.* 2016;111:168-76. <https://doi.org/10.1016/j.meatsci.2015.09.008>
 26. Parrish FCJr, Goll DE, Newcomb II WJ, Lumen BO, Chaudhry HM, Kline EA. Molecular properties of post-mortem muscle 7. changes in nonprotein nitrogen and free amino acids of bovine muscle. *J Food Sci.* 1969;34:196-202. <https://doi.org/10.1111/j.1365-2621.1969.tb00918.x>
 27. Hayashi T, Yamaguchi K, Konosu S. Sensory analysis of taste-active components in the extract of boiled snow crab meat. *J Food Sci.* 1981;46:479-83. <https://doi.org/10.1111/j.1365-2621.1981.tb04890.x>
 28. Cheon DH, Lee KH. Development of jeyuk-gui (spicy broiled pork) souce amended with concentrate of jujube flesh remaining around seeds. *J East Asian Soc Diet Life.* 2017;27:591-9. <https://doi.org/10.17495/easdl.2017.12.27.6.591>
 29. Lee SJ, Ha ES, Hu WS, Sung NJ. Comparison of quality characteristics and antioxidant activity in loin meat of imported beefs and organic hanwoo. *J Agric Life Sci.* 2016;50:149-63. <https://doi.org/10.14397/jals.2016.50.3.149>
 30. Kim YHB, Robert K, Linda MS. Effect of dry-aging on meat quality attributes and metabolite profiles of beef loins. *Meat Sci.* 2016;111:168-76. <https://doi.org/10.1016/j.meatsci.2015.09.008>
 31. Lee ES, Kim KJ, Kim JH, Hong ST. A study on the development of high functional food protein ingredient from rice bran. *J Agric Sci.* 2010;37:61-8. <https://doi.org/10.7744/cnujas.2010.37.1.061>
 32. Warren HE, Scollan ND, Enser M, Hughes SI, Richardson RI, Wood JD. Effects of breed and a concentrate or grass silage diet on beef quality in cattle of 3ages. I: animal performance, carcass quality and muscle fatty acid composition. *Meat Sci.* 2008;78:256-69. <https://doi.org/10.1016/j.meatsci.2007.06.008>
 33. Kim SY, Jeon SS, Jun ST. Effect of lotus root powder on the baking quality of white bread. *Korean J Food Cookery Sci.* 2002;18:413-25.
 34. Hur SJ, Park GB, Joo ST. A comparison of the meat qualities from the hanwoo (Korean native cattle) and Holstein steer. *Food Bioproc Tech.* 2008;1:196-200. <https://doi.org/10.1007/s11947-008-0061-2>

35. Duckett SK, Wagner DG. Effect of cooking on the fatty acid composition of beef intramuscular lipid. *J Food Compos Anal.* 1998;11:357-62. <https://doi.org/10.1006/jfca.1998.0600>
36. Alam F, Hasnain A. Studies on swelling and solubility of modified starch from Taro (*Colocasia esculenta*): effect of pH and Temperature. *Agric conspec sci.* 2009;74:45-50.
37. Femenia A, Garcia-Pascual P, Simal S, Rosselló C. Effects of heat treatment and dehydration on bioactive polysaccharide acemannan and cell wall polymers from *Aloe barbadensis* Miller. *Carbohydr Polym.* 2003;51:397-405. [https://doi.org/10.1016/S0144-8617\(02\)00209-6](https://doi.org/10.1016/S0144-8617(02)00209-6)
38. Simal S, Femenia A, Llull P, Rosello C. Dehydration of aloe vera: simulation of drying curves and evaluation of functional properties. *J Food Eng.* 2000;43:109-14. [https://doi.org/10.1016/S0260-8774\(99\)00139-9](https://doi.org/10.1016/S0260-8774(99)00139-9)
39. Johnston-Banks FA. Gelatine. In: Harris P, editor. *Food gels*. London: Elsevier Applied Science; 1990. p. 234-7.
40. Cho KH, Kang SA. Effects quality characteristics and development of global sauce using traditional gochujang. *J Korea Acad Ind Coop Soc.* 2015;16:8089-95. <https://doi.org/10.5762/KAIS.2015.16.11.8089>
41. Kim GR, Yoon SR, Lee JH, Yeo SH, Jeong YJ, Yoon KY, et al. Physicochemical properties of and volatile components in commercial fruit vinegars. *Korean J Food Preserv.* 2010;17:616-24.
42. Hwang SH. Effect of high pressure processing on freshness of meat products. *J Food Hyg Saf.* 2018;33:272-9. <https://doi.org/10.13103/JFHS.2018.33.4.272>
43. Shin JA, Choi SW, Lee KT. Prediction of Kimchi aging using electronic nose system. *Korean J Food Preserv.* 2005;12:613-6.
44. Kang SM, Kang CG, Lee SG. Analysis of differences in fat content, fatty acid composition and electronic nose by pork with varieties. In: *Proceedings of the Korean Society for Food Science of Animal Resources Conference*; 2006; Seoul. p. 107-11.