

# Interactions between *Entodinium caudatum* and an amino acid-fermenting bacterial consortium: fermentation characteristics and protozoal population *in vitro*

Tansol Park<sup>1,2</sup> and Zhongtang Yu<sup>2\*</sup>

<sup>1</sup>Department of Animal Science and Technology, Chung-Ang University, Anseong 17546, Korea

<sup>2</sup>Department of Animal Sciences, The Ohio State University, Columbus, OH 43210, USA



Received: Oct 17, 2022  
Revised: Nov 21, 2022  
Accepted: Nov 21, 2022

## \*Corresponding author

Zhongtang Yu  
Department of Animal Sciences, The Ohio State University, Columbus, OH 43210, USA.  
Tel: +1-614-292-3057  
E-mail: yu.226@osu.edu

Copyright © 2023 Korean Society of Animal Sciences and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ORCID

Tansol Park  
<https://orcid.org/0000-0002-4480-4524>  
Zhongtang Yu  
<https://orcid.org/0000-0002-6165-8522>

## Competing interests

No potential conflict of interest relevant to this article was reported.

## Funding sources

This research is based on work that is partially supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture, under award numbers 2021-67015-33393 and supported by the Chung-Ang University Research Grants in 2021.

## Abstract

Ruminal protozoa, especially entodiniomorphs, engulf other members of the rumen microbiome in large numbers; and they release oligopeptides and amino acids, which can be fermented to ammonia and volatile fatty acids (VFAs) by amino acid-fermenting bacteria (AAFB). Studies using defaunated (protozoa-free) sheep have demonstrated that ruminal protozoa considerably increase intraruminal nitrogen recycling but decrease nitrogen utilization efficiency in ruminants. However, direct interactions between ruminal protozoa and AAFB have not been demonstrated because of their inability to establish axenic cultures of any ruminal protozoan. Thus, this study was performed to evaluate the interaction between *Entodinium caudatum*, which is the most predominant rumen ciliate species, and an AAFB consortium in terms of feed degradation and ammonia production along with the microbial population shift of select bacterial species (*Prevotella ruminicola*, *Clostridium aminophilum*, and *Peptostreptococcus anaerobius*). From an *Ent. caudatum* culture that had been maintained by daily feeding and transfers every 3 or 4 days, the bacteria and methanogens loosely associated with *Ent. caudatum* cells were removed by filtration and washing. An AAFB consortium was established by repeated transfers and enrichment with casamino acids as the sole substrate. The cultures of *Ent. caudatum* alone (Ec) and AAFB alone (AAFB) and the co-culture of *Ent. caudatum* and AAFB (Ec + AAFB) were set up in three replicates and incubated at 39°C for 72 h. The digestibility of dry matter (DM) and fiber (NDF), VFA profiles, ammonia concentrations, pH, and microscopic counts of *Ent. caudatum* were compared among the three cultures. The co-culture of AAFB and *Ent. caudatum* enhanced DM degradation, VFA production, and *Ent. caudatum* cell counts; conversely, it decreased acetate: propionate ratio although the total bacterial abundance was similar between Ec and the Ec + AAFB co-culture after 24 h incubation. The ammonia production and relative abundance of *C. aminophilum* and *P. anaerobius* did not differ between AAFB alone and the Ec + AAFB co-culture. Our results indicate that *Ent. caudatum* and AAFB could have a mutualistic interaction that benefited each other, but their interactions were complex and might not increase ammoniogenesis. Further research

#### Acknowledgements

Not applicable.

#### Availability of data and material

The datasets of this study can be available from the corresponding author upon reasonable request.

#### Authors' contributions

Conceptualization: Park T, Yu Z.  
Data curation: Park T.  
Formal analysis: Park T.  
Methodology: Park T, Yu Z.  
Software: Park T.  
Validation: Park T, Yu Z.  
Investigation: Park T, Yu Z.  
Writing - original draft: Park T.  
Writing - review & editing: Park T, Yu Z.

#### Ethics approval and consent to participate

This article does not require IRB/ACUC approval because there are no human and animal participants.

should examine how such interactions affect the population dynamics of AAFB.

**Keywords:** Amino acid-fermenting bacteria, Co-culture, *Entodinium caudatum*, Intraruminal nitrogen recycling, Mutualistic interaction

## INTRODUCTION

Ruminant animals depend on a diverse microbial assembly, which consists of bacteria, archaea, protozoa, and fungi, in their rumen for their survival, growth, and production of animal proteins (beef, lamb, milk, and wool). Collectively, digestive and fermentative processes convert dietary carbohydrates, primarily starch and cellulose, and dietary nitrogen, primarily plant protein nitrogen, into carbon and nitrogen sources that ruminants can utilize. However, the utilization efficiency of dietary nitrogen in ruminants is low, which is only approximately 25% [1,2]. This low nitrogen utilization efficiency not only increases the production cost but also creates a major environmental problem. Indeed, about 70% of  $\text{NH}_3$  and 30% of  $\text{N}_2\text{O}$  released into the environment by anthropogenic activities are estimated to stem from livestock husbandry [3]. The ruminal microbiome participates and greatly affects the nitrogen utilization efficiency in ruminants. Conceptually, two metabolic processes can lead to low nitrogen utilization efficiency: microbial protein proteolysis and amino acid deamination. In the rumen, approximately 70% of dietary nitrogen (primarily as protein, often referred to as rumen degradable protein (RDP) is hydrolyzed to oligopeptides and free amino acids, which can be fermented to short-chain fatty acid (SCFA) and ammonia. Some of these nitrogen forms are used as nitrogen sources by ruminal microbes, primarily bacteria, to synthesize cellular proteins, which are the main direct nitrogen source of host animals [4–6]. However, a large portion of microbial cells (about 24% of the total ruminal bacteria daily) are engulfed by ruminal protozoa [7]; furthermore, approximately 50% of the engulfed bacterial protein is hydrolyzed by protozoa and discharged as oligopeptides and free amino acids [8], thereby promoting protein nitrogen recycling in the rumen. A significant portion of oligopeptides and free amino acids are fermented by amino acid-fermenting bacteria (AAFB) to SCFA and ammonia. Thus, ruminal protozoa can decrease the ruminal outflow of microbial protein, which is the main protein source of host animals, to the small intestines. They also increase the availability of substrates for AAFB and the production of ammonia, which is absorbed and converted into urea in the liver and excreted. Ruminal protozoa and AAFB have been explored extensively to understand their roles in nitrogen utilization efficiency in ruminant animals, but their interactions remain poorly understood.

Ruminal protozoa have been considered a nonvital group of microbes for host animals although they contribute to organic matter digestion and homeostasis of the rumen environment [9]. Although their peptidase and deaminase activities remain to be determined and are probably variable among different protozoal species [10,11], all ruminal protozoa engulf and digest cells of ruminal microbes, even small ruminal protozoa; subsequently, they degrade the microbial protein into oligopeptides and amino acids, producing substrates for AAFB [12]. Although AAFB can utilize carbohydrates, they can use amino acids as their sole energy and carbon sources [13]. Unlike the predominant proteolytic bacteria that have limited deamination activity, AAFB, especially hyper-ammonia-producing bacteria (HAB), including *Clostridium aminophilum*, *Clostridium sticklandii*, and *Peptostreptococcus anaerobius*, have high deamination activities [13–15]. Numerous studies have been conducted to understand the roles of ruminal protozoa and AAFB in ruminal protein metabolism and decrease ruminal protozoa (primarily by defaunation) and AAFB by using plant extracts [15–18]. However, effective and practical approaches are yet to be developed to

decrease intraruminal protein turnover and improve nitrogen utilization efficiency.

From an ecological perspective, ruminal protozoa and AAFB can form two relationships: prey-predator relationship and mutualism. In the former relationship, ruminal protozoa benefit from preying on AAFB, but in the latter relationship, ruminal protozoa provide substrates to AAFB. We hypothesized that a better understanding of the interactions between ruminal protozoa and AAFB could help unravel the roles of ruminal protozoa and AAFB in intraruminal protein recycling and improve nitrogen utilization efficiency in ruminants. This study aimed to evaluate the interaction between *Entodinium caudatum*, the most predominant rumen protozoal species, and AAFB in terms of feed degradation, ammoniogenesis, and dynamics of microbial populations.

## MATERIALS AND METHODS

### *Entodinium caudatum* monoculture and amino acid-fermenting bacteria consortium

An *Ent. caudatum* monoculture established from a single cell isolated from the rumen of gerenuk [19] was maintained by daily feeding of protozoal feed containing wheat grain, alfalfa, and grass and regular transfers into a fresh SP medium [20].

An AAFB consortium was established previously in our laboratory by using enrichment on casamino acids as the sole substrate [21]. Briefly, rumen fluid was collected from two rumen-fistulated Jersey dairy cows, mixed, and inoculated into a mineral medium [22] containing yeast extract (0.5 g/L) as growth factors and casamino acids (30 g/L) as the sole substrate. The culture was incubated anaerobically at 39°C and transferred every 24 h until the ammonia concentration in the culture stabilized. *Proteus mirabilis*, *Bacillus* spp., *Fusobacterium ulcerans*, *C. aminophilum*, and *P. anaerobius* with varying amino acid fermentation activities were identified in this AAFB consortium through 16S rRNA gene sequencing.

### Co-culture experiment

The monoculture of *Ent. caudatum* and the AAFB enrichment culture were incubated individually or as a co-culture to explore the interaction between ruminal protozoa and AAFB (Table 1). The *Ent. caudatum* monoculture was washed using three filter membranes with decreasing pore sizes (50, 25, and 10 mm; Sefar Filtration, New York, USA) to remove most prokaryotes present in the *Ent. caudatum* monoculture. The washed *Ent. caudatum* cells retained on the 10 mm filter membrane were then collected into the simplex buffer (modified from Williams and Coleman, 1992 [12]) and used as the *Ent. caudatum* inoculum. The AAFB culture was centrifuged at 21,000×g for 7 min. The pelleted cells were washed with the simplex buffer thrice followed by centrifugation and resuspended in an SP medium [20] as the AAFB inoculum. The cell density of the AAFB suspension was estimated based on its optical density [23]. Approximately  $8.8 \times 10^7$

**Table 1.** Experimental design of the co-culturing experiment

	Ec	AAFB	Ec + AAFB
Feed (g/culture)	0.1	0.1	0.1
SP medium (mL)	8	7	7
Washed <i>Ent. caudatum</i> monoculture (mL)	2	0	2
AAFB suspension (in SP, mL)	0	1	1
Simplex buffer (mL)	0	2	0
Total culture volume (mL)	10	10	10

Ec, *Entodinium caudatum* alone; AAFB, amino acid-fermenting bacterial consortium; Ec + AAFB, co-culture of Ec and AAFB.

AAFB cells/mL were inoculated into the AAFB culture and the Ec + AAFB co-culture. Both the medium and the buffer were made anaerobically by continuous sparging with O<sub>2</sub>-free CO<sub>2</sub> gas. The medium conditions were similar among the three treatments except for the microbial inocula and a higher ammonia concentration in AAFB (data not shown).

### Protozoal counts

Protozoal cells in the two *Ent. caudatum*-containing cultures (i.e., Ec and Ec + AAFB) were fixed and counted microscopically every 24 h of the incubation as described previously [20]. Briefly, 0.5 mL of each culture was fixed in 50% formalin solution and then mixed with 30% glycerol to dilute the culture and prevent the rapid settlement of protozoal cells during the procedures [24]. The protozoal cells were stained with brilliant green (10 mL of dye added to 1.5 mL of fixed protozoal cell suspension) to facilitate microscopic counting. A 1-mL aliquot of each stained protozoal sample was added to a Sedgewick Rafter counting chamber (no. 9851 C20, Thomas Scientific, Swedesboro, NJ, USA), and the cells within 50 different grids were counted twice and averaged.

### Fermentation characteristics

After 3 days of incubation, 1.5 mL of culture was subsampled from each replicate culture every 24 h of the incubation. Then, 1 mL of each subsample was centrifuged at 16,000×g for 10 min at 4 °C, and the pellet was used for microbial DNA extraction per the repeated bead beating (RBB) + C method [25]. The supernatant of each culture sample was used to measure pH with an Accumet AB15 pH meter (Fisher Scientific, Suwanee, GA, USA) and determine the concentrations of VFA using gas chromatography (HP 5890 series, Agilent Technologies, Santa Clara, CA, USA) and ammonia using a colorimetric assay [26]. The remaining 0.5 mL culture was used to count protozoal cells as described above. The remaining content of each culture replicate was poured into a filter bag (Ankom Technology, USA; 25 mm porosity), and the retained solid was dried in a hot-air oven at 105 °C overnight [27]. The dry matter (DM) and neutral detergent fiber (NDF) content of the fresh protozoal feed and the residual feed after fermentation were determined following the method described by Van Soest et al. [28].

### Quantitative real-time polymerase chain reaction

Quantitative real-time PCR assays were used to quantify the total bacteria and three selected AAFB species (i.e., *Prevotella ruminicola*, *C. aminophilum*, and *P. anaerobius*). The PCR primer sets used are listed in Table 2. One sample-derived real-time PCR standard was produced using PCR amplification for each target group of microbes with the respective specific primer set and a DNA sample pooled from all the replicates of the three treatments as the template as described previously [29]. Each PCR product was electrophoresed on agarose (1%) gel to check its expected size and purified using a PCR purification kit (Qiagen, Germantown, MD, USA). The copy number concentration of each standard was calculated based on its length (bp). A serial dilution (10<sup>2</sup>–10<sup>11</sup> copies/mL) of each standard was used to quantify the abundance of the target bacteria by using an Mx3000 real-time PCR system (Stratagene, La Jolla, CA, USA). The thermal cycling profile and detailed PCR conditions are the same as described previously [30], but the annealing temperatures shown in Table 2 were used.

### Statistical analysis

Data from the three replicates were shown as the mean values of each measurement and subjected to the GLIMMIX procedure followed by Tukey's honestly significant difference (HSD) test by using SAS 9.4 (SAS Institute, Cary, NC, USA). The model to analyze the data was as follows:

**Table 2. Primers used in the quantification of the total bacteria and selected bacterial species via qPCR**

Target	Primer	Sequence, 5' to 3'	Annealing T (°C)	Product size (bp)	References
Total bacteria	27f	AGAGTTTGATCMTGGCTCAG	55	1,535	[46]
	1525r	AAGGAGGTGWTCARCC			
Total bacteria (qPCR)	Eub358f	TCCTACGGGAGGCAGCAGT	60	448	[47]
	Eub806r	GGACTACCAGGTATCTAATCCTGTT			
<i>Prevotella ruminicola</i>	P. rumi-F	GGTTATCTTGAGTGAGTT	53	485	[48]
	P. rumi-R	CTGATGGCAACTAAAGAA			
<i>Clostridium aminophilum</i>	C. amin-57F	ACGGAAATTACAGAAGGAAG	57	560	[49]
	C. amin-616R	GTTTCCAAGCAATTCCAC			
<i>Clostridium sticklandii</i>	C. stick-185F	ATCAAAGAATTCGGATAGG	61	442	[49]
	C. stick-626R	CAAGTTCACCAGTTTCAGAG			
<i>Peptostreptococcus anaerobius</i>	P. anae-73F	TGCTTGCAYTRATGAAAGATG	55	570	This study
	P. anae-642R	TCTTCCAGTTTCGGAGGCTA			

qPCR, quantitative real-time polymerase chain reaction; bp, base pair.

$$Y = m + T_i + t_j + (T \times t)_{ij} + e_{ij},$$


where  $Y$  is the dependent variable,  $m$  is the overall mean,  $T_i$  is the fixed effect of culture treatment,  $t_j$  is the time effect,  $(T \times t)_{ij}$  is the interaction between culture treatment and time, and  $e_{ij}$  is residual error. A Pearson correlation matrix was calculated for the fermentation characteristics and bacterial abundance by using the CORR procedure in SAS and visualized in R 3.2.2 [31]. The effects of incubation time and culture treatment were considered significant at  $p \leq 0.05$ .

## RESULTS


Approximately 99% (about a decrease by 2 logs) of the bacteria present in the monoculture were removed when *Ent. caudatum* cells were filtered and washed, but the total bacterial population increased dramatically after 24 h of incubation (Fig. 1). *Ent. caudatum* grew significantly better in the co-culture after 48 h of incubation (Fig. 2). The DM digestibility was greater in the co-culture, especially at 48 h and thereafter, than in *Ent. caudatum*- or AAFB-alone cultures (Table 3). However, no overall difference in NDF digestibility was noted among the three cultures. The ammonia concentration was higher in AAFB and Ec + AAFB co-cultures than in Ec throughout incubation. The ammonia concentration was higher in Ec + AAFB than in AAFB at 48 h but higher in AAFB than in Ec + AAFB at 72 h.

The total VFA production was higher in the *Ent. caudatum*-containing cultures, especially in the Ec + AAFB co-culture, than in AAFB (Table 4). The molar proportions of all VFAs differed ( $p < 0.01$ ) in these two cultures and at the two incubation times. More propionate but less acetate was produced in the *Ent. caudatum*-containing cultures, especially in the co-culture; as a result, the acetate:propionate ratio in the two *Ent. caudatum*-containing cultures was significantly lower than that in the AAFB culture. The proportion of branched-chain VFAs (BCVFA) was higher in the AAFB-containing cultures than in the two other cultures at 24 h of incubation but not at 72 h of incubation.

After analyzing the fermentation characteristics and protozoal counts, we quantified the total bacteria and *P. ruminicola*, *C. aminophilum*, and *P. anaerobius* in the cultures. At the beginning of incubation, the abundance of the three bacterial species was similar in the three cultures except in



**Fig. 1.** Abundance of (A) total bacteria, (B) *P. ruminicola*, (C) *C. aminophilum*, and (D) *P. anaerobius* in the cultures at different incubation times. Ec, *Entodinium caudatum* alone; AAFB, amino acid-fermenting bacterial consortium alone; Ec + AAFB, co-culture of Ec and AAFB.



**Fig. 2.** *Entodinium caudatum* cell counts in the two *Entodinium caudatum*-containing cultures (Ec alone and Ec + amino acid-fermenting bacterial consortium [AAFB]) at different time points.

the Ec culture which contained a smaller population of *P. anaerobius* (Fig. 1). Even though the Ec culture had the lowest abundance of total bacteria (at least 70-fold lower than that in the AAFB-containing cultures), it had a total bacterial abundance similar to that of Ec + AAFB after 24 h of incubation; conversely, AAFB alone had the smallest total bacterial population. At 24 h of incubation, the populations of the three selected bacterial species were similar and much larger in the *Ent. caudatum*-containing cultures than in the AAFB culture. After 72 h of incubation in the AAFB culture, *P. ruminicola* became undetectable. The initial population of *C. aminophilum* in all three cultures was not different, but it was significantly higher in *Ent. caudatum*-containing cultures than in the AAFB culture after 24 h of incubation and thereafter. Before co-culturing was performed, *P. anaerobius* was higher in AAFB-containing cultures, but this HAB species rapidly

**Table 3.** DM and NDF digestibility, NH<sub>3</sub>-N concentration, and pH in the cultures over time

	Incubation times (h)			SEM	Contrast	
	24	48	72		Linear	Quadratic
DM digestibility (%)						
Ec	67.5 <sup>a</sup>	69.0 <sup>b</sup>	80.4 <sup>b</sup>	2.09	< 0.001	0.004
AAFB	60.7 <sup>b</sup>	67.9 <sup>b</sup>	71.5 <sup>c</sup>	1.63	< 0.001	NS
Ec + AAFB	66.1 <sup>a</sup>	72.6 <sup>a</sup>	85.5 <sup>a</sup>	2.88	< 0.001	0.011
NDF digestibility (%)						
Ec	36.5	39.1	39.9	1.19	NS	NS
AAFB	35.9	43.5	44.5	1.88	NS	NS
Ec + AAFB	33.6	43.7	44.2	1.90	0.003	NS
NH <sub>3</sub> -N (mg/dL)						
Ec	7.1 <sup>b</sup>	13.6 <sup>c</sup>	16.5 <sup>c</sup>	1.39	< 0.001	0.010
AAFB	13.8 <sup>a</sup>	18.7 <sup>b</sup>	29.6 <sup>a</sup>	2.35	< 0.001	< 0.001
Ec + AAFB	14.8 <sup>a</sup>	21.5 <sup>a</sup>	26.2 <sup>b</sup>	1.66	< 0.001	0.033
pH						
Ec	6.41 <sup>b</sup>	6.17 <sup>b</sup>	6.12	0.07	NS	NS
AAFB	6.81 <sup>a</sup>	6.44 <sup>a</sup>	6.50	0.06	0.013	0.034
Ec + AAFB	6.47 <sup>b</sup>	6.27 <sup>ab</sup>	6.32	0.03	0.003	0.003

<sup>a-c</sup>Means within a row with different superscripts differ ( $p < 0.05$ ).

DM, dry matter; NDF, neutral detergent fiber; Ec, *Entodinium caudatum* alone; AAFB, amino acid-fermenting bacterial consortium alone; NS, not significant ( $p > 0.05$ ); Ec + AAFB, co-culture of Ec and AAFB.

grew in the Ec culture during the first 24 h of incubation. This rapid growth was maintained until 72 h of incubation, and this finding was comparable with that in the Ec-AAFB culture. Because *C. sticklandii*, which was previously known as a culturable HAB [16], was not detected using its specific primer set in the AAFB enrichment culture (AAFB inoculum), it was not quantified in the cultures. The relative abundances of the three AAFB species are also shown in Table 5.

Correlations between the fermentation characteristics and the abundance of select bacteria are shown in Fig. 3. Ammonia concentration was weakly correlated positively with the abundance of *C. aminophilum* and *P. anaerobius* ( $r = 0.33$  and  $r = 0.38$ , respectively) but negatively correlated with the abundance of *P. ruminicola* ( $r = -0.34$ ). *C. aminophilum* and *P. anaerobius* were strongly correlated positively with the total VFA concentrations ( $r > 0.8$ ,  $p < 0.001$ ) but negatively correlated with the valerate concentration ( $r < -0.8$ ,  $p < 0.001$ ). Moreover, the two HAB species were positively correlated with each other ( $p < 0.001$ ). The cell counts of *Ent. caudatum* were positively correlated ( $r \geq 0.57$ ) with the abundance of the three quantified bacterial species.

## DISCUSSION

Sitting at the top of the food chain in the rumen ecosystem, protozoa form a predator-prey relationship with all members of the ruminal microbiome. Because of their proteolytic activity, including the ability to degrade microbial proteins and produce oligopeptides and amino acids, ruminal protozoa also establish other relationships with other ruminal microbes, particularly AAFB. Conceptually, ruminal protozoa can engulf and provide substrates (i.e., oligopeptides and free amino acids) to AAFB, forging a commensalistic relationship beneficial to AAFB. However, these relationships have not been deterministically investigated because of the lack of and difficulties in obtaining axenic cultures of ruminal protozoa [20]. This study was the first to explore

**Table 4.** Molar proportion of volatile fatty acids (VFA) in cultures

	Incubation times (h)			SEM	Contrast	
	24	48	72		Linear	Quadratic
Total VFA (mM)						
Ec	33.5 <sup>a</sup>	45.9 <sup>b</sup>	70.9 <sup>a</sup>	5.54	< 0.001	0.008
AAFB	23.1 <sup>b</sup>	28.0 <sup>c</sup>	32.3 <sup>b</sup>	1.40	< 0.001	NS
Ec + AAFB	33.9 <sup>a</sup>	59.2 <sup>a</sup>	75.8 <sup>a</sup>	6.15	< 0.001	NS
VFA (mol/100 mol)						
Acetate						
Ec	76.3 <sup>a</sup>	67.0 <sup>b</sup>	53.8 <sup>b</sup>	3.28	< 0.001	0.002
AAFB	76.5 <sup>a</sup>	73.5 <sup>a</sup>	72.7 <sup>a</sup>	0.60	< 0.001	0.011
Ec + AAFB	72.5 <sup>b</sup>	53.8 <sup>c</sup>	46.8 <sup>c</sup>	3.84	< 0.001	< 0.001
Propionate						
Ec	15.2 <sup>b</sup>	24.1 <sup>b</sup>	37.8 <sup>b</sup>	3.30	< 0.001	0.0115
AAFB	12.10 <sup>c</sup>	16.1 <sup>c</sup>	15.3 <sup>c</sup>	0.61	< 0.001	< 0.0001
Ec + AAFB	17.1 <sup>a</sup>	37.2 <sup>a</sup>	43.4 <sup>a</sup>	3.98	< 0.001	< 0.0001
Butyrate						
Ec	6.61 <sup>c</sup>	7.29 <sup>b</sup>	6.74 <sup>c</sup>	0.15	NS	NS
AAFB	8.75 <sup>a</sup>	8.00 <sup>a</sup>	9.24 <sup>a</sup>	0.19	0.006	< 0.001
Ec + AAFB	8.24 <sup>b</sup>	7.34 <sup>b</sup>	7.79 <sup>b</sup>	0.14	0.021	0.002
Valerate						
Ec	0.66 <sup>b</sup>	0.49 <sup>b</sup>	0.35 <sup>b</sup>	0.05	< 0.001	NS
AAFB	1.00 <sup>a</sup>	0.84 <sup>a</sup>	0.76 <sup>a</sup>	0.04	< 0.001	NS
Ec + AAFB	0.67 <sup>b</sup>	0.44 <sup>c</sup>	0.37 <sup>b</sup>	0.05	< 0.001	< 0.001
Total BCVFA						
Ec	1.21 <sup>b</sup>	1.15 <sup>b</sup>	1.36	0.06	NS	NS
AAFB	1.61 <sup>a</sup>	1.54 <sup>a</sup>	1.97	0.09	NS	NS
Ec + AAFB	1.51 <sup>a</sup>	1.27 <sup>ab</sup>	1.64	0.07	NS	0.017
Acetate:Propionate ratio						
Ec	5.03 <sup>b</sup>	2.79 <sup>b</sup>	1.43 <sup>b</sup>	0.53	< 0.001	< 0.001
AAFB	6.33 <sup>a</sup>	4.57 <sup>a</sup>	4.74 <sup>a</sup>	0.28	< 0.001	< 0.001
Ec + AAFB	4.25 <sup>c</sup>	1.45 <sup>c</sup>	1.08 <sup>c</sup>	0.50	< 0.001	< 0.001

<sup>a-c</sup>Means within a row with different superscripts differ ( $p < 0.05$ ).

Ec, *Entodinium caudatum* alone; AAFB, amino acid-fermenting bacterial consortium alone; NS, not significant ( $p > 0.05$ ); Ec + AAFB, co-culture of both Ec and AAFB; BCVFA, branched-chain VFAs.

the interactions between these two important groups/guilds of ruminal microbes by using carefully washed *Ent. caudatum* and an AAFB consortium.

At the beginning of incubation, the total bacterial population was about 2 logs smaller in the Ec culture than in the two other cultures. The comparable abundance of total bacteria between the Ec culture and the Ec + AAFB co-culture at 24 h of incubation clearly showed that the residual bacteria remained after the *Ent. caudatum* inoculum was washed and grew rapidly, reaching an abundance similar to that in the Ec culture. In previous *in vitro* studies, antibiotics are used to remove prokaryotes associated with protozoa [32–34]. However, antibiotics inhibit, directly and indirectly, the viability of *Ent. caudatum* by killing its prey [20]. Although bacteria recovered after 24 h of incubation, they were protozoan-associated populations rather than free-living populations, and they probably function with ruminal protozoa. Therefore, the interactions between *Ent.*


**Table 5.** Relative abundance (% of total bacterial copy numbers) of the three selected bacterial species in each culture

	Incubation times (h)				SEM	<i>p</i> -values <sup>1)</sup>		
	0	24	48	72		Linear	Quadratic	Cubic
<i>Prevotella ruminicola</i>								
Ec	1.823 <sup>a</sup>	7.477 <sup>a</sup>	4.063 <sup>a</sup>	4.370 <sup>a</sup>	0.732	NS	0.024	0.018
AAFB	0.028 <sup>b</sup>	0.004 <sup>b</sup>	0.004 <sup>b</sup>	0 <sup>b</sup>	0.004	< 0.001	0.006	NS
Ec + AAFB	0.027 <sup>b</sup>	10.927 <sup>a</sup>	5.967 <sup>a</sup>	4.617 <sup>a</sup>	1.285	NS	0.001	0.008
<i>Clostridium aminophilum</i>								
Ec ( $\times 10^{-3}$ )	1.96 <sup>a</sup>	0.03 <sup>ab</sup>	0.04 <sup>a</sup>	0.11 <sup>a</sup>	0.25	< 0.001	< 0.001	0.013
AAFB ( $\times 10^{-5}$ )	2.21 <sup>b</sup>	0.54 <sup>b</sup>	3.55 <sup>b</sup>	1.80 <sup>b</sup>	0.49	NS	NS	0.037
Ec + AAFB ( $\times 10^{-5}$ )	2.62 <sup>b</sup>	7.91 <sup>a</sup>	7.69 <sup>a</sup>	14.6 <sup>a</sup>	1.41	< 0.001	NS	NS
<i>Peptostreptococcus anaerobius</i>								
Ec ( $\times 10^{-4}$ )	6.12 <sup>a</sup>	0.76 <sup>a</sup>	1.13	1.80 <sup>a</sup>	0.69	< 0.001	< 0.001	NS
AAFB ( $\times 10^{-5}$ )	1.29 <sup>b</sup>	3.59 <sup>b</sup>	4.34	3.10 <sup>c</sup>	0.66	NS	NS	NS
Ec + AAFB ( $\times 10^{-5}$ )	1.64 <sup>b</sup>	5.41 <sup>ab</sup>	7.00	11.3 <sup>b</sup>	1.09	< 0.001	NS	NS

<sup>1)</sup>*p*-values were for the contrast of the relative abundance of each bacterial species over time.

<sup>a-c</sup>Means within a row with different superscripts differ ( $p < 0.05$ ).

Ec, *Entodinium caudatum* alone; AAFB, amino acid-fermenting bacterial consortium alone; NS, not significance ( $p > 0.05$ ); Ec + AAFB, co-culture of both Ec and AAFB.



**Fig. 3.** Pearson correlation matrix of fermentation characteristics with molar proportions of VFAs and abundance of bacteria. Positive and negative correlations are shown in blue and red, respectively. Only strong correlations ( $r > 0.8$  or  $r < -0.8$ ) were indicated by a colored circle. The indicated correlation coefficients were also stated in the text. DMD, dry matter digestibility; NDFD, neutral detergent fiber digestibility; VFA, volatile fatty acid; BCVFA, branch-chained VFA; *P. ruminicola*, *Prevotella ruminicola*; *C. aminophilum*, *Clostridium aminophilum*; *P. anaerobius*, *Peptostreptococcus anaerobius*.

*caudatum* and AAFB could be inferred by comparing feed digestion, fermentation characteristics, or population dynamics of bacteria between Ec and Ec + AAFB.

The Ec + AAFB co-culture had the highest DM digestibility at 48 and 72 h of incubation possibly because of the greater abundance of *Ent. caudatum*, which degrades starch and hemicellulose [12,35]. Although Ec and Ec + AAFB had a similar total bacterial abundance, the latter had a larger *Ent. caudatum* population after 48 h of incubation. The NDF digestibility did not differ among the three cultures probably because *Ent. caudatum* cannot degrade cellulose [36], and neither the *Ent. caudatum* monoculture nor the AAFB consortium contained cellulolytic microbes. The AAFB culture had the lowest total VFA concentration, but it had a higher molar proportion of acetate, butyrate, valerate, and BCVFA than the two *Ent. caudatum*-containing cultures. Amino acid fermenters, including HAB, produce these VFAs as their common fermentation products [14,37]. Indeed, *C. aminophilum* and *P. anaerobius* can ferment a broad range of amino acids or casamino acids to ammonia, acetate, and butyrate [38,39].

*Ent. caudatum* grew faster in the Ec + AAFB co-culture than in the Ec culture during incubation. This finding verified the stimulatory effects on *Ent. caudatum* from the AAFB consortium. Although the Ec culture and the Ec + AAFB co-culture had similar total bacterial abundance, the latter had a greater *Ent. caudatum* population than the former. Thus, the Ec + AAFB co-culture could have higher bacterial recycling than the other culture. These findings were supported by higher concentrations of VFA and ammonia. The higher ammonia concentration in the Ec + AAFB co-culture than in the Ec or AAFB cultures also suggests that metabolic commensalism occurred through which *Ent. caudatum* provides substrates to AAFB and that AAFB are essential for ammonia production even though this bacterial consortium represents a small guild [40]. However, the Ec + AAFB co-culture did not increase the ammonia concentration after 72 h of incubation; this observation was inconsistent with a previous study that showed a two-fold higher ammonia concentration in *Entodinium*-faunated rumen than in fauna-free rumen of sheep [41]. In the present study, the protein added to the cultures might have been a limiting factor; therefore, ammoniogenesis in the Ec + AAFB co-culture could have been limited.

*C. aminophilum* and *P. anaerobius* are two of the three known HAB species [14]. Before incubation, their relative abundance was quite low (less than 0.002%). Their relative abundance linearly increased ( $p < 0.001$ ) during incubation in the Ec culture and the Ec + AAFB co-culture but not in the AAFB culture. These results suggest that *Ent. caudatum* provided the substrates for these two species and stimulated their growth. The three quantified bacterial species had similar abundance before incubation, but their populations increased in the Ec culture and the Ec + AAFB co-culture but not in the AAFB culture (except for *P. anaerobius*) at 24 h. *P. ruminicola* gradually decreased and became undetectable at 72 h. These results suggest that *Ent. caudatum* provided the substrates for the fermentation and growth of *C. aminophilum* and *P. anaerobius* but not *P. ruminicola*. Previous studies demonstrated the exopeptidase activity in rumen protozoal samples [42,43], and *Entodinium* species have a greater peptidase activity than large entodiniomorphs and holotrichs [44]. However, studies have yet to determine if the stimulatory effect of *Ent. caudatum* on the growth of *C. aminophilum* and *P. anaerobius* observed in the present study could be attributed to the exopeptidase activity of *Ent. caudatum*.

*P. ruminicola* can degrade dietary proteins but cannot use amino acids as its energy N source unless peptides are provided [11]. This ability might explain the decrease in *P. ruminicola* abundance in the AAFB culture and the more than 400-fold increase in *P. ruminicola* abundance in the Ec culture and the Ec + AAFB co-cultures during the first 24 h of incubation. *Ent. caudatum* counts were also positively correlated ( $r \geq 0.57$ ) with the abundance of the three quantified bacterial species. The Ec + AAFB co-culture had a higher ammonia concentration than the Ec culture, but

both cultures had a similar abundance of the three quantified bacterial species. Therefore, uncultured or other AAFB populations may also contribute to the deamination activity observed in this study, as shown in another study [45].

## CONCLUSION

Nitrogen utilization efficiency in ruminants is important for the economic viability of ruminant producers and the environment. The co-habitation and interactions of ruminal protozoa and AAFB contribute to the production of ammonia in the rumen. Our results verify that commensalism occurs between *Entodinium caudatum* and AAFB which benefits both microbial groups. The presence of *Ent. caudatum* in the cultures is also beneficial to HAB *Prevotella ruminicola*, *Clostridium aminophilum*, and *Peptostreptococcus anaerobius*. The abundance of *C. aminophilum* and *P. anaerobius* is correlated positively with ammonia concentration, verifying their role in ammoniogenesis. However, the prokaryotes that remained after washing the *Ent. caudatum* cells made it difficult to interpret some of the results. Understanding the interactions among the microbes involved in ruminal nitrogen metabolism remains challenging. Omics technologies, including genome-centric metagenomics, metatranscriptomics, and metabolomics, combined with stable isotope probing (SIP), should be used in future studies to help address this challenge.

## REFERENCES

1. Kohn RA, Dinneen MM, Russek-Cohen E. Using blood urea nitrogen to predict nitrogen excretion and efficiency of nitrogen utilization in cattle, sheep, goats, horses, pigs, and rats. *J Anim Sci*. 2005;83:879-89. <https://doi.org/10.2527/2005.834879x>
2. Huhtanen P, Hristov AN. A meta-analysis of the effects of dietary protein concentration and degradability on milk protein yield and milk N efficiency in dairy cows. *J Dairy Sci*. 2009;92:3222-32. <https://doi.org/10.3168/jds.2008-1352>
3. van Aardenne JA, Dentener FJ, Olivier JGJ, Goldewijk CGMK, Lelieveld J. A 1°×1° resolution data set of historical anthropogenic trace gas emissions for the period 1890–1990. *Global Biogeochem Cycles*. 2001;15:909-28. <https://doi.org/10.1029/2000GB001265>
4. Storm E, Ørskov ER, Smart R. The nutritive value of rumen micro-organisms in ruminants. 2. The apparent digestibility and net utilization of microbial N for growing lambs. *Br J Nutr*. 1983;50:471-8. <https://doi.org/10.1079/BJN19830115>
5. Leng RA, Nolan JV. Nitrogen metabolism in the rumen. *J Dairy Sci*. 1984;67:1072-89. [https://doi.org/10.3168/jds.S0022-0302\(84\)81409-5](https://doi.org/10.3168/jds.S0022-0302(84)81409-5)
6. Hackmann TJ, Firkins JL. Maximizing efficiency of rumen microbial protein production. *Front Microbiol*. 2015;6:465. <https://doi.org/10.3389/fmicb.2015.00465>
7. Hespell R, Akin D, Dehority B, Mackie R, White B, Isaacson R. Bacteria, fungi, and protozoa of the rumen. In: Mackie RI, White BA, Isaacson RE, editors. *Gastrointestinal microbiology*. Vol. 2, *Gastrointestinal microbes and host interactions*. New York, NY: Chapman and Hall; 1997.
8. Jouany JP. Effect of rumen protozoa on nitrogen utilization by ruminants. *J Nutr*. 1996;126:1335S-46S. [https://doi.org/10.1093/jn/126.suppl\\_4.1335S](https://doi.org/10.1093/jn/126.suppl_4.1335S)
9. Newbold CJ, de la Fuente G, Belanche A, Ramos-Morales E, McEwan NR. The role of ciliate protozoa in the rumen. *Front Microbiol*. 2015;6:1313. <https://doi.org/10.3389/fmicb.2015.01313>
10. Forsberg CW, Lovelock LKA, Krumholz L, Buchanan-Smith JG. Protease activities of rumen

- protozoa. *Appl Environ Microbiol.* 1984;47:101-10. <https://doi.org/10.1128/aem.47.1.101-110.1984>
11. Wallace RJ, Onodera R, Cotta MA. Metabolism of nitrogen-containing compounds. In: Hobson PN, Stewart CS, editors. *The rumen microbial ecosystem*. Dordrecht: Springer; 1997. p. 283-328.
  12. Williams AG, Coleman GS. *The rumen protozoa*. New York, NY: Springer; 1992.
  13. Eschenlauer SCP, McKain N, Walker ND, McEwan NR, Newbold CJ, Wallace RJ. Ammonia production by ruminal microorganisms and enumeration, isolation, and characterization of bacteria capable of growth on peptides and amino acids from the sheep rumen. *Appl Environ Microbiol.* 2002;68:4925-31. <https://doi.org/10.1128/AEM.68.10.4925-4931.2002>
  14. Attwood GT, Klieve AV, Ouwerkerk D, Patel BKC. Ammonia-hyperproducing bacteria from New Zealand ruminants. *Appl Environ Microbiol.* 1998;64:1796-804. <https://doi.org/10.1128/AEM.64.5.1796-1804.1998>
  15. Harlow BE, Flythe MD, Kagan IA, Aiken GE. Biochanin A (an isoflavone produced by red clover) promotes weight gain of steers grazed in mixed grass pastures and fed dried-distillers' grains. *Crop Sci.* 2017;57:506-14. <https://doi.org/10.2135/cropsci2016.07.0590>
  16. Flythe M, Kagan I. Antimicrobial effect of red clover (*Trifolium pratense*) phenolic extract on the ruminal hyper ammonia-producing bacterium, *Clostridium sticklandii*. *Curr Microbiol.* 2010;61:125-31. <https://doi.org/10.1007/s00284-010-9586-5>
  17. Taghavi-Nezhad M, Alipour D, Flythe MD, Zamani P, Khodakaramian G. The effect of essential oils of *Zataria multiflora* and *Mentha spicata* on the in vitro rumen fermentation, and growth and deaminative activity of amino acid-fermenting bacteria isolated from Mehraban sheep. *Anim Prod Sci.* 2014;54:299-307. <https://doi.org/10.1071/AN12244>
  18. Harlow BE, Bryant RW, Cohen SD, O'Connell SP, Flythe MD. Degradation of spent craft brewer's yeast by caprine rumen hyper ammonia-producing bacteria. *Lett Appl Microbiol.* 2016;63:307-12. <https://doi.org/10.1111/lam.12623>
  19. Dehority BA. Physiological characteristics of several rumen protozoa grown in vitro with observations on within and among species variation. *Eur J Protistol.* 2010;46:271-9. <https://doi.org/10.1016/j.ejop.2010.05.002>
  20. Park T, Meulia T, Firkins JL, Yu Z. Inhibition of the rumen ciliate *Entodinium caudatum* by antibiotics. *Front Microbiol.* 2017;8:1189. <https://doi.org/10.3389/fmicb.2017.01189>
  21. Gano JM. Amino acid-fermenting bacteria from the rumen of dairy cattle-enrichment, isolation, characterization, and interaction with *Entodinium caudatum* [Master's thesis]. Columbus, OH: The Ohio State University; 2013.
  22. Attwood GT, Reilly K. Identification of proteolytic rumen bacteria isolated from New Zealand cattle. *J Appl Bacteriol.* 1995;79:22-9. <https://doi.org/10.1111/j.1365-2672.1995.tb03119.x>
  23. Sutton S. Measurement of microbial cells by optical density. *J Valid Technol.* 2011;17:46-9.
  24. Dehority BA. Ciliate protozoa. In: Makkar HPS, McSweeney CS, editors. *Methods in gut microbial ecology for ruminants*. Dordrecht: Springer; 2005. p. 67-78.
  25. Yu Z, Morrison M. Improved extraction of PCR-quality community DNA from digesta and fecal samples. *Biotechniques.* 2004;36:808-12. <https://doi.org/10.2144/04365ST04>
  26. Chaney AL, Marbach EP. Modified reagents for determination of urea and ammonia. *Clin Chem.* 1962;8:130-2. <https://doi.org/10.1093/clinchem/8.2.130>
  27. AOAC [Association of Official Analytical Chemists]. *Official methods of analysis*. 15th ed. Arlington, VA: Association of Official Analytical Chemists; 1990.
  28. Van Soest PJ, Robertson JB, Lewis BA. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J Dairy Sci.* 1991;74:3583-97. <https://doi.org/10.3168/jds1991-74-3583>

- doi.org/10.3168/jds.S0022-0302(91)78551-2
29. Yu Z, Michel FC Jr, Hansen G, Wittum T, Morrison M. Development and application of real-time PCR assays for quantification of genes encoding tetracycline resistance. *Appl Environ Microbiol.* 2005;71:6926-33. <https://doi.org/10.1128/AEM.71.11.6926-6933.2005>
  30. Li YF, Chen PH, Yu Z. Spatial and temporal variations of microbial community in a mixed plug-flow loop reactor fed with dairy manure. *Microb Biotechnol.* 2014;7:332-46. <https://doi.org/10.1111/1751-7915.12125>
  31. R Core Team. R: a language and environment for statistical computing [Internet]. R Foundation for Statistical Computing. 2014 [cited 2022 Sep 7]. <http://www.R-project.org>
  32. Lee SS, Ha JK, Cheng KJ. Relative contributions of bacteria, protozoa, and fungi to in vitro degradation of orchard grass cell walls and their interactions. *Appl Environ Microbiol.* 2000;66:3807-13. <https://doi.org/10.1128/AEM.66.9.3807-3813.2000>
  33. Nagpal R, Puniya AK, Sehgal JP, Singh K. Influence of bacteria and protozoa from the rumen of buffalo on in-vitro activities of anaerobic fungus *Caecomyces* sp. isolated from the feces of elephant. *J Yeast Fungal Res.* 2010;1:152-6.
  34. Wei YQ, Long RJ, Yang H, Yang HJ, Shen XH, Shi RF, et al. Fiber degradation potential of natural co-cultures of *Neocallimastix frontalis* and *Methanobrevibacter ruminantium* isolated from yaks (*Bos grunniens*) grazing on the Qinghai Tibetan Plateau. *Anaerobe.* 2016;39:158-64. <https://doi.org/10.1016/j.anaerobe.2016.03.005>
  35. Béra-Maillet C, Devillard E, Cezette M, Jouany JP, Forano E. Xylanases and carboxymethylcellulases of the rumen protozoa *Polyplastron multivesiculatum* *Eudiplodinium maggii* and *Entodinium* sp. *FEMS Microbiol Lett.* 2005;244:149-56. <https://doi.org/10.1016/j.femsle.2005.01.035>
  36. Coleman GS. The cellulase content of 15 species of entodiniomorphid protozoa, mixed bacteria and plant debris isolated from the ovine rumen. *J Agric Sci.* 1985;104:349-60. <https://doi.org/10.1017/S0021859600044038>
  37. Chen G, Russell JB. More monensin-sensitive, ammonia-producing bacteria from the rumen. *Appl Environ Microbiol.* 1989;55:1052-7. <https://doi.org/10.1128/aem.55.5.1052-1057.1989>
  38. Paster BJ, Russell JB, Yang CMJ, Chow JM, Woese CR, Tanner R. Phylogeny of the ammonia-producing ruminal bacteria *Peptostreptococcus anaerobius*, *Clostridium sticklandii*, and *Clostridium aminophilum* sp. nov. *Int J Syst Bacteriol.* 1993;43:107-10. <https://doi.org/10.1099/00207713-43-1-107>
  39. Vos P, Garrity G, Jones D, Krieg NR, Ludwig W, Rainey FA, et al. *Bergey's manual of systematic bacteriology*. Vol. 3, The firmicutes. New York, NY: Springer; 2011.
  40. Walker ND, Newbold CJ, Wallace RJ. Nitrogen metabolism in the rumen. In: Pfeiffer E, Hristov A, editors. *Nitrogen and phosphorus nutrition of cattle: reducing the environmental impact of cattle operations*. Cambridge, MA: CABI Publishing; 2005. p. 71-115.
  41. Ivan M. Comparison of duodenal flow and digestibility in fauna-free sheep inoculated with *Holotrich* protozoa, *Entodinium* monofauna or total mixed protozoa population. *Br J Nutr.* 2009;101:34-40. <https://doi.org/10.1017/S0007114508984245>
  42. Abou Akkada AR, Howard BH. The biochemistry of rumen protozoa. 3. The carbohydrate metabolism of *Entodinium*. *Biochem J.* 1960;76:445-51. <https://doi.org/10.1042/bj0760445>
  43. Nagasawa T, Uchida T, Onodera R. Exopeptidase activity of mixed rumen ciliate protozoa. *Anim Sci Technol (Jpn).* 1992;63:481-7.
  44. Newbold CJ, McKain N, Wallace RJ. The role of protozoa in ruminal peptide metabolism. In: Lloyd D, Coombs GH, Paget TA, editors. *Biochemistry and molecular biology of "anaerobic" protozoa*. Chur: Harwood Academic; 1989. p.42-55.

45. Krause DO, Russell JB. An rRNA approach for assessing the role of obligate amino acid-fermenting bacteria in ruminal amino acid deamination. *Appl Environ Microbiol.* 1996;62:815-21. <https://doi.org/10.1128/aem.62.3.815-821.1996>
46. Lane DJ. 16S/23S rRNA sequencing. In: Stackebrandt E, Goodfellow M, editors. *Nucleic acid techniques in bacterial systematics*. Chichester, WS: John Wiley and Sons; 1991. p. 115-75.
47. Nadkarni MA, Martin FE, Jacques NA, Hunter N. Determination of bacterial load by real-time PCR using a broad-range (universal) probe and primers set. *Microbiology.* 2002;148:257-66. <https://doi.org/10.1099/00221287-148-1-257>
48. Tajima K, Aminov RI, Nagamine T, Matsui H, Nakamura M, Benno Y. Diet-dependent shifts in the bacterial population of the rumen revealed with real-time PCR. *Appl Environ Microbiol.* 2001;67:2766-74. <https://doi.org/10.1128/AEM.67.6.2766-2774.2001>
49. Patra AK, Yu Z. Effects of vanillin, quillaja saponin, and essential oils on in vitro fermentation and protein-degrading microorganisms of the rumen. *Appl Microbiol Biotechnol.* 2014; 98:897-905. <https://doi.org/10.1007/s00253-013-4930-x>