

Estimation of annual phosphorus excretion from pigs in Korea

Jong Young Ahn, Hansol Kim, Beob Gyun Kim*

Department of Animal Science, Konkuk University, Seoul 05029, Korea

Received: Mar 23, 2024

Revised: Apr 9, 2024

Accepted: Apr 15, 2024

*Corresponding author

Beob Gyun Kim

Department of Animal Science, Konkuk University, Seoul 05029, Korea

Tel: +82-2-2049-6255

E-mail: bgkim@konkuk.ac.kr

Copyright © 2025 Korean Society of Animal Science and Technology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Jong Young Ahn

<https://orcid.org/0000-0001-7893-459X>

Hansol Kim

<https://orcid.org/0000-0002-6088-8924>

Beob Gyun Kim

<https://orcid.org/0000-0003-2097-717X>

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

The authors are grateful for the support by Rural Development Administration, Korea (PJ017087).

Acknowledgements

Not applicable

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Kim BG.

Data curation: Ahn JY, Kim H.

Formal analysis: Ahn JY.

Abstract

The objective was to estimate the annual phosphorus (P) excretion from pigs in Korea based on P and phytate-P concentrations in commercial swine diets. Fifty-eight samples from commercial diets for various growth stages of pigs were collected from 16 swine farms and analyzed for P and phytate-P concentrations. The P concentrations ranged from 0.54% to 0.66%. Phytate-P contents in the piglet phase 1 and 2 diets were less (0.19% and 0.22% vs. 0.28% to 0.31%; $p < 0.05$) than those in the growing pig or sow diets. Fecal P excretion was calculated based on total P, phytate-P, and phytase concentrations: fecal P excretion (g/day) = [total P in feed (g/kg) – apparent total tract digestible (ATTD) P in feed (g/kg)] × feed intake (kg/day). The Gompertz model was adapted to estimate feed intake from suckling to finishing pigs, from birth to 121.5 kg body weight (BW), on days 0 to 180. Feed intake for gestating and lactating sows was adapted from the NRC. The ATTD P in the feed was estimated using the following equation: $\text{ATTD P (g/kg)} = 0.135 + 0.649 \times \text{total P (g/kg)} - 0.445 \times \text{phytate-P (g/kg)} + 0.470 \times \text{phytate-P} \times (1 - e^{-0.824 \times \text{phytase}})$. The phytase concentration in all diets was assumed to be 500 FTU/kg. Urinary P excretion for the market swine was estimated using the following equation: $\text{Urinary P excretion (g/day)} = \text{BW (kg)} \times 0.007 \text{ (g/kg BW/day)}$. The urinary P excretion for reproductive sows was adapted from a previous study. The total annual P excretion for market pigs was estimated to be 1.94 kg/year, which is equivalent to a pig with a BW of 44.1 kg at 93 days of age. For gestating and lactating sows, the total annual P excretion was estimated to be 3.26 kg/year and 6.89 kg/year, respectively. Assuming a population ratio of 91:7:2 for market pigs, gestating sows, and lactating sows, the annual P excretion from all market and breeding swine in Korea was estimated to be 2.15 kg/year.

Keywords: Phosphorus excretion, Pigs, Swine diets

INTRODUCTION

Phosphorus (P) plays a crucial role in both the skeletal system and various physiological functions in pigs [1]. In cereal grains, grain by-products, and oilseed meals, approximately two-thirds of P exists in the form of phytate-P with low digestibility in pigs [2,3]. Although exogenous phytase is commonly supplemented in commercial swine diets to improve P digestibility, phytate-P is not completely digested [4–6]. Unabsorbed P is excreted through swine feces, posing a potential risk of environmental pollution

Methodology: Ahn JY, Kim H, Kim BG.
 Investigation: Ahn JY.
 Writing - original draft: Ahn JY, Kim H.
 Writing - review & editing: Ahn JY, Kim H,
 Kim BG.

Ethics approval and consent to participate

This article does not require IRB/IACUC approval because there are no human and animal participants.

such as eutrophication [7,8]. Additionally, the quantities of excreted P would vary based on the types of feed ingredients included, the phytate-P concentration, and the phytase supplementation in swine diets [4–6].

An accurate estimation of P excretion from pigs is essential on a regional or national scale to develop efficient strategies for managing P excretion. Although European Union members routinely assess P excretion in individual countries [9–13], such data are lacking in Korea. In addition, the models for estimating P excretion from swine production developed more than 20 years ago [9–11] would not represent the present pig diets in Korea as the usage of phytase in swine diets has dramatically increased during the last 2 decades. The swine NRC [1] suggests models for pig growth and feed intake that are reasonably close to the Korean pig production systems. Therefore, this study aimed to estimate the annual P excretion per pig in Korea using the total P and phytate-P concentrations in commercial swine diets employing the models in the literature.

MATERIALS AND METHODS

Sample collection and chemical analyses

A total of 58 commercial swine diet samples were collected from 16 swine farms in Korea during various growth stages of pigs (Table 1). The diet samples were categorized into piglet phase 1 (7 to 15 kg; n = 11), piglet phase 2 (15 to 25 kg; n = 11), growing phase (25 to 50 kg; n = 10), finishing phase (50 to 121.5 kg; n = 7), gestation phase (n = 9), and lactation phase (n = 9). All samples were finely ground (< 0.1 mm) and stored at 4°C in the refrigerator until chemical analyses. The diet samples were analyzed for P using the molybdenum blue method (method 995.11) by UV spectrophotometer (UV-2450, Shimadzu, Kyoto, Japan) after dry-ash sample preparation as described by the AOAC [14]. Additionally, phytate-P in the diet samples was also analyzed using the commercial phytic acid assay kit (K-PHYT, Megazyme, Bray, Ireland) and the UV spectrophotometer (UV-2450, Shimadzu).

Calculations

To estimate the daily body weight (BW) for market pigs, the NLIN procedure of SAS (SAS Institute, Cary, NC, USA) was used based on BW and age data from the NRC [1]. The equation used in the Gompertz model [15] was:

$$\text{Gompertz model: } W_t \text{ (kg)} = Ae^{-be^{-kt}}$$

where W_t represents the age at time t (day), A was the BW of the mature pig, b was the growth ratio, k was the maturing rate, and e was the natural logarithm.

$$\text{Gompertz model: BW (kg)} = 217.4e^{-4.6919e^{-0.0116t}}$$

where t represents the age of the market pigs ($R^2 = 0.999$ and $p < 0.001$). The feed intake of market pigs was estimated using the default metabolizable energy (ME) intake equations for gilts and barrows and the effective ME content of the diet suggested by the NRC [1], as follows:

$$\text{Feed intake for market pigs (kg/day, as-fed basis)} = \frac{\text{default ME intake, gilts and barrows (kcal/day)}}{\div \text{effective ME content of the diet (kcal/kg)} \div \text{feed wastage correction coefficient}}$$

where the average of default ME intake for gilts and barrows was calculated based on the default

ME intake curves suggested by the NRC [1]. The values of 3,300 kcal/kg for the effective ME content of the diet and 5% feed wastage were also applied based on the assumption suggested by the NRC [1]. The feed wastage correction coefficient was 0.95. The feed intake for gestating and lactating sows was set at 2.12 and 5.34 kg/day, respectively, adapted from the NRC [1]. Fecal P excretion from pigs was estimated using the total P concentration in the diet, apparent total tract digestible (ATTD) P concentration in the diet, and feed intake as independent variables:

$$\text{Fecal P excretion (g/day)} = [\text{total P in diet (g/kg)} - \text{ATTD P in diet (g/kg)}] \times \text{feed intake (kg/day)}$$

The ATTD P in the diet was estimated using the total P, phytate-P, and phytase concentrations in the diet as independent variables, following the prediction equation suggested by Sung and Kim [6]:

$$\text{ATTD P (g/kg)} = 0.135 + 0.649 \times \text{total P (g/kg)} - 0.445 \times \text{phytate-P (g/kg)} + 0.470 \times \text{phytate-P (g/kg)} \times (1 - e^{-0.824 \times \text{phytase}})$$

where the total P and phytate-P concentrations in the diet were based on the analyzed data, and the phytase concentration in the diet was assumed to be 500 phytase unit (FTU)/kg for all diets. Urinary P excretion for market pigs was estimated using the following equation suggested by the NRC [1]:

$$\text{Urinary P excretion (g/day)} = \text{BW (kg)} \times 0.007 \text{ (g/kg BW/day)}$$

The urinary P excretion for gestating and lactating sows was assumed to be 2.50 and 2.40 g/day, respectively, based on the report by Grez-Capdeville and Crenshaw [16].

Total P excretion from the pigs was calculated as the sum of fecal and urinary P excretions:

$$\text{Total P excretion (g/day)} = \text{fecal P excretion (g/day)} + \text{urinary P excretion (g/day)}$$

The weighted mean of P excretion for breeding sows was calculated based on an estimated country population ratio of 80:20 for gestating sows and lactating sows. Additionally, P excretion for the entire pig was calculated based on an estimated country population ratio of 91:7:2 for market pigs, gestating sows, and lactating sows [17]. An example illustrating the calculation of the weighted mean of total P excretion for the entire swine population is presented in Fig. 1.

Statistical analyses

Data for P and phytate-P in the diet were analyzed using the MIXED procedure of SAS (SAS Institute, Cary, NC, USA). Each phase was included as a fixed variable in the model. Least squares means were calculated for the dietary total P and phytate-P concentrations for each phase and were compared using the PDIF option. Each diet was considered an experimental unit. Statistical significance was set at $p < 0.05$.

RESULTS

The total P concentrations in the diet ranged from 0.54% to 0.66% and did not differ among the phases (Table 1). Dietary phytate-P concentrations in the piglet phases were lower ($p < 0.05$) than




Fig. 1. An example illustrating the calculations from total phosphorus (P) excretion per pig per day to weighted mean of total P excretion for overall swine population per pig per year.

Table 1. Phosphorus (P) and phytate-P concentrations in commercial swine diet in Korea (as-fed basis)

Category	Body weight (kg)	n	Total P (%)		Phytate-P (%)	
			Mean	SD	Mean	SD
Piglet phase 1	7 to 15	11	0.59	0.11	0.19 ^b	0.06
Piglet phase 2	15 to 25	11	0.60	0.11	0.22 ^b	0.07
Growing phase	25 to 50	10	0.58	0.10	0.30 ^a	0.06
Finishing phase	50 to 121.5	7	0.54	0.10	0.28 ^a	0.07
Gestating sow	-	9	0.66	0.12	0.30 ^a	0.05
Lactating sow	-	9	0.66	0.07	0.31 ^a	0.04
SEM	-	-	0.03	-	0.02	-
<i>p</i> -value	-	-	0.152	-	< 0.001	-

^{a,b} Means within a column without a common superscript letter differ ($p < 0.05$).

those in other phases.

Fecal, urinary, and total P excretions for market pigs from 7 to 121.5 kg BW at 27 to 180 days of age were 4.93, 0.40, and 5.33 g/day, respectively (Table 2). For gestating sows, fecal, urinary, and total P excretion were 6.70, 2.50, and 9.20 g/day, respectively. Fecal, urinary, and total P excretion of lactating sows were 19.31, 2.40, and 21.71 g/day, respectively. Estimated annual total P excretion was 1.94 kg/year for market pigs, 3.26 kg/year for gestating sows, and 6.89 kg/year for lactating sows. Collectively, the weighted mean of the total P excretion for the entire swine population in Korea was 5.90 g/day or 2.15 kg/year.

As the age of the pig increases, daily P excretion increases from 0.98 to 7.65 g/day for fecal P excretion and from 0.05 to 0.85 g/day for urinary P excretion (Fig. 2). From 7 to 121.5 kg BW at ages of 27 to 180 days, the total P excretion for market pigs is 5.33 g/day or 0.82 kg/pig per production cycle. The representative BW for the P excretion is 44.1 kg at the age of day 93.

DISCUSSION

In market pigs diets, the analyzed total P concentration of the commercial diets for piglet and

Table 2. Estimated fecal, urinary, and total phosphorus (P) excretion of various growth phases in pigs

	Market pig					Breeding sow			Entire pig ³⁾
	Piglet phase 1	Piglet phase 2	Growing phase	Finishing phase	Overall ¹⁾	Gestating sow	Lactating sow	Overall ²⁾	
Body weight range (kg)	7 to 15	15 to 25	25 to 50	50 to 121.5	7 to 121.5	-	-	-	-
Age range (day)	27 to 48	49 to 66	67 to 100	101 to 180	27 to 180	-	-	-	-
Total P in diet (g/kg)	5.91	6.01	5.78	5.44	5.65	6.57	6.64	-	-
ATTD P in diet ⁴⁾ (g/kg)	3.42	3.40	3.03	2.86	3.04	3.54	3.56	-	-
Feed intake (kg/day)	0.57	0.98	1.60	2.58	1.89	2.21	6.26	-	-
Total P intake (g/day)	3.38	5.92	9.22	14.04	10.68	14.52	41.60	-	-
ATTD P intake (g/day)	1.96	3.35	4.83	7.39	5.75	7.83	22.28	-	-
Daily P excretion (g/day)									
Fecal P excretion	1.42	2.57	4.39	6.65	4.93	6.70	19.31	9.17	5.32
Urinary P excretion	0.07	0.14	0.26	0.61	0.40	2.50	2.40	2.48	0.59
Total P excretion ⁵⁾	1.50	2.71	4.65	7.26	5.33	9.20	21.71	11.65	5.90
Yearly P excretion (kg/year)									
Fecal P excretion	0.52	0.94	1.60	2.43	1.80	2.44	7.05	3.35	1.94
Urinary P excretion	0.03	0.05	0.09	0.22	0.15	0.91	0.88	0.91	0.21
Total P excretion	0.55	0.99	1.70	2.65	1.94	3.36	7.92	4.25	2.15

¹⁾Weaning to finishing pigs from 7 to 121.5 kg body weight at ages of 27 to 180 days.

²⁾Phosphorus excretion for the breeding sows was calculated based on the assumption of a population ratio of 80:20 for gestating sows and lactating sows.

³⁾Phosphorus excretion for the entire pig was calculated based on the assumption of a population ratio of 91:7:2 for market pigs, gestating sows, and lactating sows.

⁴⁾The ATTD P in the diet was estimated using the total P, phytate-P, and phytase concentrations in the diet as independent variables, from the literature [6]. Phytase concentration was assumed to be constant at 500 FTU/kg in all diets.

⁵⁾Total P excretion = fecal P excretion + urinary P excretion.

ATTD, apparent total tract digestible.




Fig. 2. Estimated fecal, urinary, and total phosphorus (P) excretion of market pigs from 7 to 121.5 kg body weight at ages of 27 to 180 days. The fecal P excretion was estimated using the total P, phytate-P, and apparent total tract digestible (ATTD) P concentrations in the diets, and feed intake as independent variables. The ATTD P in the diet was estimated using the total P, phytate-P, and phytase concentrations in the diet as independent variables, as suggested by Sung and Kim [6]. Urinary P excretion for market pigs was estimated using the equation suggested by the NRC [1]. The mean fecal, urinary, and total P excretion for market pigs were 4.93, 0.40, and 5.33 g/day, respectively, equivalent to a pig weighing 44.1 kg BW at the age of day 93.

growing phases closely matched the requirement estimates in the NRC [1]. However, the analyzed total P concentrations in finishing diets for pigs weighing over 75 kg were greater than the NRC requirement estimates. This deviation in the finishing phase is likely attributed to the assumption that the same diet was provided to the pigs from 50 kg until marketing of pigs in the present work. As pigs grow, their nutrient requirements and concentrations in the diet gradually decrease [1]. Nevertheless, in the Korean swine production system, the diet for growing phase is sometimes fed to growing and finishing phases [18]. Feeding growing diets to finishing pigs is likely a strategy to cope with challenging production conditions, such as hot summers and cold winters in Korea, where feed replacement can be difficult. Therefore, it was assumed that P intake exceeded the requirements for finishing pigs in this study, resulting in higher P excretion compared to diets that marginally met the P requirements for finishing pigs.

In the gestating and lactating sow diets, the analyzed P concentrations were similar to or higher than the values suggested by the NRC [1]. The P content in the gestating sow diets (0.66%) exceeded the NRC [1] requirement estimates (ranging from 0.38% to 0.62%, total P basis), which are based on variables including parity, anticipated gestational weight gain, anticipated litter size, and days of gestation. Similarly, the P content in the lactating sow diets (0.66%) either exceeded or closely matched the requirement estimates (ranging from 0.54% to 0.67%, total P basis), considering variables such as parity, post-farrowing BW, litter size, lactation length, and mean daily weight gain of nursing pigs. Variations between the P concentration of diets in the present study and the suggested requirement estimates by the NRC [1] can be attributed to the inclusion of safety margins for nutrients in the diet formulation process, especially for the gestating and lactating sow diets, which are typically fed as a single diet throughout each phase.

Research on P utilization and excretion in pigs has been conducted in several countries [9–13]. P excretion can be expressed in two ways. The first expression quantifies the amount of P excreted by pigs during their entire life or during a specific production period in kilograms per pig. For example, Jongbloed et al. [11] reported a P excretion of 0.19 kg/pig for piglets weighing from 7.5 to 30 kg BW in Denmark. Additionally, CORPEN reported a P excretion of 0.74 kg/pig for pigs from wean to finish (8 to 108 kg) in France [19]. This approach is particularly relevant for the production cycle of market pigs, which consists of approximately 6 months from birth to slaughter. The second expression for P excretion was the annual amount (kg/year), typically applied to breeding sows that live for more than a year [9,10]. Our study provides information on P excretion in market pigs based on age and BW, allowing for conversion to production period-based excretion as well as daily or yearly excretion values.

The breeding sows were categorized as gestating or lactating. In typical commercial swine farms, newborn piglets have a suckling period lasting 3 to 4 weeks, with a weaning weight of approximately 6.5 to 7.5 kg. This study assumed that nutrient intake during the suckling period was solely from sow milk. Consequently, the amount of feed intake from the diet for pigs under 7.0 kg was set to zero. Therefore, the estimated P excretion of market pigs ranged from 7 to 121.5 kg BW at ages of 27 to 180 days.

The estimated P excretion for market pigs with a BW ranging from 7 to 121.5 kg in this study agreed with values reported in the literature [9–13,19]. However, P excretion for breeding sows (4.25 kg/year) in this study was relatively lower than the values reported in the literature. CORPEN [19] reported sow P excretion in France as 5.10 kg/year (with standard feeding) and 6.50 kg/year (with 2-phase feeding). Jongbloed et al. [11] also reported sow P excretion as 6.92 kg/year in Denmark, 6.71 kg/year in France, and 4.04 kg/year in the Netherlands. These discrepancies are likely due to factors not considered in the estimation of P excretion for breeding sows in this study, such as phytase efficacy [20], variations in P digestibility during gestation and lactation periods

[21], and reproductive performance. Another important factor for discrepancies is the amount of P excretion for suckling pigs. Studies on P excretion in sows have reported the amounts of P excretion measured with their offspring [9–11,13,19]. However, the present study did not consider the P excretion for suckling pigs because the estimation was based on the P and phytate-P concentrations in commercial diets. More research on P excretion from suckling pigs is needed.

The BW specifications for pigs in different growth stages vary among the NRC [1], CVB [22], and Korean Feeding Standards for Swine [23]. The Gompertz growth curve for BW was adapted using NRC [1] data. However, the calculation of ATTD P (g/kg of diet) to determine fecal P excretion utilized the actual weights of pigs fed each diet, as outlined in Table 1. Consequently, fecal P excretion in market pigs was calculated based on the phase feeding of the four different diets throughout their lifetime.

For pigs weighing over 7 kg, ATTD P was calculated based on the analyzed data for total P and phytate-P contents in each diet fed according to their weight. The prediction equation for ATTD P in the diet, as suggested by Sung and Kim [6], was applied by incorporating the total P, phytate-P, and a constant phytase concentration of 500 FTU/kg in all diets, a common phytase supplementation level in commercial swine diets [24]. As a result, the ATTD P values varied depending on the total P and phytate-P content of the diets. Therefore, Fig. 2 illustrating P excretion, confirms the discontinuation in excretion levels at the point of diet transition.

Based on the results of this study, the annual excretion of fecal P was calculated by multiplying the fecal P excretion (Table 2; kg/year) by the entire swine population in Korea (11.1 million pigs; [17]), resulting in 20,994 tons/year. We validated this annual fecal P excretion by comparing it with the production weight of swine feed in Korea. Utilizing data on the production weight of swine feed for respective phases in Korea in 2022 [18] and dietary ATTD P (Table 1; [6]), we calculated the annual fecal P excretion to be 19,062 tons/year. This amount of excretion closely approximates the estimated annual fecal P excretion in the present study, indicating that the present estimations employing the NRC [1] models were fairly reasonable.

Sung and Kim [6] reported that the prediction equation for ATTD P may not be applicable to sows because it is based on observations from growing pigs. Additionally, variations in phytase efficacy [25] and P digestibility [21,26] may occur depending on the gestating or lactating stage of the sow and even during the gestation period. However, because of the lack of an alternative prediction equation that can estimate ATTD P based solely on the dietary concentrations of P and phytate-P, the equation suggested by Sung and Kim [6] was utilized. The calculated ATTD P values for gestating and lactating sows using this equation were higher than the values recommended by the NRC [1] for gestating and lactating sows. Consequently, the P excretion estimated in this study for breeding sows may have been underestimated compared to the actual values.

Taken together, the P excretion of pigs for each growing phase was estimated based on the total P and phytate-P concentrations in commercial swine diets. The total annual P excretion for market pigs was estimated to be 1.94 kg/year, equivalent to a pig with a BW of 44.1 kg at 93 days of age. For gestating and lactating sows, the total annual P excretion was estimated to be 3.36 kg/year and 7.92 kg/year, respectively. Assuming a population ratio of 91:7:2 for market pigs, gestating sows, and lactating sows, the annual P excretion from all market and breeding swine in Korea was estimated to be 2.15 kg/year. Further research is required to validate our estimation through *in vivo* experiments using commercial diets.

REFERENCES

1. NRC (National Research Council). Nutrient requirements of swine. 11th ed. Washington, DC: National Academies Press; 2012.
2. Lee SA, Lopez DA, Stein HH. Mineral composition and phosphorus digestibility in feed phosphates fed to pigs and poultry. *Anim Biosci.* 2023;36:167-74. <https://doi.org/10.5713/ab.22.0322>
3. Stein HH, Lagos LV, Casas GA. Nutritional value of feed ingredients of plant origin fed to pigs. *Anim Feed Sci Technol.* 2016;218:33-69. <https://doi.org/10.1016/j.anifeedsci.2016.05.003>
4. Almeida FN, Stein HH. Performance and phosphorus balance of pigs fed diets formulated on the basis of values for standardized total tract digestibility of phosphorus. *J Anim Sci.* 2010;88:2968-77. <https://doi.org/10.2527/jas.2009-2285>
5. Passos AA, Moita VHC, Kim SW. Individual or combinational use of phytase, protease, and xylanase for the impacts on total tract digestibility of corn, soybean meal, and distillers dried grains with soluble fed to pigs. *Anim Biosci.* 2023;36:1869-79. <https://doi.org/10.5713/ab.23.0212>
6. Sung JY, Kim BG. Prediction models for apparent and standardized total tract digestible phosphorus in swine diets. *Anim Feed Sci Technol.* 2019;255:114224. <https://doi.org/10.1016/j.anifeedsci.2019.114224>
7. Won S, You BG, Shim S, Ahmed N, Choi YS, Ra C. Nutrient variations from swine manure to agricultural land. *Asian-Australas J Anim Sci.* 2018;31:763-72. <https://doi.org/10.5713/ajas.17.0634>
8. Lautrou M, Cappelaere L, Létourneau Montminy MP. Phosphorus and nitrogen nutrition in swine production. *Anim Front.* 2022;12:23-9. <https://doi.org/10.1093/af/vfac068>
9. Bohnenkemper O. Bilanzierung der nährstoffausscheidungen landwirtschaftlicher nutztiere. Frankfurt am Main: DLG-Verlag; 2005.
10. Jongbloed AW, Kemme PA. De uitscheiding van stikstof en fosfor door varkens, kippen, kalkoenen, pelsdieren, eenden, konijnen en parelhoeders in 2002 en 2006. Lelystad: Animal Sciences Group Wageningen UR; 2005. Rapport No.: 05/I01077.
11. Jongbloed AW, Poulsen HD, Dourmad JY, van der Peet-Schwering CMC. Environmental and legislative aspects of pig production in The Netherlands, France and Denmark. *Livest Prod Sci.* 1999;58:243-9. [https://doi.org/10.1016/S0301-6226\(99\)00012-3](https://doi.org/10.1016/S0301-6226(99)00012-3)
12. Jørgensen H, Prapasongsa T, Vu VTK, Poulsen HD. Models to quantify excretion of dry matter, nitrogen, phosphorus and carbon in growing pigs fed regional diets. *J Anim Sci Biotechnol.* 2013;4:42. <https://doi.org/10.1186/2049-1891-4-42>
13. Šebek LB, Bikker P, Vuuren AM, Krimpen M. Nitrogen and phosphorous excretion factors of livestock. Task 2: in-depth analyses of selected country reports. Wageningen: Wageningen UR Livestock Research; 2014.
14. AOAC (Association of Official Analytical Chemists) International. Official methods of analysis of AOAC International. 21st ed. Gaithersburg, MD: AOAC International; 2019.
15. Gompertz B. XXIV. On the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies. In a letter to Francis Baily, Esq. F.R.S &c. *Philos Trans R Soc Lond.* 1825;115:513-83. <https://doi.org/10.1098/rstl.1825.0026>
16. Grez-Capdeville M, Crenshaw TD. Estimation of phosphorus requirements of sows based on 24-h urinary phosphorus excretion during gestation and lactation. *Br J Nutr.* 2022;128:377-88.

<https://doi.org/10.1017/S0007114521003421>

17. Statistics Korea. 2022 Annual livestock survey report. Daejeon: Statistics Korea; 2023.
18. MAFRA (Ministry of Agriculture, Food and Rural Affairs). Livestock feed production statistics. Sejong: MAFRA; 2023.
19. CORPEN (Comité d'Orientation pour la Réduction de la Pollution des Eaux par les Nitrates et les Phosphates). Estimation des rejets d'azote - phosphore - potassium - cuivre et zinc des porcs. Influence de la conduite alimentaire et du mode de logement des animaux sur la nature et la gestion des déjections produites. Paris: CORPEN; 2003.
20. Kemme PA, Jongbloed AW, Mroz Z, Beynen AC. The efficacy of *Aspergillus niger* phytase in rendering phytate phosphorus available for absorption in pigs is influenced by pig physiological status. *J Anim Sci*. 1997;75:2129-38. <https://doi.org/10.2527/1997.7582129x>
21. Lee SA, Lagos LV, Walk CL, Stein HH. Basal endogenous loss, standardized total tract digestibility of calcium in calcium carbonate, and retention of calcium in gestating sows change during gestation, but microbial phytase reduces basal endogenous loss of calcium. *J Anim Sci*. 2019;97:1712-21. <https://doi.org/10.1093/jas/skz048>
22. Bikker P, Blok MC. Phosphorus and calcium requirements of growing pigs and sows. Wageningen: Wageningen Livestock Research; 2017. CVB Documentation Report No.: 59.
23. NIAS (National Institute of Animal Science). Korean feeding standard for swine. 4th rev. ed. Wanju: NIAS; 2022.
24. Lagos LV, Bedford MR, Stein HH. Apparent digestibility of energy and nutrients and efficiency of microbial phytase is influenced by body weight of pigs. *J Anim Sci*. 2022;100:skac269. <https://doi.org/10.1093/jas/skac269>
25. Jongbloed AW, van Diepen JTM, Kemme PA, Broz J. Efficacy of microbial phytase on mineral digestibility in diets for gestating and lactating sows. *Livest Prod Sci*. 2004;91:143-55. <https://doi.org/10.1016/j.livprodsci.2004.07.017>
26. Lee SA, Bedford MR, Stein HH. Comparative digestibility and retention of calcium and phosphorus in normal- and high-phytate diets fed to gestating sows and growing pigs. *Anim Feed Sci Technol*. 2021;280:115084. <https://doi.org/10.1016/j.anifeedsci.2021.115084>