

Efficacy and matrix value of xylanase in broiler fed an energy-deficient corn-wheat-soybean diet

Jun Seon Hong^{1#}, Myunghwan Yu^{1#}, Shan Randima Nawarathne¹, Elijah Ogola Oketch¹, Hyeonho Yun², Dinesh Darshaka Jayasena³, Jung Min Heo^{1*}

¹Department of Animal Science and Biotechnology, Chungnam National University, Daejeon 34134, Korea

²Technical Marketing Division, Protein Solution Business Unit, CJ CheilJedang Bio, Seoul 04560, Korea

³Department of Animal Science, Uva Wellassa University of Sri Lanka, Badulla 90000, Sri Lanka



Received: Mar 19, 2024

Revised: Mar 28, 2024

Accepted: Apr 4, 2024

#These authors contributed equally to this work.

*Corresponding author

Jung Min Heo

Department of Animal Science and Biotechnology, Chungnam National University, Daejeon 34134, Korea.

Tel: +82-42-821-5777

E-mail: jmheo@cnu.ac.kr

Copyright © 2025 Korean Society of Animal Science and Technology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Jun Seon Hong

<https://orcid.org/0000-0003-2142-9888>

Myunghwan Yu

<https://orcid.org/0000-0003-4479-4677>

Shan Randima Nawarathne

<https://orcid.org/0000-0001-9055-9155>

Elijah Ogola Oketch

<https://orcid.org/0000-0003-4364-460X>

Hyeonho Yun

<https://orcid.org/0009-0001-3215-822X>

Dinesh Darshaka Jayasena

<https://orcid.org/0000-0002-2251-4200>

Jung Min Heo

<https://orcid.org/0000-0002-3693-1320>

Abstract

The experiment was conducted to estimate the matrix value of xylanase and the effect of dietary xylanase supplementation on growth performance, viscosity, digestibility, and carcass traits in broiler chickens. A total of 588 one-day-old Ross 308 broiler chicks were raised with a commercial diet until day 7 and seven-day-old chicks were randomly allotted to one of seven dietary treatments with twelve replicates. Diets were corn-soybean meat based with wheat added. Dietary treatments were as follows; four basal diets (PC, energy sufficient diet; NC-1, -40 kcal/kg ME reduced from PC diet; NC-2, -80 kcal/kg ME reduced from PC diet; NC-3, -120 kcal/kg ME reduced from PC diet) and three different xylanase activity levels diet (NCX-1, 1,500 U/kg xylanase activity; NCX-2, 3,000 U/kg xylanase activity, NCX-3 4,500 U/kg xylanase activity) in the NC-3 diet. The standard xylanase dose was decided from the previous *in vitro* experiment. The weight gain and feed intake were measured and feed efficiency was calculated weekly. One bird per pen was selected and euthanized to harvest the intestinal digesta, breast meat, and leg meat samples on days 24 and 35. The linear and quadratic regression analysis and regression plateau were used to determine the xylanase recommendation and marginal level. The viscosity, digestibility, and proximate analysis of meat were analyzed from taken samples. Xylanase-added treatments were performed for higher ($p < 0.05$) body weight and body weight gain. Furthermore, xylanase-added treatments showed higher protein digestibility and lower viscosity compared to non-xylanase treatment. The maximum metabolizable energy compensation level of xylanase calculated by the regression was 120 kcal/kg and the marginal xylanase level showed maximum performances were 3,622 U/kg on the linear plateau and 4,000 U/kg on the quadratic plateau. Therefore, our experiment suggested that xylanase addition in an energy deficiency diet not only enhances growth performance but also reduces viscosity, and enhances protein digestibility and the maximum compensation level of metabolizable energy was 120 kcal/kg. The recommended levels of xylanase supplementation were determined to be 3,622 and 4,000 U/kg.

Keywords: Broiler, Carcass trait, Digestibility, Growth performance, Xylanase

Competing interests

The author Hyeonho Yun is an employee of CJ CheilJedang Bio and declares no conflicts of interest.

Funding sources

Not applicable.

Acknowledgements

The authors would like to show their appreciation to CJ CheilJedang Corp., Korea, who supported this research through financial assistance and the provision of additives.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Hong JS, Yun H, Heo JM.
Data curation: Hong JS, Heo JM.
Formal analysis: Hong JS.
Methodology: Hong JS, Yu M.
Software: Hong JS, Yu M.
Validation: Hong JS, Yu M, Jayasena DD, Heo JM.
Investigation: Hong JS, Yu M, Nawarathne SR, Oketch EO.
Writing - original draft: Hong JS, Yu M.
Writing - review & editing: Hong JS, Yu M, Nawarathne SR, Oketch EO, Yun H, Jayasena DD, Heo JM.

Ethics approval and consent to participate

All experimental procedures were revised and endorsed by the Animal Care and Use Committee of Chungnam National University, Daejeon, Korea (Protocol No. 202103A-CNU-062).

INTRODUCTION

The recent unstable international situation makes the price of ingredients used in animal diets fluctuate negatively and those changes increase the overall cost of broiler production. This circumstance makes feed formulators focus on using alternative feed ingredients and feed additives such as enzymes, probiotics, prebiotics, and functional biotics to reduce the feed cost without any disadvantage.

Corn and wheat are majorly used for animal diets, and those contain about 15% soluble and insoluble non-starch polysaccharide (NSP) [1]. Furthermore, arabinoxylans are the major components in NSP which are enriched in plant-originated ingredients. Arabinoxylans act like an anti-nutritional factor that increases viscosity in digesta and decreases feed efficiency and growth performances when they are fed to broilers [2,3]. Increased viscosity could change the dominant microbiota which might negatively affect digestion and absorption for the host animals [4]. Arabinoxylan polymers also encapsulate the nutrient unavailable by the host animal and it is been estimated about 400 to 450 kcal/kg [5]. Therefore, the supplementation of xylanase in poultry diets has become an essential factor in increasing nutrient utilization by degrading arabinoxylan complexes [6]. Xylanases are generally produced by a plethora of organisms including bacteria, algae, fungi, protozoa, gastropods, and arthropods. Xylanases are glycosidases that catalyze the endohydrolytic of 1,4- β -D-xylosidic linkages in the xylan complex to produce xylose, a primary carbon source for cell metabolism and in plant cell infection by plant pathogens [7]. Xylanases are also known for releasing encapsulated nutrients by degrading xylosidic linkage in the cell wall [8] but some recent studies suggest that degrading the cell wall and releasing nutrients are supported by very limiting *in vivo* experiments [6,9].

Previous studies about adding xylanases to poultry diets have suggested that the addition of xylanase could improve growth performance, nutrient digestibility, digesta viscosity, and carcass traits [10–14]. However, referred studies suggest different levels of xylanase activity because of the different diet formulations. The matrix value of enzymes presented approximate compensated nutrient levels by degrading and catalyzing with enzymes. Accurately measured matrix values are important when commercial feed formulators are required to use the enzyme. Overestimated matrix values can cause not only depressing performances but also animal welfare and underestimation of the potential loss. Thus, accurate and achievable matrix value estimation in the commercial should be required [15]. Therefore, this experiment was conducted to evaluate the physiological effect and estimate the matrix value of dietary xylanase in the corn-wheat-soybean diet on broiler chickens.

MATERIALS AND METHODS

All experimental procedures were revised and endorsed by the Animal Care and Use Committee of Chungnam National University, Daejeon, Korea (Protocol No. 202103A-CNU-062).

Birds and housing

An experiment was conducted using 588 Ross broilers from day 1 to 35 days of age. Seven birds were randomly allotted and raised in each wire floor cage ($76 \times 61 \times 46 \text{ cm}^3$) from day 1 to 7 as an adaptation period, and they were reallocated following similar body weight ([BW], $134.27 \pm 0.950 \text{ g}$; mean \pm SEM) and weight distribution on day 8. Each pen was installed with three nipple drinkers and a metal trough. Experimental diets and fresh water were provided via nipple drinker and metal trough on an *ad libitum* basis. All the management practices were followed by Ross 308 broiler management guidance (Table 1).

Table 1. Seven different dietary treatments of the experimental design

| Treatment | Description |
|-----------|--|
| PC | Positive control diet to meet or exceed Ross 308 specification |
| NC-1 | 40 kcal/kg ME level reduced from PC diet |
| NC-2 | 80 kcal/kg ME level reduced from PC diet |
| NC-3 | 120 kcal/kg ME level reduced from PC diet |
| NCX-1 | NC-3 diet with 1,500 U/kg xylanase supplementation |
| NCX-2 | NC-3 diet with 3,000 U/kg xylanase supplementation |
| NCX-3 | NC-3 diet with 4,500 U/kg xylanase supplementation |

NC, negative control; ME, metabolizable energy.

Experimental design and diets

The experiment was conducted using 588 one-day-old Ross 308 broiler chicks for 35 days. All birds were raised with a commercial diet from day 1 to 7 days of age as an adaptation period. On day 8, birds were reallocated to one of seven dietary treatments arranged in a completely randomized design and each treatment had 12 replicate cages. Four basal diets (PC, energy sufficient diet; NC-1, -40 kcal/kg metabolizable energy (ME) reduced from PC diet; NC-2, -80 kcal/kg ME reduced from PC diet; NC-3, -120 kcal/kg ME reduced from PC diet) were formulated based on corn, wheat, and soybean meal (Tables 2 and 3) to meet or exceed the nutrient requirements of Aviagen [16], except for energy levels, with two-phase feeding which is the starter (day 8 to 24) and the grower (day 25 to 35) phases. Experimental diets were designed with three different xylanase activity levels (1,500 U/kg, 3,000 U/kg, 4,500 U/kg) in the NC-3 diet to estimate the compensation level of ME. Xylanase activity levels were pre-determined based on a previous study (*in vitro* experiment, unpublished data). All diets contained 0.3% Cr₂O₃ (chromic oxide powder, Daejung Chemicals & Materials, Siheung, Korea) as an index for digestibility analysis on days 24 and 35.

Data collection

Growth performance parameters were measured based on a single pen. BW and feed intake were measured on days 7, 14, 21, 24, 28, and 35. Average daily gain (ADG), average daily feed intake (ADFI), and feed conversion ratio were calculated based on measured BW and feed intake every week. Mortality was measured when the dead bird was observed during the experimental period.

Sacrifice bird and sample collection were carried out on days 24 and 35 to estimate the effect of starter and grower phase diets. One bird per cage closer to the median BW of the treatment was selected, stunned by carbon dioxide, and euthanized by cervical dislocation for sample collection.

Meat samples were collected after the bird was sacrificed. Both sides of the pectoralis major and pectoralis minor were collected for breast muscle samples and the right-side drumstick with thigh was collected for leg muscle samples. Dry matter, crude protein, ether extract, and ash were analyzed with the proximate analysis method suggested by AOAC [17].

Ileal digesta samples were collected for analysis of viscosity and ileal nutrient digestibility. Fresh digesta 2 g was collected from digesta and centrifuged (12,000×g, 10 min, 20°C). Supernatant 0.5 mL was used for determining viscosity using the viscometer (Brookfield DV-III model, Brookfield Engineering Laboratories, Stoughton, MA, USA) at 25°C with a CP40 cone and shear rate of 5–500/s. Before the proximate analysis, digesta are oven-dried at 55°C for 24 h, followed by fine grinding and strained through a sieve of < 0.75 mm. Dry matter, crude protein, ether extract, crude ash, and total gross energy were analyzed with the proximate analysis method suggested by AOAC [17].

Table 2. Calculated and analyzed nutrient content and ingredient profiles of the 4 basal diets for the starter phase (as-fed basis, %)

| Ingredient | Diet ¹⁾ | | | |
|--------------------------------|--------------------|---------------------|-------|-------|
| | Control | NC-1 | NC-2 | NC-3 |
| Corn | 57.46 | 56.59 | 55.71 | 54.84 |
| Wheat | 4.00 | 5.33 | 6.67 | 8.00 |
| Soybean meal (48 %) | 30.25 | 29.89 | 29.54 | 29.18 |
| Fish meal | 1.90 | 1.97 | 2.03 | 2.10 |
| Vegetable oil | 2.35 | 2.10 | 1.85 | 1.60 |
| Limestone | 1.08 | 1.17 | 1.27 | 1.36 |
| Mono-calcium phosphate | 1.52 | 1.51 | 1.51 | 1.50 |
| Salt | 0.30 | 0.30 | 0.30 | 0.30 |
| Vit-Min premix ²⁾ | 0.30 | 0.30 | 0.30 | 0.30 |
| Lys-HCl | 0.25 | 0.25 | 0.25 | 0.25 |
| DL-Methionine | 0.29 | 0.28 | 0.28 | 0.27 |
| Cr ₂ O ₃ | 0.30 | 0.30 | 0.30 | 0.30 |
| Calculated composition | | | | |
| Metabolizable energy (kcal/kg) | 3,050 | 3,010 | 2,970 | 2,930 |
| Crude protein (%) | 21.50 | 21.50 | 21.50 | 21.50 |
| Lysine (%) | 1.32 | 1.32 | 1.32 | 1.32 |
| Methionine + Cystine (%) | 1.00 | 1.00 | 0.99 | 0.99 |
| Analyzed xylanase activity | None | NCX-1 ³⁾ | NCX-2 | NCX-3 |
| Xylanase activity (U/g) | - | 1.49 | 3.00 | 4.52 |

¹⁾NC-1, -40 kcal/kg ME reduced from PC diet; NC-2, -80 kcal/kg ME reduced from PC diet; NC-3, -120 kcal/kg ME reduced from PC diet; PC, energy sufficient diet.

²⁾Vitamin and mineral mixture provided the following nutrients per kg of diet: vitamin A, 24,000 IU; vitamin D3, 6,000 IU; vitamin E, 30 IU; vitamin K, 4 mg; thiamin, 4 mg; riboflavin, 12 mg; pyridoxine, 4 mg; folacine, 2 mg; biotin, 0.03 mg; vitamin B8 0.06 mg; niacin, 90 mg; pantothenic acid, 30 mg; Fe, 80 mg (as FeSO₄ · H₂O); Zn, 80 mg (as ZnSO₄ · H₂O); Mn, 80 mg (as MnSO₄ · H₂O); Co, 0.5 mg (as CoSO₄ · H₂O); Cu, 10 mg (as CuSO₄ · H₂O); Se, 0.2 mg (as Na₂SeO₃); I, 0.9 mg (as Ca (IO₃) · 2H₂O).

³⁾NCX-1, NC-3 diet with 1,500 U/kg xylanase; NCX-2, NC-3 diet with 3,000 U/kg xylanase; NCX-3, NC-3 diet with 4,500 U/kg xylanase.

NC, negative control; ME, metabolizable energy; PC, positive control.

Statistical analysis

Data were analyzed as a completely randomized design, using one-way analysis of variance in SPSS software (Version 26, IBM SPSS, Chicago, IL, USA, 2018). A pen is used as the experimental unit for growth performances and the individual bird is used as the experimental unit for digestibility, carcass traits, and viscosity. Tukey's multiple range test was used to compare the significant differences between different pairs of means at ($p < 0.05$) when the data showed a significant difference on ANOVA.

Xylanase assessment

The xylanase was assessed to estimate the compensation amount of ME in an energy-deficiency diet and to evaluate the ideal levels of xylanase in the diet.

The matrix value was estimated following the method suggested by Adedokun et al. [18]. Orthogonal polynomial contrasts examine responses to ME levels for BW and also xylanase for BW. Linear and quadratic regression were analyzed for xylanase with BW and ME level with BW. The regression equation for the ME levels in the diet and supplemental xylanase level for particular

Table 3. Calculated and analyzed nutrient content and ingredient profiles of the 4 basal diets for the finisher phase (as-fed basis, %)

| Ingredient | Diet ¹⁾ | | | |
|--------------------------------|--------------------|---------------------|-------|-------|
| | PC | NC-1 | NC-2 | NC-3 |
| Corn | 51.35 | 52.68 | 54.02 | 55.35 |
| Wheat | 4.00 | 4.33 | 4.67 | 5.00 |
| Soybean meal (48 %) | 27.87 | 27.60 | 27.34 | 27.07 |
| Fish meal | 2.20 | 2.13 | 2.07 | 2.00 |
| Vegetable oil | 3.30 | 2.62 | 1.94 | 1.26 |
| Corn starch | 7.00 | 6.39 | 5.77 | 5.16 |
| Limestone | 1.38 | 1.33 | 1.28 | 1.23 |
| Mono-calcium phosphate | 1.50 | 1.50 | 1.50 | 1.50 |
| Salt | 0.30 | 0.30 | 0.30 | 0.30 |
| Vit-Min premix ²⁾ | 0.30 | 0.30 | 0.30 | 0.30 |
| Lys-HCl | 0.20 | 0.21 | 0.22 | 0.23 |
| DL-methionine | 0.30 | 0.30 | 0.30 | 0.30 |
| Cr ₂ O ₃ | 0.30 | 0.30 | 0.30 | 0.30 |
| Calculated composition | | | | |
| Metabolizable energy (kcal/kg) | 3,150 | 3,110 | 3,070 | 3,030 |
| Crude protein (%) | 20.00 | 20.00 | 20.00 | 20.00 |
| Lysine (%) | 1.22 | 1.22 | 1.22 | 1.22 |
| Methionine + cystine (%) | 0.96 | 0.96 | 0.97 | 0.97 |
| Analyzed xylanase activity | None | NCX-1 ³⁾ | NCX-2 | NCX-3 |
| Xylanase activity (U/g) | - | 1.52 | 3.03 | 4.49 |

¹⁾PC, energy sufficient diet; NC-1, -40 kcal/kg ME reduced from PC diet; NC-2, -80 kcal/kg ME reduced from PC diet; NC-3, -120 kcal/kg ME reduced from PC diet.

²⁾Vitamin and mineral mixture provided the following nutrients per kg of diet: vitamin A, 24,000 IU; vitamin D3, 6,000 IU; vitamin E, 30 IU; vitamin K, 4 mg; thiamin, 4 mg; riboflavin, 12 mg; pyridoxine, 4 mg; folacin, 2 mg; biotin, 0.03 mg; vitamin B8 0.06 mg; niacin, 90 mg; pantothenic acid, 30 mg; Fe, 80 mg (as FeSO₄ · H₂O); Zn, 80 mg (as ZnSO₄ · H₂O); Mn, 80 mg (as MnSO₄ · H₂O); Co, 0.5 mg (as CoSO₄ · H₂O); Cu, 10 mg (as CuSO₄ · H₂O); Se, 0.2 mg (as Na₂SeO₃); I, 0.9 mg (as Ca (IO₃) · 2H₂O).

³⁾NCX-1, NC-3 diet with 1,500 U/kg xylanase; NCX-2, NC-3 diet with 3,000 U/kg xylanase; NCX-3, NC-3 diet with 4,500 U/kg xylanase.

response variables were equated and solved for x.

$$\text{Equation 1: } Y_m = a_m + b_m x_m \text{ (linear regression equation for ME)}$$

$$\text{Equation 2: } Y_x = a_x + b_x x_x \text{ (linear regression equation for xylanase)}$$

$$\text{Equation 1} = \text{Equation 2: } a_m + b_m x = a_x + b_x x_x$$

where Y represents the response criterion that is BW, x_m presents ME levels in diets, b_m represents the slope of the response criterion to the dietary energy, and b_x represents the slope of the response criterion to the added xylanase. The linear response equations for ME in diet and that for added xylanase were set to be equal and were solved for BW equivalency values for their respective variable.

The quadratic regression equation is also calculated the same as the linear equation as follows:

$$\text{Equation 3: } Y_m = a_m + b_m x_m + c_m x_m^2 \text{ (quadratic regression equation for ME)}$$

$$\text{Equation 4: } Y_x = a_x + b_x x_x + c_x x_x^2 \text{ (quadratic regression equation for xylanase)}$$

$$\text{Equation 3} = \text{Equation 4: } a_m + b_m x_m + c_m x_m^2 = a_x + b_x x_x + c_x x_x^2$$

Matrix values calculated by linear and quadratic regression are compared to check the accuracy of each method.

The marginal xylanase level was estimated by the methods suggested by [19,20]. The linear and quadratic plateau (broken line) models were used to determine the optimum requirement for dietary xylanase levels. The plateau model consisted of a straight or curvy line with an increasing or decreasing slope and a horizontal line.

The linear model of the one-slope broken line is as follows:

$$Y = a + b (R-x)$$

In these equations, a is the ordinate, R is the abscissa of the breakpoint in the curve, and b is the slope of the line for $x < R$. When $x > R$, the equation indicates $Y = a$.

The quadratic plateau model is as follows:

$$Y = a + bx + cx^2$$

In the quadratic plateau model, the vertex was regarded as a breakpoint, and x showed over the vertex point, Y was presented as a constant number that the vertex value substituted in the formula. In both models, X and Y indicate the xylanase level and BW. Based on the measured growth performance.

RESULTS

Experimental birds were consuming sufficient drinking water and experimental feed, and no symptoms of death or disease were found due to sudden death syndrome (SDS) or stress throughout the entire 35-day experiment.

Growth performance

BW, ADFI, average daily gain, and feed conversion ratio from days 7 to 35 were shown in Tables 4 to 7. BW on days 14, 24, and 28 showed significant differences ($p < 0.05$) during the experimental period. Xylanase treatments except for NCX-1 performed a superior BW compared to NC-3 when there was a significant difference ($p < 0.05$). NCX-1 exhibits a significant difference ($p < 0.05$) with NC-3 only on day 24. During all experimental periods, xylanase treatment showed no differences from the PC treatment ($p > 0.05$). Xylanase treatments showed improved ADG ($p < 0.05$) on week 2 compared to NC-3 and there were no differences compared to the PC diet. In the starter, grower,

Table 4. Effect of xylanase addition on the body weight of broilers from day 8 to 35 (g)

| Treatment | NC-1 ¹⁾ | NC-2 | NC-3 | NCX-1 | NCX-2 | NCX-3 | PC | SEM | p-value |
|-----------|-------------------------|------------------------|-----------------------|-------------------------|------------------------|-----------------------|-----------------------|--------|---------|
| Day 8 | 134.24 | 133.63 | 134.7 | 133.89 | 134.06 | 134.43 | 134.96 | 0.952 | 0.827 |
| Day 14 | 348.29 ^{ab} | 348.86 ^{ab} | 331.95 ^a | 346.04 ^{ab} | 359.21 ^b | 363.96 ^b | 359.88 ^b | 8.285 | 0.004 |
| Day 21 | 635.64 | 618.30 | 626.55 | 651.68 | 675.88 | 672.70 | 657.13 | 21.989 | 0.068 |
| Day 24 | 838.72 ^{ab} | 808.66 ^a | 809.55 ^a | 843.10 ^b | 867.22 ^b | 873.48 ^b | 862.91 ^b | 25.694 | 0.036 |
| Day 28 | 1,180.82 ^{abc} | 1,129.82 ^{ab} | 1,120.54 ^a | 1,168.33 ^{abc} | 1,192.33 ^{bc} | 1,211.19 ^c | 1,207.28 ^c | 34.989 | 0.037 |
| Day 35 | 1,703.23 | 1,673.24 | 1,659.35 | 1,726.64 | 1,730.48 | 1,750.60 | 1,725.47 | 52.052 | 0.494 |

¹⁾NC-1, -40 kcal/kg ME reduced from PC diet; NC-2, -80 kcal/kg ME reduced from PC diet; NC-3, -120 kcal/kg ME reduced from PC diet; NCX-1, NC-3 diet with 1,500 U/kg xylanase; NCX-2, NC-3 diet with 3,000 U/kg xylanase; NCX-3, NC-3 diet with 4,500 U/kg xylanase; PC, energy sufficient diet.

^{a-c} Values in a row with different superscripts differ significantly ($p < 0.05$).

Table 5. Effect of xylanase addition on the average daily gain of broilers from day 8 to 35 (g/d)

| Treatment | NC-1 ¹⁾ | NC-2 | NC-3 | NCX-1 | NCX-2 | NCX-3 | PC | SEM | p-value |
|-----------|---------------------|--------------------|--------------------|---------------------|--------------------|--------------------|--------------------|-------|---------|
| Day 8–14 | 30.72 ^{ab} | 30.96 ^b | 28.46 ^a | 30.31 ^{ab} | 32.16 ^b | 32.7 ^{9b} | 32.13 ^b | 1.146 | 0.007 |
| Day 15–21 | 41.05 | 38.49 | 42.09 | 43.66 | 45.24 | 44.11 | 42.46 | 2.789 | 0.279 |
| Day 22–24 | 67.69 | 62.46 | 59.67 | 63.81 | 63.78 | 66.93 | 68.59 | 3.425 | 0.124 |
| Day 25–28 | 85.53 | 80.04 | 77.25 | 81.31 | 81.28 | 84.43 | 86.09 | 4.431 | 0.118 |
| Day 29–35 | 75.06 | 77.77 | 76.97 | 79.76 | 76.88 | 77.06 | 74.03 | 4.430 | 0.909 |
| Day 8–24 | 42.26 | 40.47 | 40.64 | 41.72 | 43.13 | 43.47 | 42.82 | 1.500 | 0.292 |
| Day 25–35 | 79.68 | 80.05 | 78.71 | 80.32 | 78.48 | 79.74 | 78.41 | 3.370 | 0.995 |
| Day 8–35 | 56.80 | 56.02 | 56.88 | 56.96 | 57.02 | 57.72 | 55.59 | 1.851 | 0.942 |

¹⁾NC-1, -40 kcal/kg ME reduced from PC diet; NC-2, -80 kcal/kg ME reduced from PC diet; NC-3, -120 kcal/kg ME reduced from PC diet; NCX-1, NC-3 diet with 1,500 U/kg xylanase; NCX-2, NC-3 diet with 3,000 U/kg xylanase; NCX-3, NC-3 diet with 4,500 U/kg xylanase; PC, energy sufficient diet.

^{ab}Values in a row with different superscripts differ significantly ($p < 0.05$).

Table 6. Effect of xylanase addition on the average daily feed intake of broilers from day 8 to 35 (g/d)

| Treatment | NC-1 ¹⁾ | NC-2 | NC-3 | NCX-1 | NCX-2 | NCX-3 | PC | SEM | p-value |
|-----------|--------------------|--------|--------|--------|--------|--------|-------|--------|---------|
| Day 8–14 | 38.82 | 38.22 | 38.31 | 39.03 | 40.80 | 41.37 | 38.40 | 0.952 | 0.498 |
| Day 15–21 | 66.40 | 64.86 | 69.80 | 72.78 | 72.59 | 71.98 | 63.25 | 8.285 | 0.472 |
| Day 22–24 | 90.67 | 86.06 | 83.20 | 82.95 | 84.73 | 88.12 | 83.76 | 21.989 | 0.424 |
| Day 25–28 | 110.42 | 104.17 | 104.55 | 103.83 | 103.67 | 108.5 | 99.71 | 25.694 | 0.378 |
| Day 29–35 | 108.25 | 111.17 | 112.88 | 111.65 | 111.05 | 111.65 | 97.77 | 34.989 | 0.620 |
| Day 8–24 | 59.33 | 57.63 | 59.20 | 60.67 | 61.64 | 62.22 | 56.64 | 3.026 | 0.490 |
| Day 25–35 | 109.04 | 108.63 | 109.85 | 108.81 | 108.36 | 110.50 | 98.47 | 6.445 | 0.560 |
| Day 8–35 | 78.86 | 77.66 | 79.10 | 79.58 | 80.00 | 81.19 | 73.07 | 3.765 | 0.451 |

¹⁾NC-1, -40 kcal/kg ME reduced from PC diet; NC-2, -80 kcal/kg ME reduced from PC diet; NC-3, -120 kcal/kg ME reduced from PC diet; NCX-1, NC-3 diet with 1,500 U/kg xylanase; NCX-2, NC-3 diet with 3,000 U/kg xylanase; NCX-3, NC-3 diet with 4,500 U/kg xylanase; PC, energy sufficient diet.

Table 7. Effect of xylanase addition on the feed conversion ratio of broilers from day 8 to 35 (g/g)

| Treatment | NC-1 ¹⁾ | NC-2 | NC-3 | NCX-1 | NCX-2 | NCX-3 | PC | SEM | p-value |
|-----------|--------------------|------|------|-------|-------|-------|------|-------|---------|
| Day 8–14 | 1.26 | 1.22 | 1.30 | 1.32 | 1.27 | 1.27 | 1.20 | 0.077 | 0.737 |
| Day 15–21 | 1.60 | 1.64 | 1.66 | 1.66 | 1.60 | 1.64 | 1.51 | 0.107 | 0.809 |
| Day 22–24 | 1.31 | 1.36 | 1.37 | 1.33 | 1.34 | 1.34 | 1.24 | 0.081 | 0.750 |
| Day 25–28 | 1.28 | 1.29 | 1.33 | 1.30 | 1.28 | 1.30 | 1.17 | 0.082 | 0.575 |
| Day 29–35 | 1.45 | 1.44 | 1.48 | 1.40 | 1.44 | 1.46 | 1.34 | 0.089 | 0.757 |
| Day 8–24 | 1.41 | 1.43 | 1.46 | 1.46 | 1.43 | 1.44 | 1.33 | 0.071 | 0.596 |
| Day 25–35 | 1.37 | 1.37 | 1.40 | 1.36 | 1.38 | 1.40 | 1.27 | 0.082 | 0.719 |
| Day 8–35 | 1.39 | 1.39 | 1.43 | 1.41 | 1.40 | 1.41 | 1.30 | 0.072 | 0.610 |

¹⁾NC-1, -40 kcal/kg ME reduced from PC diet; NC-2, -80 kcal/kg ME reduced from PC diet; NC-3, -120 kcal/kg ME reduced from PC diet; NCX-1, NC-3 diet with 1,500 U/kg xylanase; NCX-2, NC-3 diet with 3,000 U/kg xylanase; NCX-3, NC-3 diet with 4,500 U/kg xylanase; PC, energy sufficient diet.

and whole experimental periods, there were no differences in ADG, ADFI, and feed conversion ratio among the treatments ($p > 0.05$).

Carcass traits

Proximate analysis results of breast and leg meat on days 24 and 35 are presented in Tables 8 and 9. There was a significant difference ($p < 0.05$) in the crude protein on leg meat and dry matter content

Table 8. The effect of xylanase addition in the poultry diet on leg meat (%)

| Treatment | NC-1 ¹⁾ | NC-2 | NC-3 | NCX-1 | NCX-2 | NCX-3 | PC | SEM | <i>p</i> -value |
|---------------|---------------------|---------------------|---------------------|--------------------|---------------------|--------------------|---------------------|-------|-----------------|
| Day 24 | | | | | | | | | |
| Moisture | 71.71 | 72.28 | 70.34 | 72.31 | 72.14 | 74.38 | 71.80 | 1.355 | 0.208 |
| Crude protein | 19.44 ^{ab} | 19.95 ^{ab} | 19.52 ^{ab} | 20.57 ^b | 19.65 ^{ab} | 18.93 ^a | 19.75 ^{ab} | 0.347 | 0.006 |
| Crude fat | 5.26 | 5.75 | 5.47 | 4.56 | 6.15 | 5.39 | 5.51 | 0.919 | 0.756 |
| Ash | 1.23 | 1.26 | 1.33 | 1.31 | 1.26 | 1.15 | 1.22 | 0.064 | 0.158 |
| Day 35 | | | | | | | | | |
| Moisture | 70.06 | 68.86 | 68.77 | 69.56 | 65.88 | 67.36 | 66.57 | 1.669 | 0.155 |
| Crude protein | 19.64 | 19.47 | 19.77 | 18.73 | 19.04 | 18.62 | 19.32 | 0.769 | 0.674 |
| Crude fat | 5.03 | 5.44 | 6.39 | 4.29 | 5.65 | 4.14 | 6.95 | 1.517 | 0.945 |
| Ash | 1.26 | 1.30 | 1.19 | 1.17 | 1.21 | 1.20 | 1.14 | 0.071 | 0.320 |

¹⁾NC-1, -40 kcal/kg ME reduced from PC diet; NC-2, -80 kcal/kg ME reduced from PC diet; NC-3, -120 kcal/kg ME reduced from PC diet; NCX-1, NC-3 diet with 1,500 U/kg xylanase; NCX-2, NC-3 diet with 3,000 U/kg xylanase; NCX-3, NC-3 diet with 4,500 U/kg xylanase; PC, energy sufficient diet.

^{a,b}Values in a row with different superscripts differ significantly ($p < 0.05$).

Table 9. The effect of xylanase addition in the poultry diet on breast meat (%)

| Treatment | NC-1 ¹⁾ | NC-2 | NC-3 | NCX-1 | NCX-2 | NCX-3 | PC | SEM | <i>p</i> -value |
|---------------|--------------------|---------------------|--------------------|---------------------|---------------------|---------------------|--------------------|-------|-----------------|
| Day 24 | | | | | | | | | |
| Moisture | 73.78 ^c | 72.64 ^{bc} | 68.23 ^a | 68.57 ^{ab} | 69.23 ^{ab} | 68.30 ^{ab} | 66.92 ^a | 1.353 | < 0.001 |
| Crude protein | 24.08 | 24.30 | 23.94 | 24.31 | 24.52 | 24.02 | 24.86 | 0.471 | 0.498 |
| Crude fat | 1.32 | 1.25 | 1.07 | 0.91 | 0.98 | 0.67 | 1.06 | 0.212 | 0.092 |
| Ash | 1.60 ^{ab} | 1.53 ^{ab} | 1.68 ^{ab} | 1.79 ^b | 1.72 ^{ab} | 1.75 ^{ab} | 1.42 ^a | 0.105 | 0.023 |
| Day 35 | | | | | | | | | |
| Moisture | 68.03 | 69.18 | 67.10 | 67.98 | 66.28 | 66.36 | 66.74 | 0.991 | 0.074 |
| Crude protein | 21.84 | 23.70 | 23.75 | 23.46 | 24.50 | 23.71 | 23.80 | 0.894 | 0.179 |
| Crude fat | 1.18 | 0.75 | 0.59 | 1.13 | 0.77 | 0.63 | 1.07 | 0.281 | 0.223 |
| Ash | 1.66 | 1.59 | 1.60 | 1.63 | 1.63 | 1.67 | 1.64 | 0.064 | 0.876 |

¹⁾NC-1, -40 kcal/kg ME reduced from PC diet; NC-2, -80 kcal/kg ME reduced from PC diet; NC-3, -120 kcal/kg ME reduced from PC diet; NCX-1, NC-3 diet with 1,500 U/kg xylanase; NCX-2, NC-3 diet with 3,000 U/kg xylanase; NCX-3, NC-3 diet with 4,500 U/kg xylanase; PC, energy sufficient diet.

^{a-c} Values in a row with different superscripts differ significantly ($p < 0.05$).

and crude ash on breast meat on day 24. However, xylanase treatments showed no significant differences compared to NC-3 treatment and PC treatment ($p > 0.05$). Furthermore, carcass traits on day 35 have no difference for all measurements ($p > 0.05$).

Viscosity and digestibility

Digesta viscosity and crude protein analysis were presented in Table 10. Xylanase treatment showed significant differences ($p < 0.05$) compared to NC-3 and PC diet on days 24 and 35. Xylanase treatments on ME, DM, and crude protein (CP) showed a significant difference ($p < 0.05$) and performed the same with the PC diet.

Matrix value evaluation

The matrix value estimation with linear and quadratic regression graphs is presented in Tables 11 and 12. The linear regression showed improved matrix value based on the xylanase in the diet increases. On day 14, the matrix value for 1,500 U/kg xylanase was 60.54 kcal/kg, 3,000 U/kg xylanase was 117.12 kcal/kg, and 4,500 U/kg xylanase was 137.53 kcal/kg. On day 24, the matrix

Table 10. Digesta analysis for viscosity and protein digestibility

| Treatment | NC-1 ¹⁾ | NC-2 | NC-3 | NCX-1 | NCX-2 | NCX-3 | PC | SEM | p-value |
|---------------------------|---------------------|--------------------|--------------------|---------------------|---------------------|---------------------|---------------------|-------|---------|
| Viscosity (mPa/s) | | | | | | | | | |
| Day 24 | 2.76 ^{cde} | 2.84 ^{de} | 2.93 ^e | 2.65a ^{bc} | 2.55 ^{ab} | 2.48 ^a | 2.73 ^{bcd} | 0.061 | 0.001 |
| Day 35 | 2.64 ^{cd} | 2.78 ^{de} | 2.86 ^e | 2.56a ^{bc} | 2.44 ^{ab} | 2.38 ^a | 2.62 ^{bcd} | 0.062 | 0.001 |
| Protein digestibility (%) | | | | | | | | | |
| Day 24 | 81.87 ^b | 81.87 ^b | 77.22 ^a | 79.62 ^{ab} | 80.52 ^{ab} | 81.52 ^b | 81.47 ^b | 0.003 | 0.011 |
| Day 35 | 77.62 ^{ab} | 77.8 ^{ab} | 76.90 ^a | 82.12 ^b | 80.75 ^{ab} | 80.75 ^{ab} | 82.02 ^{ab} | 0.009 | 0.016 |

¹⁾NC-1, -40 kcal/kg ME reduced from PC diet; NC-2, -80 kcal/kg ME reduced from PC diet; NC-3, -120 kcal/kg ME reduced from PC diet; NCX-1, NC-3 diet with 1,500 U/kg xylanase; NCX-2, NC-3 diet with 3,000 U/kg xylanase; NCX-3, NC-3 diet with 4,500 U/kg xylanase; PC, energy sufficient diet.

^{a-e} Values in a row with different superscripts differ significantly ($p < 0.05$).

Table 11. Xylanase matrix value evaluation by the linear regression graph

| Items | Linear regression ¹⁾ | R ² | Body weight (g) | | | Calculated matrix value ²⁾ (kcal/kg) | | |
|--------|---------------------------------|----------------|-----------------|------------|------------|---|------------|------------|
| | | | 1,500 U/kg | 3,000 U/kg | 4,500 U/kg | 1,500 U/kg | 3,000 U/kg | 4,500 U/kg |
| Day 14 | 331.95 + 0.23275x | 87.162 | 346.04 | 359.21 | 363.96 | 60.54 | 117.12 | 137.53 |
| Day 24 | 809.55 + 0.44467x | 88.929 | 843.10 | 867.22 | 873.48 | 75.45 | 129.69 | 143.77 |
| Day 28 | 1,120.54 + 0.72283x | 94.289 | 1,168.33 | 1,192.33 | 1,211.19 | 66.11 | 99.32 | 125.41 |

¹⁾Regression was drawn with body weight (x) and growth performance (y); x has a range from 0 to 120 kcal/kg, 0 represents the ME level of the NC-3 diet and 120 represents the ME level of the PC diet for convenience.

²⁾Calculated values are calculated by substituting body weight values to estimate the matrix value of xylanase. Estimated values over 120 kcal/kg could be biased. ME, metabolizable energy; PC, energy sufficient diet.

Table 12. Xylanase matrix value evaluation by the quadratic regression graph

| Items | Quadratic regression ¹⁾ | R ² | Body weight (g) | | | Calculated matrix value ²⁾ (kcal/kg) | | |
|--------|--|----------------|-----------------|------------|------------|---|------------|------------|
| | | | 1,500 U/kg | 3,000 U/kg | 4,500 U/kg | 1,500 U/kg | 3,000 U/kg | 4,500 U/kg |
| Day 14 | -0.0008x ² + 0.3078x + 333.43 | 88.943 | 346.04 | 359.21 | 363.96 | 46.62 | 123.22 | 192.38 |
| Day 24 | 0.0039x ² + 0.0051x + 807.71 | 96.665 | 843.10 | 867.22 | 873.48 | 94.61 | 122.88 | 129.21 |
| Day 28 | 0.0027x ² + 0.4559x + 1,117.2 | 95.726 | 1,168.33 | 1,192.33 | 1,211.19 | 77.02 | 102.53 | 120.36 |

¹⁾Regression was drawn with body weight (x) and growth performance (y); x has a range from 0 to 120 kcal/kg, 0 represents the ME level of the NC-3 diet and 120 represents the ME level of the PC diet for convenience.

²⁾Calculated values are calculated by substituting body weight values to estimate the matrix value of xylanase. Estimated values over 120 kcal/kg could be biased. ME, metabolizable energy; PC, energy sufficient diet.


value for 1,500 U/kg xylanase was 75.45 kcal/kg, 3,000 U/kg xylanase was 129.69 kcal/kg, and 4,500 U/kg xylanase was 143.77 kcal/kg. On day 28, the matrix value for 1,500 U/kg xylanase was 66.11 kcal/kg, 3,000 U/kg xylanase was 99.32 kcal/kg, and 4,500 U/kg xylanase was 125.41 kcal/kg.

The quadratic regression also showed improved matrix value based on the xylanase increases. On day 14, the matrix value for 1,500 U/kg xylanase was 46.62 kcal/kg, 3,000 U/kg xylanase was 123.22 kcal/kg, and 4,500 U/kg xylanase was 192.38 kcal/kg. On day 24, the matrix value for 1,500 U/kg xylanase was 94.61 kcal/kg, 3,000 U/kg xylanase was 122.88 kcal/kg, and 4,500 U/kg xylanase was 129.21 kcal/kg. On day 28, the matrix value for 1,500 U/kg xylanase was 77.02 kcal/kg, 3,000 U/kg xylanase was 102.53 kcal/kg, and 4,500 U/kg xylanase was 120.36 kcal/kg.

Marginal xylanase amount estimation

Marginal xylanase amount estimation analyzed by linear and quadratic plateau methods was drawn in Figs 1, 2, and 3. The linear plateau indicates the xylanase requirement of 3,500 U/kg


Linear and quadratic plateau analysis on day 14



| | Requirement (U/kg) | Standard error | p-value | R ² |
|-------------------|--------------------|----------------|---------|----------------|
| Linear plateau | 3,500 | 1.681 | 0.003 | 0.99545 |
| Quadratic plateau | 4,400 | 3.319 | 0.006 | 0.98117 |

Fig. 1. Linear and quadratic plateau analysis to evaluate the marginal xylanase level on day 14.

Linear and quadratic plateau analysis on day 24



| | Requirement (U/kg) | Standard error | p-value | R ² |
|-------------------|--------------------|----------------|---------|----------------|
| Linear plateau | 3,244 | 3.850 | 0.003 | 0.99412 |
| Quadratic plateau | 3,517 | 6.586 | 0.005 | 0.97741 |

Fig. 2. Linear and quadratic plateau analysis to evaluate the marginal xylanase level on day 24.

and the quadratic plateau indicates the xylanase requirement of 4,400 U/kg on day 14. The linear plateau indicates the xylanase requirement of 3,244 U/kg and the quadratic plateau indicates the xylanase requirement of 3,517 U/kg on day 24. On day 28, The linear plateau indicates the xylanase requirement of 3,622 U/kg and the quadratic plateau indicates the xylanase requirement of 4,000 U/kg.

DISCUSSION

Many previous studies suggest that the xylanase addition in adequate ME level diets conditioned




Fig. 3. Linear and quadratic plateau analysis to evaluate the marginal xylanase level on day 28.

with various feed ingredients also presents that xylanase addition also improves growth performances, digestibility, and viscosity [10–12,21,22]. The water-soluble arabinoxylans are known as anti-nutritional factors by increasing viscosity and encapsulating the nutrient [23]. The mode of action of xylanase is known that xylanase degrades the arabinoxylan backbone and releases the trapped nutrient. Releasing entrapped nutrients increases the amount of absorption in the small intestine [5,8]. Broken-down arabinoxylans make the viscosity of digesta lower and it allows the digesta to mix properly and allows the nutrient to be absorbed more easily [24]. Furthermore, small oligosaccharides produced by degrading the arabinoxylan also showed potential prebiotic effects [22,25]. The viscosity result that a xylanase-added diet exhibits a significantly low viscosity and a higher xylanase-added diet showed a significantly lower viscosity was also supported by previous research.

The exogenous NSP enzymes have been added to improve the nutrient value by degrading the unavailable nutrient complexes and making them available nutrients to the host animal. Especially the absence of xylanase in the poultry intestine makes endogenous xylanase addition in broiler diet experiments conducted continuously. The growth performances of the birds fed xylanase added in energy deficiency diet in the current experiment performed significantly equal to the positive control diet which meets or exceeds the nutrient requirement. Few earlier studies suggest that supplemented enzymes such as xylanase, glucanase, and protease do not affect broiler performances [26–28]. Also, some other research suggests a corn-based diet with xylanase showed no significant difference because corn contains less than 1 g/kg water-soluble NSP [29]. However, those results may be caused by various factors which nutrient levels, diet formulation, enzyme dosage, and bird conditions [30]. On the other hand, numerous studies suggest that exogenous xylanase or NSP enzyme complex addition in a broiler diet could enhance growth performances and ileal digestibility in energy deficiency conditions [31,32]. Results of xylanase addition in energy deficiency diet propose that the xylanase addition in poultry diet could compensate for the ME by degrading the xylan complex or releasing the encapsulated nutrient [5,8] and following results, the lack of ME affect negatively during the whole experimental period and it agreed with the previous xylanase experiment with a reduced ME diet [31]. This experiment also exhibits that exogenous

xylanase addition could improve growth performances, viscosity, and digestibility compared to the control diet. Therefore, previous studies' results and this experimental result suggest that the birds fed xylanase addition with a calculated energy deficiency diet could be performed equally with birds fed an adequate energy level diet. These results also suggest that the xylanase addition could be a good method to save production costs by reducing ME.

The effect of xylanase on poultry meat is controversial with various contrast studies. Some studies suggest that the xylanase addition can improve not only BW gain or final BW but also breast and leg meat weight [33,34], however, another study suggests the xylanase addition only improves growth performance but carcass traits such as breast muscle yield and relative weight of the abdominal fat pad [35]. These results could be explained that the xylanase addition on the adequate nutrient-contained diet only showed growth performance differences but some basal diet effect on the negative effect so xylanase addition compensates for enough nutrients.

The exact measurement of the amount of xylanase requirement and abilities is very important. Nevertheless, Various previous studies suggest that xylanase could improve performance, but the estimated proper xylanase activities showed diverse levels from 1,250 to 30,000 U/kg [1,11,36]. Those big variations of recommended xylanase activity might be caused by the various families of xylanase. Xylanases are classified by displaying varying folds, mechanisms of action, specifications of substrates, different rates, yields, and production of hydrolytic activities, and physicochemical characteristics [37]. Therefore, the individual xylanase requirements should be measured for maximizing animal production and for minimizing enzyme wastage. The methodology for evaluating the optimal level of xylanase follows the previous enzyme study about calculating the equivalent level with a small revision [18]. The compensation level of ME and optimal xylanase level showing the maximum compensation level was evaluated by comparing the regression of growth performance and ME and the regression of growth performance and xylanase levels. The xylanase matrix value results which 120 kcal compensation on 4,500 U/kg agreed with the previous *in vivo* experiment to estimate the matrix value. The marginal xylanase level was estimated with methods suggested by [19,20] with a small revision. The linear and quadratic regression formula with a break-point represented the maximum activity level of xylanase and performances. The linear and quadratic plateau regression presents the maximum xylanase of 3,622 U/kg in the linear plateau and the maximum xylanase of 4,000 U/kg in the quadratic plateau on day 28. The compensation levels of experimental results are acceptable when compared to the previous study which added xylanase to the energy deficiency diet [31]. The referred research also suggests that the addition of xylanase in a broiler diet could improve BW and feed conversion ratio.

CONCLUSION

Xylanase addition in the ME deficiency diet could improve growth performance, reduce viscosity, and enhance protein digestibility as much as the common diet by degrading arabinoxylans in the diet. The maximum level of xylanase in the diet was calculated as 4,500 U/kg with 120 kcal/kg ME compensation. The marginal levels of xylanase were estimated as 3,622 U/kg with the linear plateau method and estimated as 4,000 U/kg with the quadratic plateau method. These findings suggest that xylanase supplementation could be used for broilers in a ME deficiency diet.

REFERENCES

1. Van Hoeck V, Papadopoulos GA, Giannenas I, Lioliopoulou S, Tsiouris V, Mantzios T, et al. New intrinsically thermostable xylanase improves broilers' growth performance, organ weights,

- and affects intestinal viscosity and pH. *Agriculture*. 2021;11:1235. <https://doi.org/10.3390/agriculture11121235>
2. Choct M, Annison G. Anti-nutritive effect of wheat pentosans in broiler chickens: roles of viscosity and gut microflora. *Br Poult Sci*. 1992;33:821-34. <https://doi.org/10.1080/00071669208417524>
 3. Friesen OD, Guenter W, Marquardt RR, Rotter BA. The effect of enzyme supplementation on the apparent metabolizable energy and nutrient digestibilities of wheat, barley, oats, and rye for the young broiler chick. *Poult Sci*. 1992;71:1710-21. <https://doi.org/10.3382/ps.0711710>
 4. Choct M, Hughes RJ, Bedford MR. Effects of a xylanase on individual bird variation, starch digestion throughout the intestine, and ileal and caecal volatile fatty acid production in chickens fed wheat. *Br Poult Sci*. 1999;40:419-22. <https://doi.org/10.1080/00071669987548>
 5. Cowieson AJ. Strategic selection of exogenous enzymes for corn/soy-based poultry diets. *J Poult Sci*. 2010;47:1-7. <https://doi.org/10.2141/jpsa.009045>
 6. Aftab U, Bedford MR. The use of NSP enzymes in poultry nutrition: myths and realities. *Worlds Poult Sci J*. 2018;74:277-86. <https://doi.org/10.1017/S0043933918000272>
 7. Prade RA. Xylanases: from biology to biotechnology. *Biotechnol Genet Eng Rev*. 1996;13:101-32. <https://doi.org/10.1080/02648725.1996.10647925>
 8. Bedford MR. The role of carbohydrases in feedstuff digestion. In: McNab J, Boorman N, editors. *Poultry feedstuffs: supply, composition and nutritive value*. Edinburgh: CAB International; 2002. p. 319-36.
 9. Bedford MR. The evolution and application of enzymes in the animal feed industry: the role of data interpretation. *Br Poult Sci*. 2018;59:486-93. <https://doi.org/10.1080/00071668.2018.1484074>
 10. Hew LI, Ravindran V, Mollah Y, Bryden WL. Influence of exogenous xylanase supplementation on apparent metabolizable energy and amino acid digestibility in wheat for broiler chickens. *Anim Feed Sci Technol*. 1998;75:83-92. [https://doi.org/10.1016/S0377-8401\(98\)00206-5](https://doi.org/10.1016/S0377-8401(98)00206-5)
 11. Kiarie E, Romero LF, Ravindran V. Growth performance, nutrient utilization, and digesta characteristics in broiler chickens fed corn or wheat diets without or with supplemental xylanase. *Poult Sci*. 2014;93:1186-96. <https://doi.org/10.3382/ps.2013-03715>
 12. Rao SVR, Raju MVLN, Nagalakshmi D, Prakash B, Paul SS. Effect of supplementation of graded concentrations of xylanase and α -amylase on performance, slaughter variables, and energy digestibility in broiler chickens fed corn-soybean meal-based diet. *J Appl Poult Res*. 2021;30:100139. <https://doi.org/10.1016/j.japr.2021.100139>
 13. Singh AK, Mishra B, Bedford MR, Jha R. Effects of supplemental xylanase and xylooligosaccharides on production performance and gut health variables of broiler chickens. *J Anim Sci Biotechnol*. 2021;12:98. <https://doi.org/10.1186/s40104-021-00617-8>
 14. Wang J, Cao H, Bao C, Liu Y, Dong B, Wang C, et al. Effects of xylanase in corn- or wheat-based diets on cecal microbiota of broilers. *Front Microbiol*. 2021;12:757066. <https://doi.org/10.3389/fmicb.2021.757066>
 15. Bedford MR, Walk CL, Masey O'Neill HV. Assessing measurements in feed enzyme research: phytase evaluations in broilers. *J Appl Poult Res*. 2016;25:305-14. <https://doi.org/10.3382/japr/pfv073>
 16. Aviagen. *Ros broiler: nutrition specifications*. Huntsville, AL: Aviagen; 2019.
 17. AOAC (Association of Official Analytical Chemists) International. *Official methods of analysis of AOAC International*. 18th ed. Washington, DC: AOAC International; 2005.
 18. Adedokun SA, Sands JS, Adeola O. Determining the equivalent phosphorus released by an

- Escherichia coli-derived phytase in broiler chicks. *Can J Anim Sci.* 2004;84:437-44. <https://doi.org/10.4141/A03-117>
19. Pesti GM, Harper AE, Sunde ML. Choline/methionine nutrition of starting broiler chicks. Three models for estimating the choline requirement with economic considerations. *Poult Sci.* 1980;59:1073-81. <https://doi.org/10.3382/ps.0591073>
 20. Robbins KR. A method, SAS program, and example for fitting the broken-line to growth data. Knoxville, Tennessee: The University of Tennessee Agricultural Experiment Station; 1986. Report No.: 86-09.
 21. Liu N, Ru YJ, Tang DF, Xu TS, Partridge GG. Effects of corn distillers dried grains with solubles and xylanase on growth performance and digestibility of diet components in broilers. *Anim Feed Sci Technol.* 2011;163:260-6. <https://doi.org/10.1016/j.anifeedsci.2010.11.004>
 22. Zhang L, Xu J, Lei L, Jiang Y, Gao F, Zhou GH. Effects of xylanase supplementation on growth performance, nutrient digestibility and non-starch polysaccharide degradation in different sections of the gastrointestinal tract of broilers fed wheat-based diets. *Asian-Australas J Anim Sci.* 2014;27:855-61. <https://doi.org/10.5713/ajas.2014.14006>
 23. Choct M, Hughes RJ, Wang J, Bedford MR, Morgan AJ, Annison G. Increased small intestinal fermentation is partly responsible for the anti-nutritive activity of non-starch polysaccharides in chickens. *Br Poult Sci.* 1996;37:609-21. <https://doi.org/10.1080/00071669608417891>
 24. Amerah AM, Ravindran V, Lentle RG, Thomas DG. Influence of particle size and xylanase supplementation on the performance, energy utilisation, digestive tract parameters and digesta viscosity of broiler starters. *Br Poult Sci.* 2008;49:455-62. <https://doi.org/10.1080/00071660802251749>
 25. Patel S, Goyal A. Functional oligosaccharides: production, properties and applications. *World J Microbiol Biotechnol.* 2011;27:1119-28. <https://doi.org/10.1007/s11274-010-0558-5>
 26. Shalash SMM, Abou El-Wafa S, Sayed MAM, El-Gabry HE, Ramadan NA, Mohamed MS. Nutritive value of distillers dried grains with soluble and broiler performance at starter period. *Int J Poult Sci.* 2009;8:783-7. <https://doi.org/10.3923/ijps.2009.783.787>
 27. Slominski BA. Recent advances in research on enzymes for poultry diets. *Poult Sci.* 2011;90:2013-23. <https://doi.org/10.3382/ps.2011-01372>
 28. Mohammed AA, Habib AB, Eltrefi AM, Shulukh ESA, Abubaker AA. Effect of different levels of multi-enzymes (Natuzyne Plus®) on growth performance, carcass traits and meat quality of broiler chicken. *Asian J Anim Vet Adv.* 2018;13:61-6. <https://doi.org/10.3923/ajava.2018.61.66>
 29. Choct M. Feed non-starch polysaccharides: chemical structures and nutritional significance. *Feed Milling Int.* 1997;191:13-26.
 30. Cowieson AJ, Acamovic T, Bedford MR. Using the precision-feeding bioassay to determine the efficacy of exogenous enzymes—a new perspective. *Anim Feed Sci Technol.* 2006;129:149-58. <https://doi.org/10.1016/j.anifeedsci.2005.12.008>
 31. Williams MP, Klein JT, Wyatt CL, York TW, Lee JT. Evaluation of xylanase in low-energy broiler diets. *J Appl Poult Res.* 2014;23:188-95. <https://doi.org/10.3382/japr.2013-00856>
 32. González-Ortiz G, dos Santos TT, Bedford MR. Evaluation of xylanase and a fermentable xylo-oligosaccharide on performance and ileal digestibility of broiler chickens fed energy and amino acid deficient diets. *Anim Nutr.* 2021;7:488-95. <https://doi.org/10.1016/j.aninu.2020.07.008>
 33. Selle PH, Huang KH, Muir WI. Effects of nutrient specifications and xylanase plus phytase supplementation of wheat-based diets on growth performance and carcass traits of broiler chicks. *Asian-Australas J Anim Sci.* 2003;16:1501-9. <https://doi.org/10.5713/ajas.2003.1501>

34. Hu H, Dai S, Wen A, Bai X. Efficient expression of xylanase by codon optimization and its effects on the growth performance and carcass characteristics of broiler. *Animals*. 2019;9:65. <https://doi.org/10.3390/ani9020065>
35. Wu YB, Ravindran V. Influence of whole wheat inclusion and xylanase supplementation on the performance, digestive tract measurements and carcass characteristics of broiler chickens. *Anim Feed Sci Technol*. 2004;116:129-39. <https://doi.org/10.1016/j.anifeedsci.2004.02.011>
36. Masey O'Neill HVM, Mathis G, Lumpkins BS, Bedford MR. The effect of reduced calorie diets, with and without fat, and the use of xylanase on performance characteristics of broilers between 0 and 42 days. *Poult Sci*. 2012;91:1356-60. <https://doi.org/10.3382/ps.2011-01867>
37. Collins T, Gerday C, Feller G. Xylanases, xylanase families and extremophilic xylanases. *FEMS Microbiol Rev*. 2005;29:3-23. <https://doi.org/10.1016/j.femsre.2004.06.005>