

Effects of different processed forms of *Panax ginseng* on sperm motility and reproductive parameters in male dogs

Taeyoung Kil¹, Minkyu Kim^{2*}

¹Department of Social Welfare, Joongbu University, Geumsan 32713, Korea

²Division of Animal and Dairy Science, College of Agriculture and Life Science, Chungnam National University, Daejeon 34134, Korea



Received: Feb 27, 2025

Revised: Mar 21, 2025

Accepted: Mar 23, 2025

*Corresponding author

Minkyu Kim

Division of Animal and Dairy Science,
College of Agriculture and Life Science,
Chungnam National University,
Daejeon 34134, Korea

Tel: +82-42-821-5773

E-mail: kminkyu@cnu.ac.kr

Copyright © 2025 Korean Society of
Animal Science and Technology.

This is an Open Access article
distributed under the terms of the
Creative Commons Attribution
Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted
non-commercial use, distribution, and
reproduction in any medium, provided
the original work is properly cited.

ORCID

Taeyoung Kil

<https://orcid.org/0000-0003-4143-449X>

Minkyu Kim

<https://orcid.org/0000-0001-7099-9735>

Competing interests

No potential conflict of interest relevant
to this article was reported.

Funding sources

This work was supported by Joongbu
University Research Grant.

Acknowledgements

Not applicable.

Abstract

Male infertility in dogs is a significant concern in veterinary reproductive medicine, with sperm quality being a key determinant of reproductive success. Traditional herbal medicine, particularly *Panax ginseng*, is widely recognized for its potential to enhance male reproductive function. However, its effects on canine reproduction remain unexplored. This study investigated the impact of different processed forms of *Panax ginseng*—white ginseng (WG), red ginseng (RG), and black ginseng (BG)—on sperm motility, testosterone levels, and biochemical parameters in dogs. Beagle dogs were administered WG, RG, or BG daily for 60 days in a crossover design. Serum testosterone levels and biochemical markers were measured at predefined intervals, while sperm motility and velocity parameters were assessed using computer-assisted sperm analysis (CASA). The results demonstrated that BG supplementation significantly improved sperm motility and velocity parameters compared to WG and RG, with no adverse effects on biochemical markers. However, testosterone levels remained unchanged across groups. These findings suggest that BG may enhance canine sperm quality through mechanisms independent of testosterone regulation. Further research is needed to elucidate the underlying molecular pathways and optimize dosing strategies for clinical applications.

Keywords: Male infertility, *Panax ginseng*, Sperm motility, Testosterone, Canine reproduction, Computer-assisted sperm analysis

INTRODUCTION

Infertility is characterized by the inability of males or females to reproduce successfully and affects both humans and animals, including canines. Studies estimate that male reproductive disorders account for 40–50% of infertility cases in dogs [1]. Despite significant research efforts to address this issue, treatment success remains limited, with only 10% of infertility cases effectively managed [2].

Among various alternative treatments, herbal medicine—particularly *Panax ginseng*—has garnered

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Kil T.
Data curation: Kim M.
Formal analysis: Kil T.
Methodology: Kim M.
Software: Kil T.
Validation: Kim M.
Investigation: Kil T.
Writing - original draft: Kil T.
Writing - review & editing: Kil T, Kim M.

Ethics approval and consent to participate

The experimental procedures and methods used in this study were approved by the Animal Welfare and Ethics Office (CNU-00618) of Chungnam National University and performed according to the Guide for the Care and Use of Laboratory Animals published by the IACUC of Chungnam National University and ARRIVE guidelines.

significant attention for its potential role in improving reproductive health [3]. Previous studies have demonstrated its beneficial effects on male reproductive function [4]. The first report on ginseng's influence on spermatogenesis in rats highlighted its ability to enhance sperm production [5]. In human clinical trials, ginseng supplementation significantly improved sperm concentration and motility in patients with oligoasthenospermia and age-related infertility [6]. Beyond its reproductive benefits, ginseng exerts diverse pharmacological effects, supporting the immune [7], cardiovascular [8], and nervous systems [9], while also enhancing physical stamina [10]. Of particular interest, ginsenosides—the primary bioactive compounds in ginseng—share structural similarities with steroid hormones, suggesting their potential role in spermatogenesis and sexual function [11,12].

Panax ginseng is a perennial plant of the *Panax* genus and has been a fundamental component of traditional medicine in Korea and China [13]. The processing methods of ginseng significantly alter its chemical composition and biological activity, resulting in three main types: white ginseng (WG), red ginseng (RG), and black ginseng (BG). WG is produced by drying fresh roots under sunlight or hot air, whereas RG undergoes steaming before drying, giving it a reddish hue [14]. BG is subjected to multiple cycles of steaming and drying (at least three times), leading to a darker color and a higher concentration of bioactive compounds [15].

Studies have shown that both RG and BG contain a diverse range of ginsenosides with antioxidant and anti-inflammatory properties, including Rb1, Rb2, Rc, Rd, Re, Rf, Rg1, Rg2, 20(S)-Rg3, 20(R)-Rg3, Rg5, Rg6, Rh1, Rh4, Rk1, Rk3, F1, and F4. Repeated thermal processing significantly increases the levels of specific ginsenosides in BG compared to RG, particularly Rg2, Rg3, Rg5, Rg6, Rh1, Rh4, Rk1, Rk3, F1, and F4 [16]. Furthermore, research suggests that this thermal processing enhances the stability and bioavailability of key ginsenosides, including Rg3, Rg5, Rh1, and Rh2 [17,18].

However, the effects of *Panax ginseng* on canine reproductive physiology remain largely unexplored. Given these knowledge gaps, this study aims to evaluate the impact of ginseng supplementation on sperm motility and reproductive function in dogs. Three different types of processed ginseng (WG, RG, and BG) will be administered, and their effects will be assessed through blood analysis and computer-assisted sperm analysis (CASA). Specifically, this study will examine changes in testosterone levels, biochemical parameters, and sperm motility in response to ginseng supplementation. The findings will provide scientific evidence supporting the potential application of ginseng in canine reproductive health and offer valuable insights into natural supplementation strategies for improving male reproductive function in dogs.

MATERIALS AND METHODS

Unless otherwise indicated, all chemicals and reagents were purchased from Sigma-Aldrich Chemical Company (St. Louis, MO, USA).

Preparation of *Panax ginseng*

The processed *Panax ginseng*, including WG, RG, and BG, was obtained from Agricultural Corporation, GEUMSAN Black Ginseng Company (Korea), as shown in Fig. 1. The ginseng samples were ground using a blender and encapsulated into hard capsules, each containing 500 mg of ginseng powder [15,19].

Care and use of dogs

Male Beagle dogs weighing 7 to 12 kg and aged 2 to 3 years were used in this study. The animals were fed a commercial diet once daily and provided with *ad libitum* access to water. The dogs were



Fig. 1. Three different processed types of *Panax ginseng* products. (A) White ginseng: dried by sunlight or hot air, resulting in a white color. (B) Red ginseng: subjected to steaming and drying, giving it a red color. (C) Black ginseng: undergoes the same process as red ginseng but repeated at least three times, leading to a black color.

housed individually in an indoor animal facility [20]. All animal procedures were conducted in accordance with the “Guide for the Care and Use of Laboratory Animals” approved by Chungnam National University (Approval No.: CNU-00618).

Ginseng supplementation protocol

Three dogs (A, B, and C) were used in this study. Each dog received a single oral dose of one encapsulated 500 mg ginseng powder capsule per day for 60 days, followed by a 30-day washout period. After the washout period, the dogs were rotated into a different treatment group, ensuring that each dog underwent all three ginseng treatments in a crossover design [21], as illustrated in Fig. 2.




Fig. 2. The experimental design follows a switch-back group test. After 60 days of administration, dogs undergo a 30-day resting period before rotating into the next treatment group. RG, red ginseng; BG, black ginseng; WG, white ginseng.

Measurement of testosterone levels

To determine serum testosterone levels, blood samples were collected from the cephalic vein. The samples were centrifuged at 1,600 g for 10 minutes at room temperature. The radioimmunoassay for testosterone quantification was performed by Neodin (Seoul, Korea) [22]. The assay was conducted at 30-day intervals throughout the experimental period.

Biochemical analysis of serum

To assess the general health condition of the experimental dogs, serum biochemical analysis was conducted using the SPOTCHEM EZ SP-4430 (Arkray, Kyoto, Japan) [23]. Blood samples were collected from the cephalic vein and centrifuged at 1,600 g for 10 minutes at room temperature. According to the instrument manual, biochemical analysis was conducted at 30-day intervals throughout the experimental period, following the parameters listed in Table 1.

Computer-assisted sperm analysis

Semen quality was evaluated using CASA at 20-day intervals throughout the experimental period [24,25]. Semen samples were collected via digital manipulation, with only the second fraction (sperm-rich fraction) used for analysis, while the first and third fractions were discarded.

Sperm motility and kinematic parameters were assessed after appropriate dilution using the SAIS Plus system (Medical Supply, Wonju, Korea). A 5 μ L aliquot of diluted semen was placed onto a pre-warmed Makler chamber (New York Microscope Company, Hicksville, NY, USA) at 37°C. Sperm motility characteristics were examined under a 100 \times objective of an Eclipse E600 microscope (Nikon, Tokyo, Japan) equipped with a CCD camera (HVR-2000C, HVS, Seongnam, Korea). The following sperm motility parameters were measured using CASA: total motility (MOT, %), curvilinear velocity (VCL, μ m/sec), straight-line velocity (VSL, μ m/sec), average path velocity (VAP, μ m/sec), linearity (LIN, %) = (VSL/VCL) \times 100, straightness (STR, %) = (VCL/VAP) \times 100, and wobble (WOB, %) = (VAP/VCL) \times 100. Each measurement was performed in triplicate.

Statistical analysis

The results were expressed as means \pm SE. The Shapiro-Wilk test was used to assess normality, while Levene's test evaluated the homogeneity of variances. Data analysis was performed using R software (version 4.3.2, USA). One-way analysis of variance (ANOVA) was conducted, followed by Tukey's post hoc test for multiple comparisons. Pearson correlation analysis was used to assess the relationship between serum P4 levels. Statistical significance was set at $p < 0.05$.

Table 1. The index list of biochemical analysis of canine serum

Abbreviation	Full name	Reference range
T-Pro	Total protein	5.4–7.5 g/dL
Alb	Albumin	2.3–3.1 g/dL
ALP	Alkaline phosphatase	1–114 U/L
AST/GOT	Aspartate aminotransferase	13–15 U/L
GPT	Glutamic-pyruvic transaminase	10–109 U/L
Glu	Glucose	76–119 mg/dL
BUN	Blood urea nitrogen	8–28 mg/dL
Cre	Creatinine	0.5–1.7 mg/dL

RESULTS

Testosterone (T4) levels

To assess variations in T4 levels among the experimental groups (WG, RG, and BG), hormone analysis was conducted at three time points: Day 0, Day 30, and Day 60 of the experimental period. No significant differences in T4 levels were observed among the treatment groups over the 60-day period (Fig. 3). However, a trend of increasing T4 levels was noted in the ginseng-administered groups compared to the control group throughout the study.

Biochemical analysis

Serum samples were collected on Days 0, 30, and 60 of the experimental period and analyzed using a serum analyzer to assess changes in biochemical markers following ginseng supplementation. No significant changes in biochemical markers were observed, and all values remained within the normal physiological range (Figs. 4, 5, and 6). Furthermore, no adverse health effects were detected




Fig. 3. Relative levels of testosterone in dogs during experimental periods. Control, non-treatment group; WG, white ginseng group; RG, red ginseng group; BG, black ginseng group.




Fig. 4. The variation of biochemical markers in WG group during the experimental periods. T-Pro, total protein (g/dL); Alb, albumin (g/dL); ALP, alkaline phosphatase (U/L); AST/GOT, aspartate aminotransferase (U/L); GPT, glutamic-pyruvic transaminase (U/L); Glu, glucose (mg/dL); BUN, blood urea nitrogen (mg/dL); Cre, creatinine (mg/dL); WG, white ginseng.




Fig. 5. The variation of biochemical markers in RG group during the experimental periods. T-Pro, total protein (g/dL); Alb, albumin (g/dL); ALP, alkaline phosphatase (U/L); AST/GOT, aspartate aminotransferase (U/L); GPT, glutamic-pyruvic transaminase (U/L); Glu, glucose (mg/dL); BUN, blood urea nitrogen (mg/dL); Cre, creatinine (mg/dL); RG, red ginseng.




Fig. 6. The variation of biochemical markers in BG group during the experimental periods. T-Pro, total protein (g/dL); Alb, albumin (g/dL); ALP, alkaline phosphatase (U/L); AST/GOT, aspartate aminotransferase (U/L); GPT, glutamic-pyruvic transaminase (U/L); Glu, glucose (mg/dL); BUN, blood urea nitrogen (mg/dL); Cre, creatinine (mg/dL); BG, black ginseng.

in any of the experimental groups (WG, RG, and BG).

Effects of *Panax ginseng* on canine spermatozoa motility by computer-assisted sperm analysis

A two-way ANOVA was conducted to assess the effects of different treatments and time points on sperm motility (Fig. 7). On Day 40, motility in the BG group was significantly higher than in the WG group ($p < 0.05$). By Day 60, the BG group exhibited significantly higher motility than both the WG ($p < 0.001$) and RG groups ($p < 0.05$). Within-group comparisons revealed that motility in the WG group increased significantly on Day 60 compared to Day 0 and Day 20 ($p < 0.05$). Similarly, motility in the RG group showed a significant increase on Day 60 ($p < 0.01$). In the BG group, motility was significantly higher on Day 40 ($p < 0.01$) and peaked on Day 60 ($p < 0.0001$), with a further increase from Day 40 to Day 60 ($p < 0.05$).




Fig. 7. The variation of motility among treatment groups by CASA. ^{A-C} Different uppercase letters indicate significant differences within the same treatment across different time points ($p < 0.05$). ^{a,b} While different lowercase letters indicate significant differences among treatments at the same time point ($p < 0.05$). MOT, total motility; WG, white ginseng group; RG, red ginseng group; BG, black ginseng group; CASA, computer-assisted sperm analysis.

Effects of *Panax ginseng* on canine spermatozoa velocity parameters by computer-assisted sperm analysis

Sperm velocity parameters were measured using CASA. A two-way ANOVA was conducted to assess the effects of different treatments and time points on sperm VCL (Fig. 8). On Day 60, only the BG group exhibited significantly higher VCL than the RG group ($p < 0.05$). Within-group comparisons over time showed that WG VCL increased significantly on Day 40 ($p < 0.05$) and Day 60 ($p < 0.01$) compared to Day 0. Similarly, RG VCL was significantly higher on Day 40 and Day 60 compared to Day 0 ($p < 0.05$). In the BG group, VCL significantly increased on Day 40 ($p < 0.05$) and Day 60 ($p < 0.01$) relative to Day 0, with Day 60 also significantly exceeding Day 20 ($p < 0.05$).

A two-way ANOVA was also conducted to assess the effects of different treatments and time




Fig. 8. The variation of VCL among treatment groups by CASA. ^{A-C} Different uppercase letters indicate significant differences within the same treatment across different time points ($p < 0.05$). ^{a,b} While different lowercase letters indicate significant differences among treatments at the same time point ($p < 0.05$). VCL, curvilinear velocity; WG, white ginseng group; RG, red ginseng group; BG, black ginseng group; CASA, computer-assisted sperm analysis.




Fig. 9. The variation of VSL among treatment groups by CASA. ^{A-D} Different uppercase letters indicate significant differences within the same treatment across different time points ($p < 0.05$). ^{a-c} While different lowercase letters indicate significant differences among treatments at the same time point ($p < 0.05$). VSL, straight-line velocity; WG, white ginseng group; RG, red ginseng group; BG, black ginseng group; CASA, computer-assisted sperm analysis.

points on sperm VSL (Fig. 9). The BG group exhibited significantly higher VSL than the WG group on Day 20 ($p < 0.01$). On Day 40, VSL in the BG group was significantly higher than in both WG ($p < 0.05$) and RG ($p < 0.05$). By Day 60, the RG group showed significantly higher VSL than the WG group ($p < 0.01$), while the BG group exhibited the highest VSL, significantly exceeding both WG ($p < 0.0001$) and RG ($p < 0.05$). Within-group comparisons showed that WG VSL increased significantly on Day 40 ($p < 0.001$) and Day 60 ($p < 0.0001$) compared to Day 0, with Day 60 also exceeding Day 20 ($p < 0.0001$). RG VSL was significantly higher on Day 40 ($p < 0.05$) and Day 60 ($p < 0.0001$) compared to Day 0, with Day 60 also exceeding Day 20 ($p < 0.0001$) and Day 40 ($p < 0.0001$). The BG group exhibited the most substantial increase, with significantly higher VSL on Day 20 ($p < 0.05$), Day 40 ($p < 0.0001$), and Day 60 ($p < 0.0001$) compared to Day 0. Additionally, BG VSL on Day 60 was significantly higher than on Day 20 ($p < 0.0001$) and Day 40 ($p < 0.0001$).

Effects of *Panax ginseng* on canine spermatozoa average path velocity by computer-assisted sperm analysis

A two-way ANOVA was conducted to assess the effects of different treatments and time points on sperm VAP (Fig. 10). On Day 40, the BG group exhibited significantly higher VAP than the WG group ($p < 0.05$). By Day 60, VAP in the BG group was significantly higher than in the WG group ($p < 0.001$). Within-group comparisons showed a significant increase in BG VAP on Day 60 compared to Day 0 ($p < 0.01$) and Day 20 ($p < 0.05$). No other within-group comparisons reached statistical significance.

Effects of *Panax ginseng* on canine spermatozoa trajectory by computer-assisted sperm analysis

Sperm trajectory was assessed by measuring LIN, STR, and WOB. A two-way ANOVA was conducted to evaluate the effects of different treatments and time points on sperm LIN (Fig. 11). On Day 20, both the RG and BG groups displayed significantly higher LIN compared to the WG group ($p < 0.05$). By Day 60, LIN was significantly higher in the RG group than in the WG group ($p < 0.01$), while the BG group exhibited the highest LIN, significantly exceeding WG ($p < 0.001$).




Fig. 10. The variation of VAP among treatment groups by CASA. ^{A,B} Different uppercase letters indicate significant differences within the same treatment across different time points ($p < 0.05$). ^{a,b} While different lowercase letters indicate significant differences among treatments at the same time point ($p < 0.05$). WG, white ginseng group; RG, red ginseng group; BG, black ginseng group; CASA, computer-assisted sperm analysis.




Fig. 11. The variation of LIN among treatment groups by CASA. ^{A-C} Different uppercase letters indicate significant differences within the same treatment across different time points ($p < 0.05$). ^{a,b} While different lowercase letters indicate significant differences among treatments at the same time point ($p < 0.05$). LIN, linearity; WG, white ginseng group; RG, red ginseng group; BG, black ginseng group; CASA, computer-assisted sperm analysis.

Within-group comparisons showed that LIN in the WG group increased significantly on Day 40 ($p < 0.05$) and Day 60 ($p < 0.001$), with a further increase on Day 60 compared to Day 20 ($p < 0.0001$). In the RG group, LIN was significantly higher on Day 60 compared to Day 0 ($p < 0.0001$), Day 20 ($p < 0.0001$), and Day 40 ($p < 0.001$). Similarly, in the BG group, LIN was significantly higher on Day 40 ($p < 0.001$) and Day 60 ($p < 0.0001$) compared to Day 0, with further increases on Day 60 compared to Day 20 ($p < 0.0001$) and Day 40 ($p < 0.001$).

A two-way ANOVA revealed no significant differences in STR among groups at any time point ($p > 0.05$; Fig. 12). However, within-group comparisons showed significant increases over time. In the WG group, STR was significantly higher on Day 40 ($p < 0.01$) and Day 60 ($p < 0.0001$) compared to Day 0, with an additional increase on Day 40 relative to Day 20 ($p < 0.01$). Similarly, in the RG group, STR was significantly higher on Day 40 ($p < 0.05$) and Day 60 ($p < 0.0001$) compared to




Fig. 12. The variation of STR among treatment groups by CASA. ^{A-C} Different uppercase letters indicate significant differences within the same treatment across different time points ($p < 0.05$). ^a While different lowercase letters indicate significant differences among treatments at the same time point ($p < 0.05$). STR, straightness; WG, white ginseng group; RG, red ginseng group; BG, black ginseng group; CASA, computer-assisted sperm analysis.

Day 0, with further increases on Day 60 relative to Day 20 ($p < 0.0001$) and Day 40 ($p < 0.001$). The BG group exhibited the most pronounced increases, with significantly higher STR on Day 40 ($p < 0.001$) and Day 60 ($p < 0.0001$) compared to Day 0, as well as significant increases on Day 60 relative to Day 20 ($p < 0.0001$) and Day 40 ($p < 0.01$).

A two-way ANOVA was conducted to assess the effects of different treatments and time points on sperm WOB (Fig. 13). The RG group exhibited significantly higher WOB than the WG group ($p < 0.05$), while the BG group had significantly higher WOB than the WG group ($p < 0.01$). No significant differences were observed between the BG and RG groups at any time point. Within-group comparisons showed no significant changes in WOB over time for either WG or RG.




Fig. 13. The variation of WOB among treatment groups by CASA. ^A Different uppercase letters indicate significant differences within the same treatment across different time points ($p < 0.05$). ^{a,b} While different lowercase letters indicate significant differences among treatments at the same time point ($p < 0.05$). WOB, wobble; WG, white ginseng group; RG, red ginseng group; BG, black ginseng group; CASA, computer-assisted sperm analysis.

DISCUSSION

The present study investigated the effects of three types of processed ginseng (WG, RG, and BG) on male canine reproductive physiology. This was evaluated through testosterone (T4) measurement, biochemical serum analysis, and sperm quality assessment.

In 2007, a study reported that a dosage of 1 g of ginsenoside was used in human trials, with no adverse effects [19]. However, no prior research has examined the effects of ginseng supplementation in dogs. Therefore, in this study, the dosage was set at 500 mg per day, half of the dosage used in humans.

Testosterone, a primary male steroid hormone, is produced by Leydig cells in the testes and plays a crucial role in libido regulation [26]. Previous studies have demonstrated that feeding rats a diet containing 5% *Panax ginseng* for 60 days significantly increased serum testosterone levels [27]. Additionally, supplementation with ginsenoside Rg1 (10 mg/kg) significantly increased serum testosterone levels and enhanced sexual behavior in animal models [28]. However, in this study, no significant differences in testosterone levels were observed. Despite this, no adverse effects were detected in any of the experimental groups (WG, RG, or BG). Biochemical analysis of serum samples indicated that all measured markers remained within normal physiological ranges throughout the study. No significant deviations were observed, suggesting that oral administration of WG, RG, and BG did not negatively affect liver, kidney, or general health functions.

Semen quality is influenced by various factors, including health, nutrition, and oxidative stress [29,30]. *Panax ginseng* has long been studied for its reproductive benefits, particularly its ability to enhance sperm motility [4]. BG, which undergoes repeated steaming and drying, contains higher levels of secondary ginsenosides, such as Rg3, Rg5, and Rk1, compared to WG and RG, contributing to its greater bioactivity [16]. Recent studies suggest that ginsenosides improve sperm function by enhancing mitochondrial activity and reducing oxidative stress [31]. In particular, BG may protect sperm from oxidative damage by activating the Nrf2/ARE pathway, which upregulates antioxidant enzymes such as SOD and GPx [32]. Additionally, ginsenosides regulate lipid metabolism, a key factor in maintaining sperm membrane integrity and motility [33]. El-Shimi et al. reported that ginseng mitigates reproductive toxicity caused by environmental pollutants like bisphenol A, preventing mitochondrial dysfunction and sperm DNA damage [11]. Moreover, Zhang et al. demonstrated that ginseng-derived soluble dietary fiber enhances spermatogenesis via the MAPK signaling pathway, indicating a role in cellular energy regulation [34]. These mechanisms align with our findings, where CASA analysis showed that BG significantly improved sperm motility and trajectory parameters.

Despite these promising findings, several limitations should be considered. Although ginseng supplementation improved sperm motility, it did not lead to significant changes in testosterone levels. This suggests that its mechanism of action may not be directly testosterone-mediated, warranting further investigation. Additionally, while no adverse effects on liver, kidney, or overall health functions were observed, the long-term effects remain unclear. Future studies should evaluate prolonged supplementation to determine potential cumulative effects.

The role of specific ginsenosides requires further investigation. While Rg3 has been shown to exhibit bioactive properties [35], this study did not isolate individual ginsenosides. Future research should focus on identifying key active compounds and elucidating their mechanisms of action. Additionally, varying dosages and supplementation durations should be tested to optimize reproductive benefits.

Given the structural similarities between ginsenosides and steroid hormones [36], potential endocrine interactions should also be explored. While this study provides valuable insights, further

research is needed to validate these findings and refine the applications of ginseng in reproductive medicine.

REFERENCES

1. Lopate C. Advances in canine semen evaluation techniques. *Clin Theriogenol.* 2009;1:169-94.
2. Johnston SD, Kustritz MV, Olson PS. Canine and feline theriogenology. Philadelphia, PA: Saunders; 2001.
3. Leung KW, Wong AS. Ginseng and male reproductive function. *Spermatogenesis.* 2013;3:e26391.
4. Wang H, Zhang J, Ma D, Zhao Z, Yan B, Wang F. The role of red ginseng in men's reproductive health: a literature review. *Basic Clin Androl.* 2023;33:27.
5. Yamamoto M, Takeuchi N, Kumagai A, Yamamura Y. Stimulatory effect of *Panax ginseng* principles on DNA, RNA, protein and lipid synthesis in rat bone marrow. *Arzneimittelforschung.* 1977;27:1169-73.
6. Salvati G, Genovesi G, Marcellini L, Paolini P, De Nuccio I, Pepe M, et al. Effects of *Panax ginseng* C.A. Meyer saponins on male fertility. *Panminerva Med.* 1996;38:249-54.
7. Riaz M, Rahman NU, Zia-Ul-Haq M, Jaffar HZ, Manea R. Ginseng: a dietary supplement as immune-modulator in various diseases. *Trends Food Sci Technol.* 2019;83:12-30.
8. Hyun SH, Bhilare KD, In G, Park CK, Kim JH. Effects of *Panax ginseng* and ginsenosides on oxidative stress and cardiovascular diseases: pharmacological and therapeutic roles. *J Ginseng Res.* 2022;46:33-8.
9. Kim KH, Lee D, Lee HL, Kim CE, Jung K, Kang KS. Beneficial effects of *Panax ginseng* for the treatment and prevention of neurodegenerative diseases: past findings and future directions. *J Ginseng Res.* 2018;42:239-47.
10. Choi JY, Woo TS, Yoon SY, dela Peña IC, Choi YJ, Ahn HS, et al. Red ginseng supplementation more effectively alleviates psychological than physical fatigue. *J Ginseng Res.* 2011;35:331-8.
11. El-Shimi BI, Mohareb RM, Ahmed HH, Abohashem RS, Mahmoud KF, Hanna DH. Mechanistic insights into bisphenol a-mediated male infertility: potential role of *Panax ginseng* extract. *Chem Biodivers.* 2024;21:e202400480.
12. Ratan ZA, Haidere MF, Hong YH, Park SH, Lee JO, Lee J, et al. Pharmacological potential of ginseng and its major component ginsenosides. *J Ginseng Res.* 2021;45:199-210.
13. Zhang H, Abid S, Ahn JC, Mathiyalagan R, Kim YJ, Yang DC, et al. Characteristics of *Panax ginseng* cultivars in Korea and China. *Molecules.* 2020;25:2635.
14. Nam KY. The comparative understanding between red ginseng and white ginsengs, processed ginsengs (*Panax ginseng* CA Meyer). *J Ginseng Res.* 2005;29:1-18.
15. Jang AY, Sueng YC, Ji JG. The comparative study on physiological activity of white ginseng, red ginseng and black ginseng extract. *J Digit Converg.* 2016;14:459-71.
16. Jo HK, Sung MC, Ko SK. The comparison of ginseng prosapogenin composition and contents in red and black ginseng. *Korean J Pharmacogn.* 2011;42:361-5.
17. Metwaly AM, Lianlian Z, Luqi H, Deqiang D. Black ginseng and its saponins: preparation, phytochemistry and pharmacological effects. *Molecules.* 2019;24:1856.
18. Lee JH, Shen GN, Kim EK, Shin HJ, Myung CS, Oh HJ, et al. Preparation of black ginseng and its antitumor activity. *Korean Orient J Physiol Pathol.* 2006;20:951-6.
19. De Andrade E, De Mesquita AA, de Almeida Claro J, De Andrade PM, Ortiz V, Paranhos M, et al. Study of the efficacy of Korean red ginseng in the treatment of erectile dysfunction. *Asian*

- J Androl. 2007;9:241-4.
20. Kim DE, Lee JH, Ji KB, Lee EJ, Li C, Oh HJ, et al. Prime editor-mediated correction of a pathogenic mutation in purebred dogs. *Sci Rep.* 2022;12:12905.
 21. Jones B, Kenward MG. Design and analysis of cross-over trials. Chapman and Hall/CRC; 2003.
 22. Choi EG, Yin XJ, Lee HS, Kim LH, Shin HD, Kim NH, et al. Reproductive fertility of cloned male cats derived from adult somatic cell nuclear transfer. *Cloning Stem Cells.* 2007;9:281-90.
 23. Gülersoy E, Ekici YE. Assessment of hematological and serum biochemistry parameters in dogs with acute diarrhea due to different etiologies. *Maced Vet Rev.* 2022;45:149-56.
 24. Kutzler MA. Semen collection in the dog. *Theriogenology.* 2005;64:747-54.
 25. Li C, Oh HJ, Liu H, Kim MK. Schisandrin B protects boar spermatozoa against oxidative damage and increases their fertilization ability during in vitro storage. *Theriogenology.* 2023;198:194-201.
 26. Nassar GN, Leslie SW. Physiology, testosterone [Internet]. StatPearls. 2025 [cited 2025 Jan 4]. <https://www.ncbi.nlm.nih.gov/books/NBK526128/>
 27. Seftel AD, Mack RJ, Secrest AR, Smith TM. Restorative increases in serum testosterone levels are significantly correlated to improvements in sexual functioning. *J Androl.* 2004;25:963-72.
 28. Wang X, Chu S, Qian T, Chen J, Zhang J. Ginsenoside Rg1 improves male copulatory behavior via nitric oxide/cyclic guanosine monophosphate pathway. *J Sex Med.* 2010;7:743-50.
 29. Salas-Huetos A, James ER, Aston KI, Jenkins TG, Carrell DT. Diet and sperm quality: nutrients, foods and dietary patterns. *Reprod Biol.* 2019;19:219-24.
 30. Jurewicz J, Radwan M, Sobala W, Ligocka D, Radwan P, Bochenek M, et al. Lifestyle and semen quality: role of modifiable risk factors. *Syst Biol Reprod Med.* 2014;60:43-51.
 31. Koppula S, Kopalli SR, Kang HH, Kim SK. Benefits of Panax ginseng on male reproductive systems: a comprehensive review. *Food Suppl Biomater Health.* 2023;3:e32.
 32. Barakat N, Alkhen MA, Khater Y, Khirallah SM. Effect of melatonin and ginseng on rat testis and sperm quality against cadmium toxicity via inhibiting oxidative stress and autophagy pathways. *J Trace Elem Med Biol.* 2025;88:127614.
 33. Khan MT, Khan MI-u-R, Yasmin T, Khan GS, Riaz A. Effect of ginseng on blood lipid profile, testosterone level and epididymal sperm quality of aged BALB/C mice. *Pakistan J Zool.* 2025;57:501-63. <https://doi.org/10.17582/journal.pjz/20230323090320>
 34. Zhang Y, Yu Y, Bai C, Li Z, Huo X, Li W, et al. Ginseng soluble dietary fiber enhances spermatogenic potential in obese mice via the MAPK signaling pathway. *J Food Biochem.* 2024;2024:6235198.
 35. Yoon SJ, Park JY, Choi S, Lee JB, Jung H, Kim TD, et al. Ginsenoside Rg3 regulates S-nitrosylation of the NLRP3 inflammasome via suppression of iNOS. *Biochem Biophys Res Commun.* 2015;463:1184-9.
 36. Shi ZY, Zeng JZ, Wong AST. Chemical structures and pharmacological profiles of ginseng saponins. *Molecules.* 2019;24:2443.