

# Anti-Müllerian hormone and antral follicle count as predictors for optimal selection of Hanwoo donor cows in superstimulated oocyte collection

Joonho Moon<sup>1#</sup>, Jae Jung Ha<sup>2#</sup>, Woo-sung Kwon<sup>3</sup>, Daehyun Kim<sup>4\*</sup>,  
Junkoo Yi<sup>5,6\*</sup>

<sup>1</sup>GenNBio Co., Ltd., Pyeongtaek 17796, Korea


<sup>2</sup>Gyeongbuk Livestock Research Institute, Yeongju 36052, Korea

<sup>3</sup>Department of Animal Science and Biotechnology, Kyungpook National University, Sangju 37224, Korea

<sup>4</sup>Department of Animal Science, Chonnam National University, Gwangju 61186, Korea

<sup>5</sup>School of Animal Life Convergence Science, Hankyong National University, Anseong 17579, Korea

<sup>6</sup>Gyeonggi Regional Research Center, Hankyong National University, Anseong 17579, Korea



Received: Mar 5, 2024

Revised: May 1, 2024

Accepted: May 4, 2024

#These authors contributed equally to this work.

## \*Corresponding author

Daehyun Kim  
Department of Animal Science,  
Chonnam National University, Gwangju  
61186, Korea  
Tel: +82-62-530-2123  
E-mail: kimdhbio@jnu.ac.kr

Junkoo Yi  
School of Animal Life Convergence  
Science, Hankyong National University,  
Anseong 17579, Korea  
Tel: +82-31-670-5092  
E-mail: junkoo@hknu.ac.kr

Copyright © 2025 Korean Society of  
Animal Science and Technology.  
This is an Open Access article  
distributed under the terms of the  
Creative Commons Attribution  
Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted  
non-commercial use, distribution, and  
reproduction in any medium, provided  
the original work is properly cited.

## ORCID

Joonho Moon  
<https://orcid.org/0000-0001-6513-557X>

## Abstract

This study explored the correlation between anti-Müllerian hormone (AMH) levels and reproductive efficiency in native Korean cattle (Hanwoo) as potential oocyte donors in ovum pick-up (OPU) programs. In an effort to enhance the efficiency of oocyte collection, this research explored the correlation between AMH levels and various factors, including the quantity of follicles, retrieved oocytes, and the proportion of transferable embryos. A total of 42 Hanwoo cows were included in this study with AMH levels ranging from 276.5 to 2,212.5 pg/mL. These cows were categorized into three groups based on AMH concentration: high (H), medium (M), and low (L), along with the control group. To monitor the quantity of antral follicles in each group during OPU, Ultrasound scanner was used, and the retrieved oocytes were duly recorded. The implantable embryos produced from the retrieved oocytes were quantified. The results show a significant positive correlation between AMH levels and the numbers of antral follicles ( $R^2 = 0.5785$ ,  $p < 0.0001$ ), retrieved oocytes ( $R^2 = 0.6857$ ,  $p < 0.0001$ ) and transferable embryos ( $R^2 = 0.4049$ ,  $p < 0.0001$ ), indicating that higher AMH levels correspond to increased numbers of antral follicles and retrieved oocytes. However, the correlation between AMH levels and the proportion of transferable embryos was not statistically significant ( $R^2 = 0.1476$ ,  $p = 0.5225$ ). In conclusion, AMH levels were significantly correlated with the number of antral follicles and retrieved oocytes, demonstrating their potential as indicators for selecting oocyte donors for Hanwoo cattle. Although the relationship with the proportion of transferable embryos was not statistically significant, this study offers valuable insights for the improvement of the efficiency of oocyte donation plans in Hanwoo cattle by considering the AMH levels.

**Keywords:** Anti-Müllerian hormone, Antral follicles count, Ovum pick-up, Hanwoo donor, *In vitro* embryo production

Jae Jung Ha  
<https://orcid.org/0000-0001-6785-6346>  
 Woo-sung Kwon  
<https://orcid.org/0000-0002-0848-7189>  
 Daehyun Kim  
<https://orcid.org/0000-0002-4820-4438>  
 Junkoo Yi  
<https://orcid.org/0000-0003-2593-6529>

#### Competing interests

No potential conflict of interest relevant to this article was reported.

#### Funding sources

This work was supported by a research grant from Hankyong National University in the year of 2023.

#### Acknowledgements

Not applicable.

#### Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

#### Authors' contributions

Conceptualization: Moon J, Ha JJ.  
 Data curation: Moon J, Ha JJ.  
 Formal analysis: Kwon WS.  
 Methodology: Moon J, Yi J.  
 Software: Moon J, Kim DH.  
 Validation: Kim DH, Yi J.  
 Investigation: Ha JJ, Kim DH, Yi J.  
 Writing - original draft: Moon J, Ha JJ.  
 Writing - review & editing: Moon J, Ha JJ, Kwon WS, Kim DH, Yi J.

#### Ethics approval and consent to participate

This study was conducted in accordance with the guidelines of the Declaration of Helsinki and was approved by the Institutional Animal Care and Use Committee of the Gyeongsangbukdo Livestock Research Institute in Gyeongsangbuk-do (Approval No. GAEC/161/23, approved on 14 December 2022).

## INTRODUCTION

The production and transfer of embryos in the bovine species has been widely used as an effective method for genetic improvement in many livestock-producing countries. This trend is also increasing in Korea, where awareness of embryo transfer for breed improvement and genetic resource conservation is growing. Moreover, reproductive efficiency is strongly correlated with the economic viability of dairy and beef industries [1–7]. Consequently, assisted reproductive technologies such as *in vitro* embryo production (IVEP) using ovum pick-up (OPU), have been globally adopted to rapidly obtain genetically superior traits in cows [8]. The number of large-scale farms and intensive production systems for Hanwoo (native Korean) and Holstein cattle is increasing in Korea. To achieve successful breeding, the importance of embryo production as a valuable trait is increasing, not only in traditional breeding programs such as artificial insemination, but also in enhancing the pace of genetic improvement. In Korea, cows with excellent carcass characteristics or high genetic value are preferred for breeding, and multiparous cows are primarily used as valuable donors. Factors such as growth rate, pedigree, market situation, temperament, and meat quality are prioritized as criteria for selecting donor cows, often neglecting indicators related to the inherent oocyte or embryo production capacity [9–13]. Moreover, recent reports have indicated significant variability in the response to superstimulation treatments and the quantity of oocytes retrieved via OPU [14–16]. Recent studies have suggested that in cows, anti-Müllerian hormone (AMH) measurements and antral follicle count (AFC) through ovarian ultrasound scanning can serve as predictive variables for the quantity of oocytes collected, thereby assisting in forecasting the ovarian response to superstimulation treatment [17,18]. However, there is a lack of research exploring the correlation between AMH levels and AFC with respect to the quantity of oocytes retrieved and transferable embryos in *Bos taurus* breeds, such as Hanwoo. Also, the efficiency of donor selection using the superstimulation method, which has become a global trend, needs to be demonstrated. Consequently, the development and application of predictive methods to determine the inherent oocyte or embryo production capacity of donor cows are essential.

During male fetal differentiation, Sertoli cells in the testes secrete AMH, leading to regression of the Müllerian ducts. In the ovaries, granulosa cells from preantral or antral follicles also produce AMH [19–22]. The physiological functions of AMH are not completely understood; however, it is thought to regulate follicular recruitment and selection [23]. Serum AMH concentration is closely associated with the quantity of AFC and remains relatively stable throughout the estrous cycle [24]. Serum AMH levels have recently been proposed as a good indicator of ovarian reserve, showing a strong correlation with the quantity of oocytes retrieved by OPU [25]. However, recent studies have emphasized the need for reliable research on the use of AMH and AFC as selection indicators of the quantity of oocytes by superstimulation, rather than OPU in a random ovarian cycle state without superstimulation treatment in cows [26–29]. In Hanwoo cattle, recent research has shown a correlation between AMH concentrations and the quantity of embryos retrieved from donors with normal ovarian cyclicity [30]. This suggests the value of early evaluation of AMH concentrations when selecting potential Hanwoo embryo donors. Based on these findings, the measurement of AMH concentration and AFC in Hanwoo cows during superstimulation, along with follicle-stimulating hormone (FSH) treatment, could be a valuable predictive tool for assessing the reproductive potential of donor cows.

The quantity of embryos produced by the OPU-IVEP varies according to the quantity of oocytes retrieved [31]. Choosing donors specifically for the OPU-IVEP appears to be most effective strategy for enhancing the yield of superior oocytes and embryos [30]. Thus, our study aimed to improve genetics and embryo production efficiency through OPU sessions. Hanwoo donors were

selected using AMH concentration measurements as a basis for commercial farms. To correlate AMH as an indicator of ovarian function in superstimulated donors in this study, we measured AMH hormone concentrations in a total of 42 donors, and 12 donors with concentrations close to the mean were used as non-superstimulated controls. The 30 donors were used as the superstimulated group, and the group was divided into high, medium, and low groups according to concentration. Additionally, we evaluated the correlations between AFC measurements before OPU testing and the quantity of oocytes retrieved, embryos, and embryo-to-oocyte ratio after OPU testing among the groups.

## MATERIALS AND METHODS

### Animals and blood sampling


All procedures involving animals in this study were in accordance with relevant national laws and guidelines for animal care and use. Approval for the study was obtained from the Institutional Animal Care and Use Committee of the Gyeongsangbukdo Livestock Research Institute (Approval No. GAEC/161/23, approved on 14 December 2022). The experiment was carried out with Hanwoo donors ( $n = 42$ ). The Hanwoo donors had a normal cycle and were  $4.1 \pm 0.6$  (mean  $\pm$  SEM) years of age and were kept on a commercial farm in Gyeongsangbukdo, Korea, from March to September 2023. The mean body condition score was  $3.0 \pm 0.2$  (mean  $\pm$  SEM) on a scale of 1 to 5 (1 = very thin; 5 = very fatty, respectively) [32]. They had unrestricted access to water and mineralized salts. Venous blood samples were obtained from the donor's jugular vein immediately before determining their eligibility as oocyte donors. The blood was drawn into tubes and centrifuged at  $1,314 \times g$  for 10 min to separate the plasma. The recovered plasma was preserved at a temperature of  $-80^\circ\text{C}$  until subsequent AMH testing was conducted.

### Quantification of anti-Müllerian hormone plasma concentration and donor classification

A bovine AMH ELISA kit (Ansh Labs, Webster, MA, USA) was used to assess usual plasma AMH concentrations prior to oocytes retrieval, in accordance with a previous report [33]. Donors were categorized into three treatment groups and one control group based on their AMH concentrations. Group H included the top 30%, Group M included the next 30%, and Group L included donors with the lowest concentration at 40%. The control group comprised 12 donors whose AMH concentrations were closest to the mean value of the 42 donors. Nine donors from group H, 9 from group M, and 12 from group L were selected as on-farm donors for OPU.

### Superstimulation and ovum pick-up

OPU was performed on 42 selected cows six times every two weeks by two skilled technicians. Prior to OPU handling, each donor's follicular waves were synchronized with controlled intravaginal drug-release (1.38 g of Progesterone, CIDR DEVICES®, Zoetis, Rhodes, New South Wales, Australia) insertion and simultaneous administration of estradiol benzoate (2.0 mg/cow, Esron®, Samyang-Anipharma, Seoul, Korea) to induce follicular wave present in the ovary on day one. On Day 3, 5.0 mg of intramuscular prostaglandin F $_{2\alpha}$  (PGF $_{2\alpha}$ ) (Lutalyse®, Zoetis, Brussels, Belgium) was administered. Starting four days after insertion, FSH (Antrin R-10®, Kyoritsu Seiyaku, Tokyo, Japan) was given twice daily over a span of two days, with decreasing doses given at 12-hour intervals. The dosage regimens used were 3.6, 3.6, 2.4, and 2.4 AU on days four and five. On day seven, OPU was conducted 36 h after the final FSH injection, following removal of the progesterone (P $_4$ ) device (Fig 1). The follicle count was assessed by ultrasound immediately



**Fig. 1. OPU was conducted six times at 2-week intervals.** Hanwoo donors (superstimulated group = 30, control group = 10) were divided into two groups and synchronized. The superstimulated group received four intramuscular FSH injections of 12 AU, 12 h apart, on days four and five, whereas the control group received no further treatment. Dosages were 3.6, 3.6, 2.4, and 2.4 AU for FSH, 2.0 mg for EB, 5.0 mg for PG, and 1.38 g for the P<sub>4</sub> device. AFC, antral follicle count; EB, estradiol benzoate; PG, prostaglandin F<sub>2</sub>α; FSH, follicle-stimulating hormone; OPU, ovum pick-up; P<sub>4</sub> device, progesterone.

before oocyte retrieval, specifically by counting the quantity of antral follicles ranging in diameter from 1 to 15 mm. Ovarian ultrasound scanner (4Vet Slim, Draminski Tech, Olsztyn, Poland) were conducted with a 6.5 MHz OPU endovaginal probe (BLUE, Draminski Tech) on day seven, immediately before the OPU session. During the OPU, the aspiration medium (MK\_OPU®, MK biotech, Daejeon, Korea) used to retrieve the oocytes was anticoagulated with heparin to prevent blood clotting. A 19-G disposable hypodermic needle was used to perform the follicular puncture. A vacuum pump was used to maintain the vacuum for aspiration between 45 and 60 mmHg (BV-003, WTA, Cravinhos, Brazil) during the OPU procedure in both ovaries. Follicular contents were recovered using a 100 cm long tube with an internal diameter of 1.1 mm (WTA). The recovered cumulus–oocyte complexes (COCs) were washed once in wash medium (MK\_WM®, MK biotech, Daejeon, Korea) with an oocyte filter (100 µm nylon screen; Mini IVF Filter, WTA) and classified four groups: 1 (excellent), 2 (fair), 3 (poor) and 4 (dead). Immediately after aspiration, a single technician evaluated the COCs using the most common criteria used to select and classify a standard collection of bovine oocytes [34–36].

### ***In vitro* embryo production procedures: media and culture conditions**

For *in vitro* maturation (IVM), the COCs were cultured for 22 h in 450 µL of TCM-199 media that contained 0.005 AU/mL FSH (F2293, Sigma-Aldrich, St. Louis, MO, USA), 10% fetal bovine serum ([FBS] GIB16000-044, Thermo Fisher, Waltham, MA, USA), 1 µg/mL 17β-estradiol (E4389, Sigma-Aldrich), and 100 µM cysteamine (M6500, Sigma-Aldrich). The IVM cultures incubated under a humidified environment with 5% CO<sub>2</sub> at 38.5°C. For *in vitro* fertilization (IVF), sperm preparation was conducted using BoviPure® Gradient following the manufacturer's instructions (Nidacon, Gothenburg, Sweden) [37]. Layering 2 mL of BoviPure® bottom medium with 2 mL of BoviPure® top medium was meticulously done in a 15 mL centrifuge tube. Following this, thawed semen (500 µL) was mixed with BoviPure® extender in a warm test tube at a 1:1 ratio. The prepared semen (800 µL) was then gently loaded onto the top of the gradient and

centrifuged at 1,500 rounds per minute (RPM) for 20 minutes. After centrifugation, the liquid above the sperm pellet was carefully removed. Subsequently, the pellet was resuspended in 5 mL of BoviWash and centrifuged at 1700 RPM for 5 minutes. The resulting pellet was resuspended in 100  $\mu$ L of IVF medium (VibroFert™, ART Lab Solutions, Adelaide, Australia). The supplements found in the IVF medium consist of 10 IU/mL of heparin, 25 mM of penicillamine, 12.5 mM of hypotaurine, and 1.25 mM of epinephrine. Finally, on day 0, the oocytes were inseminated with  $1-2 \times 10^6$  spermatozoa/mL for 18 hours in an IVF medium within a humidified atmosphere of 5% CO<sub>2</sub> at 38.5°C. Following co-culture of COC with sperm, referred to as day 1, potential zygotes were mechanically cleared of cumulus cells by repeated pipetting into wash medium. They were then washed once in cleavage medium (VibroCleave™, ART Lab Solutions) and six embryos were placed in 20  $\mu$ L drops of pre-conditioned cleavage medium covered with paraffin oil. On day 5, embryos were washed once in blastocyst medium (VibroBlast™, ART Lab Solutions) and groups of six embryos were transferred to 20  $\mu$ L drops of pre-equilibrated blastocyst medium, also covered with paraffin oil. Embryos were then cultured until day 8. All maintained at 38.5°C in an atmosphere consisting of 5% O<sub>2</sub>, 5% CO<sub>2</sub> and 90% N<sub>2</sub>.

### Statistical analysis

GraphPad Prism (version 10.2.0, GraphPad Software, Boston, MA, USA) was applied for statistical analysis in this study. Quantity of follicles, retrieved oocytes and transferable embryos according to the AMH level were analyzed by two way-ANOVA, and the level of significance was  $p < 0.05$ , then the outcomes was displayed as the mean  $\pm$  SEM. Regression analysis was used for the correlation between total quantity of follicles and plasma AMH concentration, and the larger the slope of the regression curve.

## RESULTS

### Plasma anti-Müllerian hormone concentration, classification, and selection of donor

Among the 42 Hanwoo donors, concentrations of AMH in plasma varied between 276.5 and 2,212.5 pg/mL, with a mean AMH concentration ( $\pm$  SEM) of  $820.8 \pm 172.4$  pg/mL (Table 1). Based on these concentrations, we categorized the donors into three groups: the H group (top 30% of donors,  $n = 9$ ), M group (next 30% of donors,  $n = 9$ ), and L group (lowest 40% of donors,  $n = 12$ ), with the control group being the 12 donors closest to the mean concentration. The AMH concentrations for each group were as follows: H group ( $1,484.9 \pm 129.2$  pg/mL), M group ( $775.5 \pm 39.0$  pg/mL), L group ( $371.6 \pm 17.8$  pg/mL), and control group ( $806.1 \pm 16.6$  pg/mL). The range of AMH concentrations in each group of Hanwoo donors is shown in Fig. 2. Furthermore, the age of donors in each group was as follows: H group (age,  $4.1 \pm 0.7$  years), M group (age,  $4.3 \pm 0.8$  years), L group (age,  $4.1 \pm 0.6$  years), and control group (age,  $3.9 \pm 0.6$  years) with age differences between groups not significant.


**Table 1.** Summary of anti-Müllerian hormone (AMH) concentration, antral follicle count, oocyte retrieval, and transferable embryos production<sup>1)</sup>

No. of donors	No. of session	AMH (pg/mL)	AFC (n)	COCs (n)	TE (n)	TE rate (%)
42	252	$820.8 \pm 172.4$	$16.8 \pm 2.1$	$11.8 \pm 1.7$	$5.1 \pm 1.0$	$40.9 \pm 4.2$

<sup>1)</sup>Values are presented as means  $\pm$  SEM.

AFC, antral follicle count; COCs, cumulus-oocyte complexes; TE, transferable embryo.





**Fig. 2.** Distribution of concentrations of anti-Müllerian hormone (AMH) in donors in the three superstimulation treatment groups (n = 30). H group (highest 30% of donors, n = 9), M group (next 30% of donors, n = 9), L group (lowest 40% of donors, n = 12), and control group (mean AMH concentration for all donors, n = 12).

**Comparison of ultrasound-monitored follicles, retrieved oocytes, and transferable embryos by group**

In total, 252 OPU sessions were conducted across both the superstimulated and control groups. AMH concentration, AFC, oocyte retrieval, and transferable embryo production by donors are provided in Table 2. We divided the cows into control and superstimulation groups. The AMH concentration was  $806.1 \pm 16.6$  pg/mL in the control group and  $826.8 \pm 172.4$  pg/mL in the superstimulated group, which was not significantly different between the two groups. However, the difference in the number of AFCs increased by 3.8, from  $14.1 \pm 1.9$  in the control group to  $17.9 \pm 1.3$  in the superstimulated group, and the difference in the number of COCs increased by 3.3, from  $9.5 \pm 1.6$  in the control group to  $12.8 \pm 1.0$  in the superstimulated group. The difference in the number of transferable embryos increased by 1.3, from  $4.1 \pm 0.9$  in the control group to  $5.4 \pm 0.6$  in the superstimulated group, but the difference in the transferable embryo rate was not statistically significant, from  $40.0 \pm 4.5$  in the control group to  $41.3 \pm 2.6$  in the superstimulated group. The superstimulated group was divided into three groups according to AMH concentration to evaluate correlation of AFC, oocyte retrieval, and transferable embryo production with AMH concentration. The AFC confirmed by ultrasound monitoring during OPU in groups H, M, L, and the control group are presented in Fig. 3A. Significantly higher average quantities of follicles with diameters ranging from 1 to 15 mm were observed in groups H and M compared to the control group ( $p$


**Table 2.** Comparison of ovum pick-up (OPU)- *in vitro* embryo production (IVEP) outcomes with and without superstimulation in Hanwoo cows segmented by plasma AMH level<sup>1)</sup>

	Control group <sup>2)</sup>	Superstimulated group	High AMH $\geq 70\%$	Medium $40\% < \text{AMH} < 70\%$	Low AMH $\leq 40\%$
AMH (pg/mL)	806.1 $\pm$ 16.6	826.8 $\pm$ 172.4	1,484.8 $\pm$ 129.2	775.5 $\pm$ 39.0	371.6 $\pm$ 17.8
AFC (n)	14.1 $\pm$ 1.9	17.9 $\pm$ 1.3	23.4 $\pm$ 2.2	21.2 $\pm$ 1.8	11.4 $\pm$ 0.8
COCs (n)	9.5 $\pm$ 1.6	12.8 $\pm$ 1.0	18.9 $\pm$ 1.8	13.2 $\pm$ 1.0	7.9 $\pm$ 0.5
TE (n)	4.1 $\pm$ 0.9	5.4 $\pm$ 0.6	7.4 $\pm$ 1.2	6.9 $\pm$ 1.1	2.9 $\pm$ 0.3
TE rate (%)	40.0 $\pm$ 4.5	41.3 $\pm$ 2.6	36.8 $\pm$ 4.1	51.1 $\pm$ 5.7	36.8 $\pm$ 3.0

<sup>1)</sup>Values are presented as means  $\pm$  SEM.

<sup>2)</sup>Control group (non-superstimulated) (n = 12), Superstimulated group (n = 30).

AMH, anti-Müllerian hormone; AFC, antral follicle count; COCs, cumulus-oocyte complexes; TE, transferable embryo.




**Fig. 3.** Numbers of follicles, oocytes, and transferable embryos observed by ultrasound in ovum pick-up (OPU) trials. (A) Number of follicles per group according to anti-Müllerian hormone (AMH) concentration, (B) number of oocytes collected per group according to AMH concentration, and (C) number of transferable embryos per group according to AMH concentration. a, significant difference ( $p < 0.05$ ); b, significant difference ( $p < 0.01$ ); c, significant difference ( $p < 0.001$ ).

$< 0.05$ ). However, no significant difference was noted between the control group and group L. Furthermore, the average quantity of follicles in groups H and M was significantly higher than that in group L ( $p < 0.05$ ), with no significant difference between groups H and M. Fig. 3B displays the number of oocytes recovered after OPU. The average quantity of retrieved oocytes was significantly higher in group H than in the control group ( $p < 0.05$ ), whereas no significant difference was noted between control and groups M and L. Moreover, there was a significant difference in the average quantity of retrieved oocytes among groups H, M, and L ( $p < 0.05$ ). Transferable embryos derived from oocytes retrieved after OPU are shown in Fig. 3C. No significant difference was observed in the quantity of transferable embryos between the three AMH level groups and the control group. However, in the groups in which AMH levels were measured, the H and M groups demonstrated a significant difference from the L group ( $p < 0.05$ ).

### Correlation between anti-Müllerian hormone and the quantity of follicles, retrieved oocytes, and the proportion of transferable embryos

Individuals exhibiting higher levels of AMH tend to possess a greater quantity of antral follicles, as evidenced by a strong positive correlation observed between AMH level and follicle quantity. This correlation was statistically significant ( $R^2 = 0.5785$ ,  $p < 0.0001$ ); as shown in Fig. 4A. Similarly, a strong positive correlation was evident between AMH concentrations and the quantity of retrieved oocytes, suggesting that individuals with higher AMH levels tend to have a greater quantity of retrieved oocytes. This correlation was statistically significant ( $R^2 = 0.6857$ ,  $p < 0.0001$ ); as shown



**Fig. 4. Correlations between plasma anti-Müllerian hormone (AMH) concentration and number of follicles, collected oocytes, embryo quantity, and transferable embryo rate.** (A) Correlation between total follicle count and plasma AMH concentration ( $R^2 = 0.5785$ ,  $p < 0.0001$ ). (B) Correlation between total oocytes retrieved and AMH concentration in plasma ( $R^2 = 0.6857$ ,  $p < 0.0001$ ). (C) Correlation between total number of transferable embryos and AMH concentration in plasma ( $R^2 = 0.4049$ ,  $p < 0.0001$ ). (D) Correlation between transferable embryo rate and AMH concentration in plasma ( $R^2 = 0.1476$ ,  $p = 0.5225$ ).

in Fig. 4B. The total number of transferable embryos and plasma AMH concentration were also correlated. ( $R^2 = 0.4049$ ,  $p < 0.0001$ ), as shown in Fig. 4C. When the correlation between AMH levels and the proportion of transferable embryos was assessed, a weak positive correlation was observed in individuals with higher AMH levels. This correlation was not significant ( $R^2 = 0.1476$ ,  $p = 0.5225$ ) as shown in Fig. 4D.

## DISCUSSION

Efficient embryo production and optimal donor selection for OPU in beef cattle are crucial to save labor and time. Recently, Ghanem et al. [30] reported that plasma AMH profiles correlated with AFC after random-cycle OPU in Hanwoo cows, as well as with the retrieval of oocytes, suggesting that AMH could serve as a useful indicator of donor selection. Therefore, in this study, we aimed to validate the hypothesis of predicting donor selection using AMH testing under conditions that induce superstimulation of follicles. Consistent with previous studies, the precise synchronization of follicle waves and induction of superstimulation, as conducted in this study, are the most effective production methods [19,38,39]. Controlled internal drug release (CIDR) was used in all donors to maintain stable  $P_4$  concentrations, with 12 AU FSH administered over four treatments within two days. Overall, the cattle responded well to superstimulation induction with six rounds of OPU per cow. The average rate of COCs recovery was  $11.8 \pm 1.7$ , the average quantity of transferable embryos was  $5.1 \pm 1.0$ , and the ratio of transferable embryos was  $40.9 \pm 4.2$  per cow. These efficient rates of oocyte production were consistent with those reported in previous studies, making a comparison between superstimulation and AMH feasible.

Previous studies have shown that repeated AMH tests provide information that is very similar to that provided by a single test [40]. Our study involved conducting a single AMH test, in line with the understanding that a single measurement of AMH concentration offers adequate information to estimate ovarian reproductive capacity. While previous studies have indicated a pattern of increasing



and decreasing AFC up to the age of five years in both beef and dairy cows, our study used donor averaging  $4.1 \pm 0.6$  years old. This suggests that the use of a single threshold value for AMH level is practical for selecting donors, making it possible to predict ovarian reserves and overproduction capacity using AMH testing, which is a significant advantage in real-world applications. AMH concentrations vary widely between animal breeds. A comparison between dairy and beef breeds showed that among dairy breeds, *Bos indicus* (Nelore breed), which is known for its high genetic AFC, exhibits higher AMH levels than other dairy breeds, including Holsteins, Jerseys, and crossbred cattle following in ascending order [30,41–43]. Thus, depending on the species, characteristics, and age of the animal, AMH is an important indicator of AFC, oocyte recovery, and follicular production capacity [44,45]. Hanwoo cattle are commonly used for beef production in Republic of Korea. This study measured AMH levels in Hanwoo donors and categorized them into high, medium, and low groups according to their concentration. These concentrations varied within the range of 276.5 to 2212.5 pg/mL, and the average AMH concentrations for each group were as follows: H group,  $1,484.9 \pm 129.2$  pg/mL; MH group,  $775.5 \pm 39.0$  pg/mL; L group,  $371.6 \pm 17.8$  pg/mL. This indicates a higher tendency for AMH concentrations in Hanwoo cows aimed at beef production than in breeds with reduced breeding capacity, and similar levels were observed in dairy breeds [42,43]. In conclusion, Hanwoo cattle belong to the *Bos taurus* lineage and exhibit AMH levels more in line with those of *Bos indicus* breeds, supporting previous research that Hanwoo cows, even though they are beef breeds, display similar tendencies as Holstein dairy cows [44–46].

The strong association between the AMH concentration in the plasma, the AFC, the number of COCs and the capacity to produce embryos has been confirmed in several animal species [27–29,47,48]. According to Widodo et al. [49], Holstein AMH concentration positively correlates with number of COCs and embryos from individual OPU donors. Consistent with these findings, our study revealed a positive correlation between AMH concentrations and the quantity of superstimulated antral follicles ( $R^2 = 0.58$ ) and the number of retrieved COCs ( $R^2 = 0.69$ ), indicating that higher numbers are associated with higher AMH levels. Furthermore, although a positive correlation was observed between AMH levels and the quantity of transferable embryos ( $R^2 = 0.40$ ), there was a weak positive correlation between high AMH levels and the proportion of transferable embryos in each group, albeit statistically insignificant ( $R^2 = 0.15$ ). Recently, Ghanem et al. [30] reported a high correlation between AMH profiles and AFC, quantity of retrieved COCs, and number of embryos produced by each donor in Hanwoo with random estrus cycle. Similarly, Batista et al. [43] reported that high concentrations of AMH in Nelore and Holstein cows showed a strong correlation with AFC and retrieved COCs in donors of these breeds.

In line with other results, AMH concentration proved to be very accurate in predicting ovarian production associated with standardized superstimulation protocols [40,50–53]. Thus, strong correlations between AMH concentration and AFC, COC count, and the number of transferable embryos suggest that increasing the superstimulation response enhances these correlations. In this study, we divided the donors into three groups according to AMH concentration. Comparing the AFC, COCs and number of transferable embryos between each group, we found that group H had a 105.26% increase in AFC, 139.24% increase in COCs and 155.17% increase in transferable embryos compared to group L. Compared to group M, group H had a 10.37% increase in AFC, 43.13% increase in COCs and 7.2% increase in transferable embryos. The M group also showed an 85.96% increase in AFC, 67.08% increase in COCs and 137.93% increase in transferable embryos compared to the L group. Therefore, selecting high-AMH donors using the same housing, feeding, FSH, and labor costs would result in nearly half the cost of embryo production compared to cows that do not consider AMH levels. Consequently, considering AMH levels in donor selection can be a strategic approach for reliable donor selection by embryo production and transfer specialists,

offering a significant advantage in reducing embryo production costs.

In summary, AMH shows a strong correlation with the response to superstimulation and the potential for embryo production in each donor. Categorizing AMH concentrations into different groups revealed a strong correlation between high AMH levels and AFC, the number of retrieved COCs, and the number of embryos produced by individual donor cows. Therefore, the results of this study provide a valuable practical method for enhancing the efficiency of Hanwoo donor cow selection and embryo transfer programs during the superstimulation response protocol and OPU procedures, indicating that AMH testing could serve as a reliable indicator for predicting the IVEP capacity of Hanwoo donors.

## References

1. Giordano JO, Fricke PM, Wiltbank MC, Cabrera VE. An economic decision-making support system for selection of reproductive management programs on dairy farms. *J Dairy Sci.* 2011;94:6216-32. <https://doi.org/10.3168/jds.2011-4376>
2. Giordano JO, Kalantari AS, Fricke PM, Wiltbank MC, Cabrera VE. A daily herd Markov-chain model to study the reproductive and economic impact of reproductive programs combining timed artificial insemination and estrus detection. *J Dairy Sci.* 2012;95:5442-60. <https://doi.org/10.3168/jds.2011-4972>
3. Kalantari AS, Cabrera VE. The effect of reproductive performance on the dairy cattle herd value assessed by integrating a daily dynamic programming model with a daily Markov chain model. *J Dairy Sci.* 2012;95:6160-70. <https://doi.org/10.3168/jds.2012-5587>
4. Galvão KN, Federico P, De Vries A, Schuenemann GM. Economic comparison of reproductive programs for dairy herds using estrus detection, timed artificial insemination, or a combination. *J Dairy Sci.* 2013;96:2681-93. <https://doi.org/10.3168/jds.2012-5982>
5. Overton MW, Cabrera VE. Monitoring and quantifying the value of change in reproductive performance. In: Beede DK, editor. *Large dairy herd management*. Champaign, IL: American Dairy Science Association; 2017. p. 549-64.
6. Ricci A, Li M, Fricke PM, Cabrera VE. Short communication: economic impact among 7 reproductive programs for lactating dairy cows, including a sensitivity analysis of the cost of hormonal treatments. *J Dairy Sci.* 2020;103:5654-61. <https://doi.org/10.3168/jds.2019-17658>
7. De Vries A, Kaniyamattam K. A review of simulation analyses of economics and genetics for the use of in-vitro produced embryos and artificial insemination in dairy herds. *Anim Reprod.* 2020;17:e20200020. <https://doi.org/10.1590/1984-3143-AR2020-0020>
8. Guerreiro BM, Batista EOS, Vieira LM, Sá Filho MF, Rodrigues CA, Castro Netto A, et al. Plasma anti-mullerian hormone: an endocrine marker for in vitro embryo production from *Bos taurus* and *Bos indicus* donors. *Domest Anim Endocrinol.* 2014;49:96-104. <https://doi.org/10.1016/j.domaniend.2014.07.002>
9. van Wagtenonk-de Leeuw AM. Ovum pick up and in vitro production in the bovine after use in several generations: a 2005 status. *Theriogenology.* 2006;65:914-25. <https://doi.org/10.1016/j.theriogenology.2005.09.007>
10. Rizos D, Clemente M, Bermejo-Alvarez P, De La Fuente J, Lonergan P, Gutiérrez-Adán A. Consequences of in vitro culture conditions on embryo development and quality. *Reprod Domest Anim.* 2008;43:44-50. <https://doi.org/10.1111/j.1439-0531.2008.01230.x>
11. Telfer EE, Sakaguchi K, Clarkson YL, McLaughlin M. In vitro growth of immature bovine follicles and oocytes. *Reprod Fertil Dev.* 2019;32:1-6. <https://doi.org/10.1071/RD19270>

12. Lonergan P, Fair T. Maturation of oocytes in vitro. *Annu Rev Anim Biosci.* 2016;4:255-68. <https://doi.org/10.1146/annurev-animal-022114-110822>
13. Blondin P, Sirard MA. Oocyte and follicular morphology as determining characteristics for developmental competence in bovine oocytes. *Mol Reprod Dev.* 1995;41:54-62. <https://doi.org/10.1002/mrd.1080410109>
14. Hirayama H, Kageyama S, Naito A, Fukuda S, Fujii T, Minamihashi A. Prediction of superovulatory response in Japanese Black cattle using ultrasound, plasma anti-Müllerian hormone concentrations and polymorphism in the ionotropic glutamate receptor AMPA1/GRIA1. *J Reprod Dev.* 2012;58:380-3. <https://doi.org/10.1262/jrd.11-129S>
15. Hirata TI, Sato M, Sasaki S, Sasaki O, Osawa T. Effect of suckling on embryo production by repeated ovum pick-up before and after timed artificial insemination in early postpartum Japanese black cows. *J Reprod Dev.* 2008;54:346-51. <https://doi.org/10.1262/jrd.20007>
16. Takuma T, Sakai S, Ezoe D, Ichimaru H, Jinnouchi T, Kaedei Y, et al. Effects of season and reproductive phase on the quality, quantity and developmental competence of oocytes aspirated from Japanese black cows. *J Reprod Dev.* 2010;56:55-9. <https://doi.org/10.1262/jrd.09-071H>
17. Hasler JF. Forty years of embryo transfer in cattle: a review focusing on the journal *Theriogenology*, the growth of the industry in North America, and personal reminiscences. *Theriogenology.* 2014;81:152-69. <https://doi.org/10.1016/j.theriogenology.2013.09.010>
18. Bó GA, Mapletoft RJ. Historical perspectives and recent research on superovulation in cattle. *Theriogenology.* 2014;81:38-48. <https://doi.org/10.1016/j.theriogenology.2013.09.020>
19. Behringer RR, Finegold MJ, Cate RL. Müllerian-inhibiting substance function during mammalian sexual development. *Cell.* 1994;79:415-25. [https://doi.org/10.1016/0092-8674\(94\)90251-8](https://doi.org/10.1016/0092-8674(94)90251-8)
20. Lee MM, Donahoe PK, Hasegawa T, Silverman B, Crist GB, Best S, et al. Mullerian inhibiting substance in humans: normal levels from infancy to adulthood. *J Clin Endocrinol Metab.* 1996;81:571-6. <https://doi.org/10.1210/jcem.81.2.8636269>
21. Laven JSE, Mulders AGMGJ, Visser JA, Themmen AP, de Jong FH, Fauser BCJM. Anti-Müllerian hormone serum concentrations in normoovulatory and anovulatory women of reproductive age. *J Clin Endocrinol Metab.* 2004;89:318-23. <https://doi.org/10.1210/jc.2003-030932>
22. van Rooij IAJ, Broekmans FJM, te Velde ER, Fauser BCJM, Bancsi LFJMM, de Jong FH, et al. Serum anti-Müllerian hormone levels: a novel measure of ovarian reserve. *Hum Reprod.* 2002;17:3065-71. <https://doi.org/10.1093/humrep/17.12.3065>
23. Durlinger ALL, Kramer P, Karels B, de Jong FH, Uilenbroek J TJ, Grootegoed JA, et al. Control of primordial follicle recruitment by anti-Müllerian hormone in the mouse ovary. *Endocrinology.* 1999;140:5789-96. <https://doi.org/10.1210/endo.140.12.7204>
24. Gruijters MJG, Visser JA, Durlinger ALL, Themmen APN. Anti-Müllerian hormone and its role in ovarian function. *Mol Cell Endocrinol.* 2003;211:85-90. <https://doi.org/10.1016/j.mce.2003.09.024>
25. Guibourdenche J, Lucidarme N, Chevenne D, Rigal O, Nicolas M, Luton D, et al. Anti-Müllerian hormone levels in serum from human fetuses and children: pattern and clinical interest. *Mol Cell Endocrinol.* 2003;211:55-63. <https://doi.org/10.1016/j.mce.2003.09.011>
26. Mossa F, Evans ACO. Review: the ovarian follicular reserve – implications for fertility in ruminants. *Animal.* 2023;17:100744. <https://doi.org/10.1016/j.animal.2023.100744>
27. Rico C, Fabre S, Médigue C, di Clemente N, Clément F, Bontoux M, et al. Anti-Müllerian hormone is an endocrine marker of ovarian gonadotropin-responsive follicles and can help to predict superovulatory responses in the cow. *Biol Reprod.* 2009;80:50-9. <https://doi.org/10.1093/biolreprod/80.1.50>

- org/10.1095/biolreprod.108.072157
28. Rico C, Médigue C, Fabre S, Jarrier P, Bontoux M, Clément F, et al. Regulation of anti-Müllerian hormone production in the cow: a multiscale study at endocrine, ovarian, follicular, and granulosa cell levels. *Biol Reprod.* 2011;84:560-71. <https://doi.org/10.1095/biolreprod.110.088187>
29. Monniaux D, Barbey S, Rico C, Fabre S, Gallard Y, Larroque H. Anti-Müllerian hormone: a predictive marker of embryo production in cattle? *Reprod Fertil Dev.* 2010;22:1083-91. <https://doi.org/10.1071/RD09279>
30. Ghanem N, Jin JI, Kim SS, Choi BH, Lee KL, Ha AN, et al. The anti-Müllerian hormone profile is linked with the in vitro embryo production capacity and embryo viability after transfer but cannot predict pregnancy outcome. *Reprod Domest Anim.* 2016;51:301-10. <https://doi.org/10.1111/rda.12681>
31. Merton JS, de Roos APW, Mullaart E, de Ruigh L, Kaal L, Vos PLAM, et al. Factors affecting oocyte quality and quantity in commercial application of embryo technologies in the cattle breeding industry. *Theriogenology.* 2003;59:651-74. [https://doi.org/10.1016/S0093-691X\(02\)01246-3](https://doi.org/10.1016/S0093-691X(02)01246-3)
32. Ferguson JD, Galligan DT, Thomsen N. Principal descriptors of body condition score in Holstein cows. *J Dairy Sci.* 1994;77:2695-703. [https://doi.org/10.3168/jds.S0022-0302\(94\)77212-X](https://doi.org/10.3168/jds.S0022-0302(94)77212-X)
33. Cardoso CJT, de Oliveira JS Jr, Kischel H, da Silva WAL, Arruda EDS, Souza-Cáceres MB, et al. Anti-Müllerian hormone (AMH) as a predictor of antral follicle population in heifers. *Anim Reprod.* 2018;15:12-6. <https://doi.org/10.21451/1984-3143-2017-AR887>
34. Seneda MM, Esper CR, Garcia JM, de Oliveira JA, Vantini R. Relationship between follicle size and ultrasound-guided transvaginal oocyte recovery. *Anim Reprod Sci.* 2001;67:37-43. [https://doi.org/10.1016/S0378-4320\(01\)00113-0](https://doi.org/10.1016/S0378-4320(01)00113-0)
35. Leibfried L, First NL. Characterization of bovine follicular oocytes and their ability to mature in vitro. *J Anim Sci.* 1979;48:76-86. <https://doi.org/10.2527/jas1979.48176x>
36. Hazeleger NL, Hill DJ, Stubbing RB, Walton JS. Relationship of morphology and follicular fluid environment of bovine oocytes to their developmental potential in vitro. *Theriogenology.* 1995;43:509-22. [https://doi.org/10.1016/0093-691X\(94\)00043-T](https://doi.org/10.1016/0093-691X(94)00043-T)
37. Samardzija M, Karadjole M, Getz I, Makek Z, Cergolj M, Dobranic T. Effects of bovine spermatozoa preparation on embryonic development in vitro. *Reprod Biol Endocrinol.* 2006;4:58. <https://doi.org/10.1186/1477-7827-4-58>
38. Bergfelt DR, Bo GA, Maplettoft RJ, Adams GP. Superovulatory response following ablation-induced follicular wave emergence at random stages of the oestrous cycle in cattle. *Anim Reprod Sci.* 1997;49:1-12. [https://doi.org/10.1016/S0378-4320\(97\)00064-X](https://doi.org/10.1016/S0378-4320(97)00064-X)
39. Bó GA, Baruselli PS, Moreno D, Cutaia L, Caccia M, Tríbulo R, et al. The control of follicular wave development for self-appointed embryo transfer programs in cattle. *Theriogenology.* 2002;57:53-72. [https://doi.org/10.1016/S0093-691X\(01\)00657-4](https://doi.org/10.1016/S0093-691X(01)00657-4)
40. Souza AH, Carvalho PD, Rozner AE, Vieira LM, Hackbart KS, Bender RW, et al. Relationship between circulating anti-Müllerian hormone (AMH) and superovulatory response of high-producing dairy cows. *J Dairy Sci.* 2015;98:169-78. <https://doi.org/10.3168/jds.2014-8182>
41. Moolhuijsen LME, Visser JA. Anti-Müllerian hormone and ovarian reserve: update on assessing ovarian function. *J Clin Endocrinol Metab.* 2020;105:3361-73. <https://doi.org/10.1210/clinem/dgaa513>
42. Ribeiro ES, Bisinotto RS, Lima FS, Greco LF, Morrison A, Kumar A, et al. Plasma

- anti-Müllerian hormone in adult dairy cows and associations with fertility. *J Dairy Sci.* 2014;97:6888-900. <https://doi.org/10.3168/jds.2014-7908>
43. Batista EOS, Macedo GG, Sala RV, Ortolan MDDV, Sá Filho MF, Del Valle TA, et al. Plasma antimullerian hormone as a predictor of ovarian antral follicular population in *Bos indicus* (Nelore) and *Bos taurus* (Holstein) heifers. *Reprod Domest Anim.* 2014;49:448-52. <https://doi.org/10.1111/rda.12304>
  44. Baldrighi JM, Sá Filho MF, Batista EOS, Lopes RNVR, Visintin JA, Baruselli PS, et al. Anti-Mullerian hormone concentration and antral ovarian follicle population in Murrah heifers compared to Holstein and Gyr kept under the same management. *Reprod Domest Anim.* 2014;49:1015-20. <https://doi.org/10.1111/rda.12430>
  45. Pfeiffer KE, Jury LJ, Larson JE. Determination of anti-Müllerian hormone at estrus during a synchronized and a natural bovine estrous cycle. *Domest Anim Endocrinol.* 2014;46:58-64. <https://doi.org/10.1016/j.domaniend.2013.05.004>
  46. Batista EOS, Guerreiro BM, Freitas BG, Silva JCB, Vieira LM, Ferreira RM, et al. Plasma anti-Müllerian hormone as a predictive endocrine marker to select *Bos taurus* (Holstein) and *Bos indicus* (Nelore) calves for in vitro embryo production. *Domest Anim Endocrinol.* 2016;54:1-9. <https://doi.org/10.1016/j.domaniend.2015.08.001>
  47. Monniaux D, Baril G, Laine AL, Jarrier P, Poulin N, Cognié J, et al. Anti-Müllerian hormone as a predictive endocrine marker for embryo production in the goat. *Reproduction.* 2011;142:845-54. <https://doi.org/10.1530/REP-11-0211>
  48. Lahoz B, Alabart JL, Monniaux D, Mermillod P, Folch J. Anti-Müllerian hormone plasma concentration in prepubertal ewe lambs as a predictor of their fertility at a young age. *BMC BMC Vet Res.* 2012;8:118. <https://doi.org/10.1186/1746-6148-8-118>
  49. Widodo OS, Nishihara S, Pambudi D, Kusakabe KT, Taura Y, Nishi Y, et al. Relationship between ovary size and anti-Müllerian hormone levels in Holstein-Friesian cows. *Front Vet Sci.* 2022;9:828123. <https://doi.org/10.3389/fvets.2022.828123>
  50. Umer S, Zhao SJ, Sammad A, Weldegebriall Sahl B, Pang YW, Zhu H. AMH: could it be used as a biomarker for fertility and superovulation in domestic animals? *Genes.* 2019;10:1009. <https://doi.org/10.3390/genes10121009>
  51. Hirayama H, Naito A, Fukuda S, Fujii T, Asada M, Inaba Y, et al. Long-term changes in plasma anti-Müllerian hormone concentration and the relationship with superovulatory response in Japanese Black cattle. *J Reprod Dev.* 2017;63:95-100. <https://doi.org/10.1262/jrd.2016-019>
  52. Sevgi R, Erdem H, Kardeşin T, Yılmaz MA, Satılmış M, Okuroğlu A, et al. Determination of the relationship between serum anti-Müllerian hormone level and superovulatory response in Simmental cows. *Reprod Domest Anim.* 2019;54:1322-9. <https://doi.org/10.1111/rda.13506>
  53. Alward KJ, Bohlen JF. Overview of anti-Müllerian hormone (AMH) and association with fertility in female cattle. *Reprod Domest Anim.* 2020;55:3-10. <https://doi.org/10.1111/rda.13583>