


Growth performance of broiler chickens fed diets containing granulated L-methionine compared with DL-methionine

June Hyeok Yoon^{1, #}, Su Hyun An^{2, #}, Yangsu Kim³, Hyeon-Jin Kim³, Changsu Kong^{1, 2, 4, *}



¹Department of Animal Science and Biotechnology, Kyungpook National University, Sangju 37224, Korea

²Research Institute for Innovative Animal Science, Kyungpook National University, Sangju 37224, Korea

³AA Tech Solution, CJ BIO, Seoul 04560, Korea

⁴Department of Animal Science, Kyungpook National University, Sangju 37224, Korea

Received: Feb 15, 2024

Revised: Apr 27, 2024

Accepted: May 1, 2024

[#]These authors contributed equally to this work.

*Corresponding author

Changsu Kong
Department of Animal Science and Biotechnology, Kyungpook National University, Sangju 37224, Korea
Tel: +82-54-530-1225
E-mail: changsukong@knu.ac.kr

Copyright © 2025 Korean Society of Animal Science and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

June Hyeok Yoon
<https://orcid.org/0000-0002-1862-1281>
Su Hyun An
<https://orcid.org/0000-0001-6236-6815>
Yangsu Kim
<https://orcid.org/0000-0001-7509-1988>
Hyeon-Jin Kim
<https://orcid.org/0000-0003-1641-0859>
Changsu Kong
<https://orcid.org/0000-0002-3876-6488>

Abstract

A novel granulated L-methionine (Met) has been developed using a simplified purification process, however its replacement with DL-Met has not yet been explored. The objective of the present study was to investigate the growth performance of broilers fed diets containing granulated L-Met (90% purity) compared to a diet containing DL-Met (99% purity). A total of 192 one-day-old broilers were allocated in four dietary treatments with six replicates (eight birds/cage) in a randomized complete block design based on body weight as the blocking factor. Twelve experimental diets were used, with four for each of the three growth stages: pre-starter (day 0 to 7), starter (day 7 to 21), and grower (day 21 to 28). The experimental diets consisted of: (1) a diet containing DL-Met at 100% of the digestible Met requirement, (2) a diet containing granulated L-Met at 85% of the digestible Met requirement, (3) a diet containing granulated L-Met at 90% of the digestible Met requirement, and (4) a diet containing granulated L-Met at the same inclusion rate (approximately 95% of the digestible Met requirement) as diet 1. The broilers were fed experimental diets during the pre-starter, starter, and grower stages, and growth performance was recorded by correcting mortality throughout the experiment period. Over the entire 28-day period, body weight gain and feed intake of broilers fed diets containing granulated L-Met increased linearly ($p < 0.05$) with an increase in dietary granulated L-Met supplementation. However, the growth performance of broilers fed diets containing granulated L-Met did not differ from those fed a diet containing DL-Met. The bioefficacy of L-Met relative to DL-Met for body weight gain and gain-to-feed ratio during the pre-starter stage was 116.9% and 104.0%, respectively. During the starter stage, the bioefficacy of L-Met relative to DL-Met was 127.5% and 111.0% for body weight gain and gain-to-feed ratio, respectively. Results of the present study reveal that the growth performance of broilers fed diets containing granulated L-Met was comparable to those fed a diet containing DL-Met, despite the lower dietary Met intake than digestible Met requirement. This suggests that L-Met might exhibit greater bioefficacy relative to DL-Met.

Keywords: L-methionine, DL-methionine, Growth performance, Broilers

Competing interests

No potential conflict of interest relevant to this article was reported. Yangsu Kim and Hyeon-Jin Kim are employees of CJ BIO (Seoul, Korea).

Funding sources

The authors received the product used in this study from CJ BIO (Seoul, Korea). This work was performed with the financial support from CJ BIO.

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: An SH, Kim Y, Kim HJ, Kong C.

Data curation: Yoon JH, An SH, Kim Y, Kim HJ, Kong C.

Formal analysis: Yoon JH, An SH.

Methodology: An SH, Kim Y, Kim HJ, Kong C.

Investigation: Yoon JH, An SH.

Writing - original draft: Yoon JH, An SH.

Writing - review & editing: Yoon JH, An SH, Kim Y, Kim HJ, Kong C.

Ethics approval and consent to participate

The animal study protocol was approved by the Kyungpook National University Institute for Animal Care and Use Committee, Korea (KNU 2021-0035).

INTRODUCTION

Methionine (Met) is the first-limiting amino acid (AA) in practical corn-soybean meal-based diets for broiler chickens [1]. DL-Met, which is a racemic mixture of D- and L-Met, has been commonly used to meet Met requirements in animal diets [2]. In fact, L-Met is the only biologically available form of Met that can be readily absorbed by animal intestinal cells and directly involved in protein synthesis [3]. However, D-Met must be converted to L-Met through enzymatic conversion processes in the liver and kidney [4]. Conversion of D-Met requires oxidative deamination by D-AA oxidase to produce α -keto- γ -methiolbutyric acid, which must then be transaminated by transaminases to form L-Met [5]. Therefore, it could be hypothesized that the incorporation of L-Met into the broiler diet may result in efficient utilization and protein synthesis, leading to improved growth performance of broilers compared to that of DL-Met.

Crystalline AA can be produced by bacterial fermentation, and the purification process is the first step toward obtaining pure crystalline AA [6]. During this process, specific crystalline AA is separated from a culture medium. However, newly developed granulated AA products were obtained through batch fermentation using a simplified purification process for use in swine and poultry production. Wensley et al. [7] reported that granulated tryptophan, threonine, and valine with respective biomass were equally bioavailable and usable as alternatives to feed-grade crystalline AA for growing pigs and broilers. However, to the best of our knowledge, there is limited literature investigating the growth performance of broilers fed diets containing newly developed granular L-Met compared to DL-Met commonly used in animal production. Therefore, the objective of the present study was to compare the growth performance of broilers fed diets containing granulated L-Met with DL-Met from day 1 to 28. The hypothesis of the current study was that broilers fed diets containing granulated L-Met would yield comparable growth performance to DL-Met even with lower dietary Met concentrations.

MATERIALS AND METHODS

Experimental procedures were reviewed and approved by the Kyungpook National University Institute for Animal Care and Use Committee, Korea (approval number: KNU 2021-0035).

Ingredient and dietary treatments

Four experimental diets were formulated and fed to broilers at each of three growth stages, i.e., pre-starter (day 0 to 7), starter (day 7 to 21), and grower (day 21 to 28) stages, resulting in 12 experimental diets in total (Table 1). At each stage, the experimental diets contained (1) a diet containing DL-Met (99% purity) at 100% of the digestible Met requirement (representing 100% of the digestible sulfur-containing AA [SAA] requirement), (2 and 3) diets containing granulated L-Met (90% purity, CJ BIO, Seoul, Korea) at 85% and 90% of the digestible Met requirement (representing 92% and 95% of the digestible SAA requirement), and (4) a diet containing granulated L-Met at the same inclusion rate (weight-to-weight) as diet 1 (representing approximately 95% and 97% for the digestible Met and SAA requirement, respectively). All experimental diets were formulated to meet or exceed the recommended concentrations of energy, nutrients, and AA, except for Met, according to Hoehler et al. [8] and Aviagen [9]. The Met sources (DL-Met and granulated L-Met) were assumed to be 100% standardized ileal digestible (SID). Within each growth stage, experimental diets were formulated in a single common batch to minimize unintended variations owing to potential mixing errors. Experimental diets within a growth stage had comparable ingredient compositions, except for cornstarch, glutamic acid,

Table 1. Ingredient and chemical compositions of experimental diets, as-fed basis¹⁾

Item	Digestible Met requirement											
	Pre-starter (day 1 to 7)				Starter (day 7 to 21)				Grower (day 21 to 28)			
	DL-Met		Granulated L-Met		DL-Met		Granulated L-Met		DL-Met		Granulated L-Met	
	100%	85%	90%	95%	100%	85%	90%	95%	100%	85%	90%	95%
Ingredient compositions (g/kg)												
Corn	556.0	556.0	556.0	556.0	550.0	550.0	550.0	550.0	540.0	540.0	540.0	540.0
Soybean meal	320.0	320.0	320.0	320.0	308.0	308.0	308.0	308.0	260.0	260.0	260.0	260.0
Cornstarch	9.4	9.4	9.5	9.4	44.1	44.1	44.1	44.1	99.1	99.0	99.2	99.1
Glutamic acid	8.9	9.5	9.1	8.9	7.0	7.5	7.3	7.0	7.1	7.7	7.3	7.1
Soybean oil	28.0	28.0	28.0	28.0	30.0	30.0	30.0	30.0	32.0	32.0	32.0	32.0
L-arginine	3.2	3.2	3.2	3.2	1.2	1.2	1.2	1.2	2.0	2.0	2.0	2.0
L-histidine	0.5	0.5	0.5	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
L-isoleucine	2.8	2.8	2.8	2.8	1.3	1.3	1.3	1.3	1.7	1.7	1.7	1.7
L-leucine	1.4	1.4	1.4	1.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
L-lysine-HCl	5.7	5.7	5.7	5.7	3.3	3.3	3.3	3.3	3.7	3.7	3.7	3.7
Granulated L-Met ²⁾	0.0	2.2	2.5	2.8	0.0	1.8	2.0	2.3	0.0	1.9	2.1	2.4
DL-Met	2.8	0.0	0.0	0.0	2.3	0.0	0.0	0.0	2.4	0.0	0.0	0.0
L-cysteine	2.3	2.3	2.3	2.3	1.3	1.3	1.3	1.3	1.4	1.4	1.4	1.4
L-phenylalanine	0.8	0.8	0.8	0.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
L-threonine	2.7	2.7	2.7	2.7	1.1	1.1	1.1	1.1	1.5	1.5	1.5	1.5
L-tryptophan	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1
L-valine	3.3	3.3	3.3	3.3	1.6	1.6	1.6	1.6	1.9	1.9	1.9	1.9
Limestone	15.8	15.8	15.8	15.8	14.4	14.4	14.4	14.4	13.5	13.5	13.5	13.5
Dicalcium phosphate	16.4	16.4	16.4	16.4	14.5	14.5	14.5	14.5	13.7	13.7	13.7	13.7
Salt	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5	4.5
Sodium bicarbonate	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0
Vitamin premix ³⁾	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
Mineral premix ⁴⁾	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0
Choline chloride	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
Calculated compositions (g/kg)												
AMEn (kcal/kg)	3,033	3,033	3,033	3,033	3,081	3,081	3,081	3,081	3,201	3,201	3,201	3,201
Crude protein	222.0	220.0	220.0	220.0	200.0	200.0	200.0	200.0	180.0	180.0	180.0	180.0
Calcium	9.6	9.6	9.6	9.6	8.7	8.7	8.7	8.7	8.1	8.1	8.1	8.1
Non-phytate phosphorus	4.8	4.8	4.8	4.8	4.3	4.3	4.3	4.3	4.0	4.0	4.0	4.0
SID Met	5.1	4.3	4.6	4.8	4.7	4.0	4.3	4.5	4.5	3.9	4.1	4.3
SID lysine	13.1	13.1	13.1	13.1	11.5	11.5	11.5	11.5	10.6	10.6	10.6	10.6
SID threonine	8.3	8.3	8.3	8.3	7.3	7.3	7.3	7.3	6.9	6.9	6.9	6.9
SID cysteine	4.6	4.6	4.6	4.6	4.0	4.0	4.0	4.0	3.8	3.8	3.8	3.8
SID tryptophan	2.1	2.1	2.1	2.1	1.9	1.9	1.9	1.9	1.7	1.7	1.7	1.7
Analyzed compositions (g/kg)												
Crude protein	210.6	215.1	214.9	220.9	198.7	199.5	197.5	203.3	182.3	186.0	185.4	187.1
Total Met	5.0	4.1	4.2	4.5	4.6	4.0	3.8	4.1	4.6	3.8	4.1	4.4

¹⁾Experimental diets consisted of: (1) diet containing DL-Met at 100% of the digestible Met requirement; (2,3) diets containing granulated L-Met at 85% and 90% of the digestible Met requirement; (4) diet containing granulated L-Met at same inclusion rate (weight-to-weight) as diet 1 (approximately 95% of the digestible Met requirement).

²⁾L-Met Pro®, CJ BIO (Seoul, Korea).

³⁾Supplies the following per kilogram of diet: retinyl acetate, 24,000 IU; cholecalciferol, 8,000 IU; DL- α -tocopherol acetate, 160 mg/kg; menadione nicotinamide bisulfite, 8 mg/kg; thiamine mononitrate, 8 mg/kg; riboflavin, 20 mg/kg; pyridoxine hydrochloride, 12 mg/kg; D-calcium pantothenate, 40 mg/kg; folic acid, 4 mg/kg; nicotinamide, 12 mg/kg.

⁴⁾Supplies the following per kilogram of diet: iron, 120 mg/kg; copper, 320 mg/kg; zinc, 200 mg/kg; manganese, 240 mg/kg; cobalt, 2 mg/kg; selenium, 0.6 mg/kg; iodine, 2.5 mg/kg. Met, methionine; AMEn, apparent nitrogen-corrected metabolizable energy; SID, standardized ileal digestible.

and Met sources; consequently, the supplemented Met intake was derived only from DL-Met or granulated L-Met.

Animal and experimental design

A total of 192 one-day-old male Ross 308 broiler chickens were obtained from a local hatchery (Samhwa Breeding, Hongseong, Korea) and tagged with identification numbers. On day 1, all birds were individually weighed and allocated to four dietary treatments with six replicates (eight birds/pen) in a randomized complete block design with body weight as the blocking factor using the Experimental Animal Allotment Program [10], as described by An and Kong [11]. Birds were fed experimental diets in mash form corresponding to the pre-starter, starter, and grower stages, and feed and water were offered *ad libitum* throughout the 28-day experimental period. The birds were housed in wire-floored battery cages (60 × 50 × 60 cm) in an environmentally controlled room under continuous light. The room temperature was maintained at 33°C for the first 3 days and gradually decreased by 2°C each week for 4 weeks [12].

Measurement and chemical analyses

On day 1, 7, 21, and 28 post-hatch, body weight, feed supply, and feed leftovers per cage were recorded. Using these data, body weight gain (BWG), feed intake (FI), and gain-to-feed ratio (G:F) were calculated by correcting the mortality of birds [13]. The experimental diets were ground in a mill grinder (CT293 Cyclotec, Foss, Hillerød, Denmark) through a 1.0-mm screen for nutrient analysis. The dry matter (method 930.15; [14]) and crude protein contents (method 990.03) in the diets were determined. Ingredients and experimental diets were analyzed for total AA contents (method 982.30 E [a and b]).

Statistical analysis

The data were analyzed using the GLM procedure of SAS (SAS Institute, Cary, NC, USA) with dietary treatments as a fixed variable in the model. Mean separation was performed with Tukey's adjustment for multiple comparisons. Orthogonal polynomial contrast coefficients were generated using the IML procedure of SAS. The linear and quadratic effects of dietary granulated L-Met supplementation were determined using orthogonal polynomial contrasts. Statistical significance was considered less than 0.05, and the experimental unit was a cage.

Quantitative estimates of the bioefficacy of L-Met relative to DL-Met were estimated using the standard curve methodology as described by Dilger and Baker [1]. Supplemented L-Met intake (g) was calculated by subtracting the dietary SID Met intake (g) for corn and soybean meal fractions from the total dietary SID Met intake (g). Response criteria of BWG and G:F (dependent variables) were regressed against the supplemented L-Met intake (independent variable) using linear regression analysis. Estimated supplemental L-Met intake (g) was determined to achieve growth performance equivalent to supplemented DL-Met intake with 100% of the digestible Met requirement. The estimated supplemental L-Met intake was obtained by interpolating the growth performance (BWG and G:F) of birds fed a diet containing DL-Met through the L-Met standard curve. The bioefficacy of L-Met relative to DL-Met was calculated by dividing the supplemented DL-Met intake by the estimated supplemental L-Met intake.

RESULTS

The growth performance of broilers in the pre-starter stage was not affected by the experimental diets (Table 2). The body weight at day 7 and BWG of broilers in the pre-starter stage increased

Table 2. Growth performance of broiler chickens fed diets containing methionine (Met) sources from day 1 to 28 of age^{1,2)}

Item	Digestible Met requirement				SEM	<i>p</i> -values ³⁾		
	DL-Met	Granulated L-Met				ANOVA	Linear	Quadratic
	100%	85%	90%	95%				
Pre-starter (day 1 to 7)								
BW at day 1 (g)	52.1	52.1	52.1	52.1	0.04	0.946	0.847	0.911
BW at day 7 (g)	181.3	174.9	182.0	183.5	2.59	0.141	0.035	0.390
BWG (g/bird)	129.2	122.9	129.8	131.4	2.60	0.147	0.036	0.407
FI (g/bird)	125.1	123.5	128.4	127.4	2.26	0.436	0.236	0.307
G:F (g:g)	1.04	0.99	1.01	1.03	0.013	0.146	0.071	0.840
Starter (day 7 to 21)								
BW at day 21 (g)	877.7 ^{ab}	849.0 ^b	879.9 ^{ab}	922.7 ^a	14.23	0.018	0.002	0.803
BWG (g/bird)	695.7 ^{ab}	673.6 ^b	696.7 ^{ab}	739.3 ^a	13.37	0.024	0.003	0.614
FI (g/bird)	827.7	818.4	843.6	871.2	14.06	0.083	0.018	0.991
G:F (g:g)	0.84	0.82	0.83	0.85	0.007	0.059	0.018	0.306
Grower (day 21 to 28)								
BW at day 28 (g)	1361.2	1315.2	1329.3	1398.5	21.38	0.064	0.015	0.338
BWG (g/bird)	483.5	466.2	449.3	475.8	13.24	0.329	0.614	0.205
FI (g/bird)	762.4	754.5	745.9	773.3	11.60	0.415	0.269	0.234
G:F (g:g)	0.63	0.62	0.60	0.62	0.010	0.198	0.834	0.231
Overall period (day 1 to 28)								
BWG (g/bird)	1,309.2	1,263.1	1,277.1	1,346.4	21.40	0.064	0.015	0.308
FI (g/bird)	1,703.2	1,688.5	1,709.3	1,767.6	26.05	0.194	0.049	0.564
G:F (g:g)	0.77 ^a	0.75 ^a	0.75 ^a	0.76 ^a	0.006	0.039	0.115	0.289

¹⁾Experimental diets consisted of: (1) diet containing DL-Met at 100% of the digestible Met requirement; (2,3) diets containing granulated L-Met at 85% and 90% of the digestible Met requirement; (4) diet containing granulated L-Met at same inclusion rate (weight-to-weight) as diet 1 (approximately 95% of the digestible Met requirement).

²⁾Each value represents least squares means of 6 replicate cages.

³⁾ANOVA, *p*-value for model of analysis of variance; Linear, *p*-value for linear effect of dietary granulated L-Met supplementation; Quadratic, *p*-value for quadratic effect of dietary granulated L-Met supplementation.

^{a,b}Least squares means within a row without a common superscript differ at *p* < 0.05.

BW, body weight; BWG, body weight gain; FI, feed intake; G:F, gain-to-feed ratio.

linearly ($p < 0.05$) in response to an increase in dietary granulated L-Met supplementation. During the starter stage, the body weight at day 21 and BWG were different ($p = 0.018$ and $p = 0.024$, respectively) between dietary treatments, however the body weight at day 21 and BWG of broilers fed a diet containing DL-Met did not differ from broilers fed diets containing granulated L-Met. The body weight at day 21, BWG, FI, and G:F of broilers in the starter stage increased linearly ($p < 0.05$) as dietary granulated L-Met increased. During the grower stage, the growth performance of broilers was not affected by the experimental diet. However, the body weight of broilers at day 28 increased linearly ($p = 0.015$) in response to an increase in dietary granulated L-Met supplementation. Throughout the 28-day experimental period, the growth performance of broilers fed diets containing granulated L-Met was not different from that of broilers fed a diet containing DL-Met, even at lower dietary Met concentration. However, the BWG and FI of broilers increased linearly ($p = 0.015$ and $p = 0.049$, respectively) as the dietary granulated L-Met supplementation increased.

Based on the growth performance of broilers fed a diet containing granulated L-Met in the present study, linear regression equations were established to determine an estimated supplemental L-Met intake for growth performance equivalent to that achieved with DL-Met supplementation

(Fig. 1). During the pre-starter stage, the bioefficacy of L-Met relative to DL-Met for BWG and G:F was 116.9% and 104.0%, respectively. The bioefficacy of L-Met relative to DL-Met for BWG and G:F was 127.5% and 111.0% in the starter stage. The bioefficacy of L-Met during the grower stage was not estimated due to the lack of linearity and quadraticity in BWG and G:F.

DISCUSSION

Met is involved in several important biochemical functions [15], including providing methyl groups for the methylation process [16], and can also serve as a precursor for antioxidant enzymes such as glutathione and taurine, which play vital roles in protecting cells from oxidative stress [17,18]. Additionally, as Met is one of the building blocks for protein and peptide synthesis in animals, Met deficiency can lead to reduced growth performance [18,19]. Wen et al. [20] reported that Met supplementation in a Met-deficient diet improved nitrogen retention and muscle protein accretion in pigs by increasing protein synthesis, leading to enhanced growth performance. Similarly, the growth performance of broilers fed Met-deficient diets in the present study increased linearly with increasing dietary Met supplementation.

According to our findings, the growth performance of broilers fed diets containing granulated L-Met did not differ from those fed a diet containing DL-Met, even though dietary Met intake




Fig. 1. Body weight gain and G:F of birds fed graded levels of granulated L-Met in Met-deficient diets. (A and B) The BWG and G:F of birds fed experimental diets during the pre-starter stage (day 1 to 7). (C and D) The BWG and G:F of birds fed experimental diets during the starter stage (day 7 to 21). Methionine-deficient diets contain granulated L-Met at approximately 85, 90, and 95% of the digestible Met requirement. Standard-curve analysis based on BWG and G:F (relative to DL-Met) showed bioefficacy of L-Met of 116.9% and 104.0% in the pre-starter stage, and 127.5% and 111.0% in the starter stage. Black triangle and red diamond symbols represent data from birds fed diets containing granulated L-Met and DL-Met, respectively. Each data represents least squares means of 6 replicate cages. Parentheses indicate the standard error of the slope and intercept of the regression equation. Bioefficacy of L-Met relative to DL-Met was calculated by dividing the supplemented DL-Met intake by the estimated supplemental L-Met intake. RMSE, root mean square error; Met, methionine; BWG, body weight gain; G:F, gain-to-feed ratio.

was lower than the digestible Met requirement. This may be partially because of the greater bioavailability of L-Met compared to DL-Met. The bioefficacy of L-Met relative to DL-Met in the present study was greater than 100% for BWG and G:F during pre-starter and starter stages. Therefore, the birds need to consume more than 100 units of DL-Met to achieve BWG and G:F values equivalent to those obtained by consuming 100 units of L-Met. These results could be explained by the fact that dietary L-Met can be directly incorporated into protein synthesis in intestinal cells [21], thereby improving the efficiency of L-Met for birds compared to DL-Met. Shen et al. [22] showed consistent results with the present study, reporting that the bioavailability of L-Met relative to DL-Met in young broiler chicks was 138.2% and 140.7% for average daily gain and G:F, respectively. Zhang et al. [23] conducted a slope-ratio assay to determine the bioavailability of L-Met in Pekin ducks and reported that the bioavailability of L-Met relative to DL-Met in the starter stage (day 1 to 14) was 137.6% and 121.0% for average daily gain and feed efficiency, respectively. Using eviscerated weight and breast muscle weight of birds as response criteria, the bioavailability of L-Met relative to DL-Met in 21-day-old broilers was 122.9% and 116.8%, respectively [24]. Additionally, the greater bioavailability of birds fed diets containing L-Met compared with those fed diets containing DL-Met may be partially due to L-Met improving redox status and intestinal development, which has been previously established in chicks [22] and turkeys [25].

However, previous studies have shown conflicting results between L-Met and DL-Met isomers. Dilger and Baker [1] conducted a standard-curve analysis to determine the efficacy values for DL-Met against L-Met in growing chicks (day 8 to 20), which were 102.8% and 119.9% for weight gain and G:F, respectively. Cenesiz et al. [26] performed a slope-ratio assay in 35-day-old broilers and reported that the relative bioavailability of L-Met to DL-Met was 123% and 91.5% for BWG and feed conversion ratio, respectively, whereas the relative bioavailability for breast meat yield was 88%. Kong et al. [27] noted that the response criteria for relative bioavailability may lead to conflicting results between Met isomers. The discrepancy in bioavailability between L- and DL-Met may also be partially due to differences in the age of birds [22]. The key enzyme, D-AA oxidase, is present only in the liver and kidney and is required to convert D-Met to L-Met [28]. D'Aniello et al. [29] reported that the expression levels of this enzyme were very low in young animals, suggesting that they may not readily utilize D-Met.

In conclusion, the growth performance of broilers fed diets containing granulated L-Met was not different from the broilers fed a diet containing DL-Met, despite the dietary Met intake being lower than the digestible Met requirement. These results of the present study might be attributed to the greater bioefficacy of L-Met compared to DL-Met.

References

1. Dilger RN, Baker DH. DL-Methionine is as efficacious as L-methionine, but modest L-cystine excesses are anorexigenic in sulfur amino acid-deficient purified and practical-type diets fed to chicks. *Poult Sci.* 2007;86:2367-74. <https://doi.org/10.3382/ps.2007-00203>
2. Kim BG, Lindemann MD, Rademacher M, Brennan JJ, Cromwell GL. Efficacy of DL-methionine hydroxy analog free acid and DL-methionine as methionine sources for pigs. *J Anim Sci.* 2006;84:104-11. <https://doi.org/10.2527/2006.841104x>
3. Fang Z, Luo H, Wei H, Huang F, Qi Z, Jiang S, et al. Methionine metabolism in piglets fed DL-methionine or its hydroxy analogue was affected by distribution of enzymes oxidizing these sources to keto-methionine. *J Agric Food Chem.* 2010;58:2008-14. <https://doi.org/10.1021/jf903317x>

4. Baker DH. Comparative species utilization and toxicity of sulfur amino acids. *J Nutr.* 2006;136:1670S-5S. <https://doi.org/10.1093/jn/136.6.1670S>
5. Hasegawa H, Shinohara Y, Akahane K, Hashimoto T. Direct detection and evaluation of conversion of D-methionine into L-methionine in rats by stable isotope methodology. *J Nutr.* 2005;135:2001-5. <https://doi.org/10.1093/jn/135.8.2001>
6. Leuchtenberger W, Huthmacher K, Drauz K. Biotechnological production of amino acids and derivatives: current status and prospects. *Appl Microbiol Biotechnol.* 2005;69:1-8. <https://doi.org/10.1007/s00253-005-0155-y>
7. Wensley MR, Woodworth JC, Derouchey JM, Dritz SS, Tokach MD, Goodband RD, et al. Effects of amino acid biomass or feed-grade amino acids on growth performance of growing swine and poultry. *Transl Anim Sci.* 2020;4:49-58. <https://doi.org/10.1093/tas/txz163>
8. Hoehler D, Lemme A, Ravindran V, Bryden WL, Rostagno HS. Feed formulation in broiler chickens based on standardized ileal amino acid digestibility. In: *Avances en Nutrición Acuicola VIII. VIII Simposium Internacional de Nutrición Acuicola*; 2006; Monterrey, Nuevo León, México.
9. Aviagen. Ross broiler: nutrition specification [Internet]. Aviagen. 2022 [cited 2024 Feb 14]. https://aviagen.com/assets/Tech_Center/Ross_Broiler/Ross-BroilerNutritionSpecifications2022-EN.pdf
10. Kim BG, Lindemann MD. A spreadsheet method for experimental animal allotment. *J Anim Sci.* 2007;85:112.
11. An SH, Kong C. Influence of age and type of feed ingredients on apparent and standardized ileal amino acid digestibility in broiler chickens. *J Anim Sci Technol.* 2022;64:740-51. <https://doi.org/10.5187/jast.2022.e43>
12. Cho I, An SH, Yoon JH, Namgung N, Kong C. Growth performance and nitrogen excretion of broiler chickens fed low protein diets supplemented with crystalline amino acids. *J Anim Sci Technol.* 2024;66:145-55. <https://doi.org/10.5187/jast.2023.e131>
13. An SH, Kong C. Standardized ileal digestible methionine requirements for 22 to 29-d-old male broilers based on the growth performance and feather development. *J Anim Sci Technol.* 2024;66:555-66. <https://doi.org/10.5187/jast.2023.e139>
14. AOAC [Association of Official Analytical Chemists] International. Official methods of analysis of AOAC International. 18th ed. Gaithersburg, MD: AOAC International; 2005.
15. Zhou J, Yue S, Xue B, Wang Z, Wang L, Peng Q, et al. Enhanced supply of methionine regulates protein synthesis in bovine mammary epithelial cells under hyperthermia condition. *J Anim Sci Technol.* 2021;63:1126-41. <https://doi.org/10.5187/jast.2021.e93>
16. Wu G. Amino acids: biochemistry and nutrition. Boca Raton, FL: CRC Press; 2013.
17. Zhang S, Gilbert ER, Saremi B, Wong EA. Supplemental methionine sources have a neutral impact on oxidative status in broiler chickens. *J Anim Physiol Anim Nutr.* 2018;102:1274-83. <https://doi.org/10.1111/jpn.12946>
18. Yang Z, Htoo JK, Liao SF. Methionine nutrition in swine and related monogastric animals: beyond protein biosynthesis. *Anim Feed Sci Technol.* 2020;268:114608. <https://doi.org/10.1016/j.anifeedsci.2020.114608>
19. Kim CH. Effect of essential amino acid deficient diets in feeding response and c-fos expression in rats brain in response to methionine deficiency. *J Anim Sci Technol.* 2002;44:727-38. <https://doi.org/10.5187/JAST.2002.44.6.727>
20. Wen C, Chen X, Chen GY, Wu P, Chen YP, Zhou YM, et al. Methionine improves breast muscle growth and alters myogenic gene expression in broilers. *J Anim Sci.* 2014;92:1068-73. <https://doi.org/10.2527/jas.2013-6485>

21. Saunderson CL. Effect of fasting and of methionine deficiency on L-methionine, DL-methionine and DL-2-hydroxy-4-methylthiobutanoic acid metabolism in broiler chicks. *Br J Nutr.* 1987;57:429-37. <https://doi.org/10.1079/BJN19870050>
22. Shen YB, Ferket P, Park I, Malheiros RD, Kim SW. Effects of feed grade L-methionine on intestinal redox status, intestinal development, and growth performance of young chickens compared with conventional DL-methionine. *J Anim Sci.* 2015;93:2977-86. <https://doi.org/10.2527/jas.2015-8898>
23. Zhang YN, Xu RS, Min L, Ruan D, Kim HY, Hong YG, et al. Effects of L-methionine on growth performance, carcass quality, feather traits, and small intestinal morphology of Pekin ducks compared with conventional DL-methionine. *Poult Sci.* 2019;98:6866-72. <https://doi.org/10.3382/ps/pez438>
24. Wang W, Wang J, Wu S, Dong X, Guo C, Zhang H, et al. Bioavailability of L-methionine relative to DL-methionine in broiler chickens. *Ital J Anim Sci.* 2019;18:1231-8. <https://doi.org/10.1080/1828051X.2019.1641433>
25. Park I, Pasquetti T, Malheiros RD, Ferket PR, Kim SW. Effects of supplemental L-methionine on growth performance and redox status of turkey poults compared with the use of DL-methionine. *Poult Sci.* 2018;97:102-9. <https://doi.org/10.3382/ps/pex259>
26. Çenesiz AA, Çiftci I, Ceylan N. Effects of DL- and L-methionine supplementation on growth performance, carcass quality and relative bioavailability of methionine in broilers fed maize-soybean-based diets. *J Anim Feed Sci.* 2022;31:142-51. <https://doi.org/10.22358/jafs/147800/2022>
27. Kong C, Park CS, Ahn JY, Kim BG. Relative bioavailability of DL-methionine compared with L-methionine fed to nursery pigs. *Anim Feed Sci Technol.* 2016;215:181-5. <https://doi.org/10.1016/j.anifeedsci.2016.03.011>
28. Bauriedel WR. The effect of feeding D-methionine on the D-amino acid oxidase activity of chick tissues. *Poult Sci.* 1963;42:214-7. <https://doi.org/10.3382/ps.0420214>
29. D'Aniello A, D'Onofrio G, Pischetola M, D'Aniello G, Vetere A, Petrucelli L, et al. Biological role of D-amino acid oxidase and D-aspartate oxidase: effects of D-amino acids. *J Biol Chem.* 1993;268:26941-9. [https://doi.org/10.1016/S0021-9258\(19\)74201-X](https://doi.org/10.1016/S0021-9258(19)74201-X)