

Control of alkaline phosphatase activity and pH stability by taurine in liquid boar semen

Wijesooriya Mudhiyanselage Nadeema Dissanayake¹,
Malavi Arachchi Gamage Nayodya Dananjane Malavi Arachchi Gamage¹,
Jung Min Heo², Young-Joo Yi^{1*}

¹Department of Agricultural Education, College of Education, Suncheon National University, Suncheon 57922, Korea

²Department of Animal Science and Biotechnology, College of Agriculture and Life Sciences, Chungnam National University, Daejeon 34134, Korea



Received: Aug 2, 2024
Revised: Sep 23, 2024
Accepted: Sep 26, 2024

*Corresponding author

Young-Joo Yi
Department of Agricultural Education,
College of Education, Suncheon
National University, Suncheon 57922,
Korea
Tel: +82-61-750-3352
E-mail: yij@scnu.ac.kr

Copyright © 2025 Korean Society of Animal Science and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Wijesooriya Mudhiyanselage Nadeema Dissanayake
<https://orcid.org/0000-0001-7130-7737>
Malavi Arachchi Gamage Nayodya Dananjane Malavi Arachchi Gamage
<https://orcid.org/0009-0009-3194-6548>
Jung Min Heo
<https://orcid.org/0000-0002-3693-1320>
Young-Joo Yi
<https://orcid.org/0000-0002-7167-5123>

Competing interests

No potential conflict of interest relevant to this article was reported.

Abstract

Alkaline phosphatase (ALP) is an enzyme present in various tissues and plays an important role in biochemical processes, by catalyzing the hydrolysis of organic phosphates and energy metabolism. Despite significant ALP activity in seminal plasma across species, its role in sperm function remains unclear. Several studies have revealed the important role of taurine in male reproductive functions, due to its antioxidant and membrane-stabilizing activity, and ability to enhance sperm quality. Therefore, this study examined the influence of taurine on ALP activity and pH stability in liquid boar semen during storage. Boar spermatozoa were exposed to different concentrations of ALP (0–5 IU/mL), and sperm motility, viability, acrosome integrity, and ALP activity were examined. In addition, liquid boar semen with varying concentrations of taurine (10–80 mM) was stored at 17°C for 7 days, and the ALP activity, pH level, and fertilization competence of spermatozoa were investigated through the storage period. Higher ALP activity was detected in fresh spermatozoa compared with capacitated spermatozoa. Motility, viability, and acrosome integrity decreased significantly in sperm incubated with 1–5 IU/mL ALP. An immunofluorescence assay revealed that ALP was localized on the acrosome, equatorial segment, and tail, and the fluorescence intensity indicated that ALP levels gradually decreased during storage. When the pH of liquid boar semen was maintained at 7.4 during storage, it showed higher ALP activity and sperm quality compared to sperm stored in a medium where pH was not maintained at a stable level. Also, when boar spermatozoa were stored in the diluent containing taurine, the pH and ALP activity were stable during the storage period. The generation of intracellular reactive oxygen species decreased in sperm stored with taurine, and higher levels of sperm motility and viability were observed. The expression of mRNA associated with fertilization competence was higher in the sperm stored with 40 mM taurine compared to that of sperm stored without taurine. Thus, it can be concluded that ALP activity and pH stability are crucial for maintaining sperm quality during liquid boar semen storage, and these factors can be regulated by the addition of taurine, suggesting its applicability in assisted reproductive techniques in mammals.

Keywords: Alkaline phosphatase, Liquid boar semen, pH, Taurine, Storage

Funding sources

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT) (RS-2023 00245276).

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Heo JM, Yi YJ.

Data curation: Dissanayake WMN, Yi YJ.

Formal analysis: Dissanayake WMN, Dananjane Malavi Arachchi Gamage MAGN.

Methodology: Dissanayake WMN, Yi YJ. Software: Dissanayake WMN, Dananjane Malavi Arachchi Gamage MAGN.

Validation: Dissanayake WMN, Dananjane Malavi Arachchi Gamage MAGN.

Investigation: Dissanayake WMN, Dananjane Malavi Arachchi Gamage MAGN.

Writing - original draft: Dissanayake WMN, Heo JM, Yi YJ.

Writing - review & editing: Dissanayake WMN, Dananjane Malavi Arachchi Gamage MAGN, Heo JM, Yi YJ.

Ethics approval and consent to participate

This article does not require IRB/IACUC approval because there are no human and animal participants.

INTRODUCTION

The preservation of boar semen in liquid form is crucial for successful artificial insemination (AI) practices in swine as it maintains sperm quality and has high fertilization rates over extended periods [1]. The extended storage of liquid boar semen often leads to a decline in sperm motility, viability, and acrosome integrity, primarily due to oxidative stress and unstable pH. Therefore, it is important to develop improved preservation methods to sustain the fertilizing capacity of spermatozoa during storage [2]. Alkaline phosphatase (ALP) an enzyme present on the surface of cell membranes, is crucial for the proper functioning of the male reproductive system [3,4]. ALP activity has been identified in the seminal fluid of several species, especially in boars, ALP is primarily secreted by the epididymis [5]. The enzyme is involved in hydrolyzing phosphate groups from different substrates and helps in transporting chemicals across membranes. Previous studies have demonstrated that ALP activity correlates with semen concentration and fertility, which indicates its potential role as a marker for sperm quality. Also, ALP has been associated with sperm maturation, since it facilitates the dephosphorylation process, which is important for sperm development and motility [6–8], suggesting its importance in maintaining sperm functionality. The pH of seminal fluid is another critical factor in sperm preservation. An alkaline pH helps protect sperm in acidic environments, and insufficient semen volume or reduced alkalinity can compromise this buffering capacity, negatively impacting fertility [9]. Maintaining proper pH is essential for sperm respiration and motility, and buffering agents are often added to semen diluents to stabilize the pH during storage [10]. These buffering agents neutralize the pH changes caused by sperm metabolism and ensure the optimal conditions of sperm functions such as motility and viability which are important for the fertilizing capacity [11].

Taurine, an amino acid naturally found in the body, plays a vital role in various mammalian organs. It is crucial for the growth and function of skeletal muscles [12], the detoxification of foreign substances [13], the stability of cell membranes [14], and the regulates the central nervous system [15]. The specific physiological roles of taurine have been reported to be antioxidation, immunoregulation, detoxification, osmoregulation, and neuromodulation [16,17]. Taurine has been evidenced to preserve sperm quality through the storage period in several species, including boar [18] and sheep [19]. It has been evidenced to reduce the production of reactive oxygen species (ROS) and enhance the rabbit sperm quality [20], rams [21], and bulls [22]. However, the effects of taurine on the preservation of boar semen in liquid form and its effect on ALP activity and pH have not been clarified yet. Therefore, the purpose of this study was to investigate the effect of taurine in regulating ALP activity and stabilizing pH to maintain sperm quality throughout the storage of liquid boar semen.

MATERIALS AND METHODOLOGY

Sample preparation

Boar semen was purchased from a local artificial insemination (AI) center, and only samples with an initial motility of over 80% were used for the experiment. The sperm samples were washed, reconstituted with Beltsville thawing solution (BTS; [23]), and stored at 17°C for 7 days. Unless otherwise stated, other all chemicals used in the present study were obtained from Sigma-Aldrich Chemical.

Experimental designs

Experiment 1: Sperm capacitation was induced by incubating the boar spermatozoa in Tyrode's

lactate-N-2-hydroxyethylpiperazine-N¹-2-ethanesulfonic acid-polyvinyl alcohol (PVA) medium with the addition of 5 mM sodium pyruvate, 11 mM glucose, and 2% bovine serum albumin (BSA) for 4 h at 38.5 °C, 5% CO₂ in the air [24]. The ALP activity was measured in fresh sperm and capacitated sperm, respectively.

Experiment 2: Liquid boar semen was incubated with or without different concentrations of ALP (0.5–5 IU/mL) at 37.5 °C for 2 h. Sperm motility, viability, and intact acrosome were examined after incubation.

Experiment 3: The effect of pH was evaluated by storing liquid boar semen at 17 °C for 7 days at a pH of 7.4 and compared with a group in which pH balance was not maintained (a pH-balance group *vs.* a control group without pH balance). To maintain the pH at 7.4 during the storage period in the pH-balance group, the pH of the semen was adjusted using pH adjustment solutions (HCl and NaOH; Sigma-Aldrich Chemical). The ALP activity, motility, and viability of the spermatozoa were examined on days 1, 3, 5, and 7.

Experiment 4: Liquid boar semen was stored in BTS containing different concentrations of taurine (10–100 mM). The level of pH, ALP activity, motility, viability, and production of intracellular ROS of spermatozoa were examined on days 1, 3, 5, and 7.

Measurement of alkaline phosphatase activity in boar spermatozoa

Spermatozoa (2×10^9 cells/ml) were washed with phosphate-buffered saline (PBS) by centrifugation for 3 min at 200×g (1730R, Labogene). The resulting sperm pellet was sonicated at 60 Hz in PBS for 10 sec (Daihan Scientific), then centrifuged at 13,000×g (Labogene) for 15 min at 4 °C. The protein concentration of the sperm extract was determined by Bradford's assay, and BSA was used as the protein standard. ALP activity was assessed using the Senso-Lyte[®] p-nitrophenyl phosphate (pNPP) ALP assay kit (AnaSpec) according to the manufacturer's guidelines. The standard curve was generated by performing twofold serial dilutions of the top standard to the concentrations of 100, 50, 25, 12.5, 6.2, 3.1, and 0 ng/mL. The assay to measure ALP activity was conducted by adding 100 µL sample and 50 µL of 5 mM pNPP solution to each well. The reaction was carried out at 25 °C for 1 h in a dark environment. Finally, 20 µL of stop solution was added to terminate the reaction. ALP activity was determined by monitoring the transformation of pNPP (colorless) into para-nitrophenol (yellow), which is measured by reading the absorbance at 405 nm using a microplate reader (Absorbance 96, Byoany). The enzyme activity was expressed as IU/mL.

Evaluation of sperm motility

Sperm motility was examined using a computer-assisted system (Sperm Class Analyzer[®], Microptic). A 2 µL of sperm was placed on a counting chamber (Leja products B.V.), and 10 separate fields were evaluated at 37.5 °C. At least 500 spermatozoa were analyzed per sample. The percentage of total motile sperm (%), progressively motile sperm (%), and hyperactive sperm (%) was analyzed.

Evaluation of sperm viability

Sperm viability was analyzed using the LIVE/DEAD[®] sperm viability kit (Molecular Probes), supplemented with DNA binding dyes SYBR14 (100 nM) and propidium iodide (PI; 10 µM). Sperm cells (1×10^8 cells/mL) were washed in PBS containing 0.1% PVA (PBS-PVA). Then spermatozoa were stained, and images were captured by a Nikon Eclipse Ci microscope (Nikon Instruments), a DS-Fi2 camera (Nikon), and imaging software (version 4.30, Nikon). Viable sperm cells exhibit green fluorescence (SYBR14) while dead sperm cells exhibit red fluorescence (PI).

Evaluation of acrosome integrity

Spermatozoa were fixed in 95% ethanol and incubated for 30 min at 4°C. Following fixation, the sperm were air-dried on the slides and stained for 10 min with fluorescein isothiocyanate-labeled *Pisum sativum* agglutinin (FITC-PSA; 5 µg/mL) [25]. A Fluorescence microscope and camera (Nikon), along with imaging software (version 4.30, Nikon) were used to analyze the acrosome integrity. Sperm heads displaying green fluorescence indicated an intact acrosome, while partial or no green fluorescence in the head indicated that the acrosome reacted or damaged spermatozoa.

Evaluation of intracellular reactive oxygen species in spermatozoa

The sperm cells were rinsed in 0.1% PBS-PVA and then incubated with 1 µM 5-(and-6) carboxy-2',7'-dichlorodihydrofluorescein diacetate (carboxy-H₂DCFDA, Invitrogen) at 37°C for 10 min. Thereafter, sperm cells were mounted on slides, and the ROS fluorescence intensity was analyzed using a fluorescence microscope (Nikon).

Immunofluorescence of boar spermatozoa

Spermatozoa were mounted on poly-L-lysine-treated coverslips in KMT medium (100 mM KCl, 2 mM MgCl₂, 10 mM TRIS-HCl, pH 7.0) and kept for 3 min to allow attachment. After that cells were fixed with 2% formaldehyde for 40 min at room temperature (RT), washed with PBS, followed by permeabilized in PBS and 0.1% Triton-X 100 (PBS-TX) for 40 min at RT. The blocking was done by PBS-TX with 5% normal goat serum for 25 min. Afterward, spermatozoa were incubated with mouse monoclonal immunoglobulin G (IgG) anti-ALP (A-10) antibody (1:100 dilution, #sc-271431, Santa Cruz Biotechnology) for 40 min. After washing with PBS-TX, cells were incubated with goat anti-mouse IgG (H+L) secondary antibody conjugated to FITC (626511, Invitrogen ThermoFisher Scientific) for 40 min at RT. DNA was stained with 4,6-diamidino-2-phenylindole (Molecular Probes), and images were captured using a Nikon fluorescence microscope.

Western blotting

Protein was extracted from the sperm pellets by boiling with loading buffer containing 50 mM Tris (pH 6.8), 150 mM NaCl, 2% sodium dodecyl sulfate, 20% glycerol, 5% b-mercaptoethanol, 0.02% bromophenol blue. Proteins were separated using 10% sodium dodecyl sulfate-polyacrylamide gel and electrophoretically transferred to polyvinylidene difluoride membranes (Bio-Rad Laboratories). Then the membranes were blocked with 5% skim milk in tris-buffered saline containing Tween-20 (TBS-T) for 1 h at RT, followed by overnight incubation with anti-ALP antibody (mouse monoclonal IgG, 1:1,000 dilution, #sc-271431, Santa Cruz Biotechnology) at 4°C. The membranes were then incubated with goat anti-mouse IgG-horseradish peroxidase secondary antibody (#31430, 1:10,000, ThermoFisher Scientific) for 1 h. at RT. The β-tubulin antibody (rabbit polyclonal IgG, 1:1000, #sc-9104, Santa Cruz) was used as a reference. Immunoreactive bands were observed using chemiluminescence reagents (SuperSignal™ West Femto, Thermo Fisher Scientific) and captured with an imaging system (Davinch-K).

Real-time polymerase chain reaction

Sperm samples were washed with PBS before extracting RNA with the PureLink™ RNA Mini Kit (Thermo Fisher Scientific), following slight modifications. RNA concentrations were quantified using a nanodrop spectrophotometer (DeNovix DS-11FX, DeNovix). According to the manufacturer's instructions, complementary DNA (cDNA) was synthesized from the extracted RNA using the TOYOBO ReverTra Ace qPCR RT kit (TOYOBO). Quantitative real-time polymerase chain

reaction (qRT-PCR) was conducted using SYBR™ Premix Ex Taq™ II (Bioneer) on a MyGo Pro PCR cycler (Diagnostic Technology). The expression of the target gene mRNA was quantified and normalized against glyceraldehyde-3-phosphate dehydrogenase, as the internal reference gene. The primer sequences for target genes of outer dense fiber of sperm tails protein 2 (ODF2), zona pellucida binding protein 2 (ZPBP2), and A-kinase anchor proteins 3 (AKAP3) and 4 (AKAP4) were designed using Primer-BLAST software from the National Center for Biotechnology Information (NCBI), Bethesda, USA. (<https://www.ncbi.nlm.nih.gov/tools/primer-blast/>) (Table 1).

Statistical analysis

The experimental data were displayed as mean \pm SEM and were statistically analyzed using one-way ANOVA in GraphPad PRISM® (GraphPad Software). A completely randomized design was employed, followed by Tukey's test for multiple comparisons across treatment groups. Additionally, sperm motility, viability, and ALP activity between the two pH groups were analyzed using an unpaired two-tailed t-test. Statistical significance was defined at * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$.

RESULTS

ALP activity was measured in fresh and capacitated spermatozoa, respectively. The ALP activity of fresh boar spermatozoa was 2.81 ± 0.4 IU/mL, whereas capacitated spermatozoa showed reduced ALP activity of 0.57 ± 0.2 IU/mL (Fig. 1A; $p < 0.01$). To evaluate the effect of ALP, the sperm cells were exposed to different concentrations of ALP (0.5–5 IU/mL) at 37.5 °C for 2 h. Sperm incubated with ALP exhibit a dose-dependent reduction in motility compared to the sperm without (W/O) ALP (84.6% in control [W/O] vs. 68.5%–81.3% in 0.5–5 IU/mL ALP, $p < 0.05$ & $p < 0.001$; Fig. 1B). The viability percentage was reduced significantly in the sperm incubated with 1–5 IU/mL ALP compared to that of sperm incubated without ALP (77.3% in control vs. 56.6%–63.5% 1–5 IU/mL ALP, $p < 0.01$ & $p < 0.001$; Fig. 1C). Intact acrosome percentage was lower in sperm samples incubated with 1–5 IU/mL ALP (57.9% in control vs. 39.5%–47.3% at 1–5 IU/mL ALP, $p < 0.01$, $p < 0.001$; Fig. 1D) compared to the control samples.

Spermatozoa were treated with different concentrations of ALP (0.5–5 IU/mL), and then sperm proteins were extracted. Western blotting with an anti-ALP antibody detected specific bands around 70 kDa (Fig. 2A). Analysis revealed significant variations in ALP levels among the five different treatment groups. ALP expression was significantly reduced in samples treated with 2–5 IU/mL of ALP compared to the control group (without ALP), with significant differences at $p < 0.05$ and $p < 0.01$ (Fig. 2B).

Localization of ALP in spermatozoa was assessed by immunofluorescence assay using an anti-

Table 1. Nucleotide sequences of primers for real-time PCR (RT-PCR)

Gene	Forward	Reverse
ODF2 (CV864529.1)	AGGCAGGTGGAACAAACCAA	GTTGGTGCTCTCTGACTGCT
ZPBP2 (CV870104.1)	GCGGTTTGGTCAGCAATGAG	TGTCCCGGCTTGCCATAAT
AKAP3 (NM001195324.1)	GCCGCCTCAGAGCTCAATGT	TCATAGCGCAGCACC GACTG
AKAP4 (XM_005657807.3)	CCAGTGCTGAGAAAGTCGGT	TGTCCTGGCATTGGTCTTCC
GAPDH (NM_001206359.1)	GTCGGAGTGAACGGATTTGGC	CACCCCATTTGATGTTGGCG

ODF2, outer dense fiber of sperm tails proteins 2; ZPBP2, zona pellucida binding protein 2; AKAP, A-kinase anchor protein; GAPDH, glyceraldehyde-3-phosphate dehydrogenase.




Fig. 1. Comparison of alkaline phosphatase (ALP) activity between fresh and capacitated boar spermatozoa (A). Spermatozoa were incubated in Beltsville thawing solution (BTS) in the absence (W/O) or presence of varying concentrations of ALP for 2 h. The motility (B), viability (C), and intact acrosome (D) of the spermatozoa were examined after incubation. Values are expressed as mean \pm SEM. The superscript denotes significance at $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.




Fig. 2. Western blotting of boar sperm incubated with varying concentrations of alkaline phosphatase (ALP) or without ALP (W/O). Beta-tubulin antibody was used as a control (A). Relative protein density was measured in each treatment band (B). Values are expressed as mean \pm SEM. The superscript denotes significance at $*p < 0.05$, $**p < 0.01$.

ALP antibody (Fig. 3). We observed from the images obtained that this enzyme was present in the acrosome region, equatorial segment, and tail (a-a"; Fig. 3A). To monitor the changes in ALP activity during the period of sperm storage, we performed immunofluorescence staining on the samples on storage days 1–7 (Fig. 3A). Our analysis indicated a significant decrease in immunofluorescence intensity over the storage period, implying a notable reduction in ALP activity during this period ($p < 0.05, p < 0.01$; Figs. 3A [b-b", c-c", and d-d"] and 3B).

The ALP activity, motility, and viability of sperm stored in BTS without pH balance (a control) and sperm stored in BTS with pH balance (maintained at pH 7.3–7.4 during storage) were compared (Fig. 4) to evaluate the association between pH balance and ALP activity on the fertilizing competence of spermatozoa during storage. Our findings indicated that the pH of liquid boar semen without pH balance gradually declined from day 3 ($p < 0.01, p < 0.001$; Fig. 4A). The ALP activity decreased during the storage period, and lower ALP activity was indicated in the control without pH balance compared to that of sperm stored in BTS with pH balance (32.1 ± 1.5 – 40.1 ± 1.3 IU/mL ALP in the control *vs.* 35.2 ± 1.2 – 43.3 ± 1.1 in the pH balance group, $p < 0.05$; Fig. 4B). In a similar pattern, sperm motility significantly lower in the control compared to the pH balance group ($64 \pm 2.0\%$ – $75 \pm 2.2\%$ in the control *vs.* $70 \pm 2.2\%$ – $79.5 \pm 2.4\%$ in the pH balance group, $p < 0.05$; Fig. 4C). Also, sperm viability showed the same reduction during storage, but the sperm viability in the pH balance group was significantly higher compared to the control group on days 5–7. ($62.5 \pm 1.5\%$ – $70 \pm 1.0\%$ in the control *vs.* $67.5 \pm 1.8\%$ – $75 \pm 1.7\%$ pH balance group, $p < 0.01$; Fig. 4D).

Boar spermatozoa were stored in BTS with different concentrations of taurine (10–80 mM), and the ALP activity, motility, viability, and intracellular ROS were examined over the storage period. All experimental groups showed a pH of 7.4 on day 1 (Fig. 5A). By day 3, the level of pH was stable in the sperm stored in BTS with 10–40 mM taurine, compared to the control without taurine or 80 and 100 mM taurine (pH 7.3–7.2 in control [W/O] or 80, 100 mM taurine *vs.* pH




Fig. 3. Liquid boar semen was stored for 7 days. Immunofluorescent staining was performed on days 1 (a-a"), 3 (b-b"), 5 (c-c"), and 7 (d-d"), respectively. DNA was stained using 4,6-diamidino-2-phenylindole (DAPI, blue). The alkaline phosphatase (green) was localized in the acrosome, equatorial segments and tail (a white arrow; a), and the fluorescence intensity of alkaline phosphatase gradually decreased with longer storage periods (B). Values are expressed as mean \pm SEM. The superscript denotes significance at $*p < 0.05, **p < 0.01$.




Fig. 4. Liquid boar semen was stored at 17°C for 7 days. During the storage period, the liquid boar semen was adjusted and maintained at pH 7.4 (a pH-balance group vs. a control without pH balance). The pH (A), alkaline phosphatase (ALP) activity (B), motility (C), and viability (D) of the spermatozoa were examined on days 1, 3, 5, and 7. Values are expressed as mean \pm SEM. The superscript denotes significance at * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

7.3–7.4 in 10–40 mM taurine, $p < 0.05$, $p < 0.01$, $p < 0.001$; Fig. 5A). On day 5, the pH level further declined, particularly in the sperm stored without taurine and with 80–100 mM taurine (pH 7.1–7.0 in the control or 80, 100 mM taurine vs. pH 7.2–7.4 in 10–40 mM taurine, $p < 0.05$, $p < 0.001$; Fig. 5A). On day 7, there was a significant overall decrease in the pH across all concentrations. However, the medium containing 10 and 40 mM taurine maintained an alkaline pH status (7.1–7.2) compared to the control without taurine or sperm stored with 100 mM taurine, which showed lower pH levels (6.7–6.6, $p < 0.05$, $p < 0.001$; Fig. 5A).

On the first day of storage, ALP activity was somehow different between the groups (Fig. 5B). However, a significant decrease was observed in the sperm stored without taurine and stored with 100 mM taurine on day 3, while the sperm stored with 40 mM taurine showed significantly higher ALP activity compared to the other groups (3.0–3.1 IU/mL in control and 100 mM taurine vs. 3.5–3.9 IU/mL in 10–40 mM taurine, $p < 0.05$, $p < 0.001$; Fig. 5B). The pattern was similar on day 5 (2.2–2.5 IU/mL in control and 100 mM taurine vs. 2.8–3.1 IU/mL in 10–40 mM taurine, $p < 0.05$, $p < 0.001$; Fig. 5B) and day 7 (1.7–2.1 IU/mL in control and 100 mM taurine vs. 2.2–2.7 IU/mL in 10–40 mM taurine, $p < 0.01$, $p < 0.001$; Fig. 5B) of the storage.

On the third day of the storage, sperm motility was higher in the sperm stored in BTS with the 10–80 mM taurine-included groups compared to the control without taurine or 100 mM (83.4%–86.5% in W/O and 100 mM taurine vs. 86.1%–87.7% in 10–80 mM taurine), while the highest motility was observed in the 40 mM group (92.2%, $p < 0.05$, $p < 0.001$; Fig. 5C). Similarly, on day 5, motility further declined, particularly in the control and at the highest taurine concentrations (60.5%–72.0% in the W/O and 100 mM taurine vs. 76.7%–77.1% in 10–80 mM taurine, $p < 0.05$, $p < 0.001$; Fig. 5C), while by day 7, there was a significant overall decrease across all concentrations, and sperm stored in BTS with 10–40 mM taurine maintained better motility compared to the control without taurine or sperm stored in the presence of 80 mM and 100 mM taurine (50.2% in W/O, 45.5%–49.9% in 80–100 mM taurine vs. 51.1%–54.9% in 10–40 mM taurine, $p < 0.05$, $p < 0.001$; Fig. 5C).

The viability trends were also similar, starting around 80%–85% on day 1 across all taurine




Fig. 5. Liquid boar semen was stored in Beltsville thawing solution (BTS) in the absence (W/O) or presence of varying concentrations of taurine for 7 days. The pH level (A), alkaline phosphatase (ALP) activity (B), motility (C), viability (D), and the production of reactive oxygen species (ROS) were examined on days 1, 3, 5, and 7, respectively. Values are expressed as mean \pm SEM. The superscript denotes significance at * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

concentrations. By day 3, the percentages of viable cells were higher in the 10–80 mM taurine-included groups compared to the control without taurine or 100 mM taurine, (65.4%–69.5% in W/O and 100 mM taurine vs. 69.5%–72.5% in 10–80 mM taurine, while the highest viability was observed at 40 mM taurine group (76.5%, $p < 0.05$, $p < 0.001$; Fig. 5D). Similarly, on day 5, viability further declined, particularly in the control and the highest taurine concentrations (56.6% in W/O vs. 58.4%–63.3% in 10–80 mM taurine, $p < 0.05$, $p < 0.001$; Fig. 5D). By day 7, a significant overall decrease was observed across all concentrations. However, sperm stored in BTS with 10 mM and 80 mM taurine maintained significantly higher viability compared to the control without taurine or sperm stored in the presence of 100 mM taurine (36.7%–46.0% in the W/O and 100 mM taurine vs. 39.5%–44.8% in 10–80 mM taurine, $p < 0.05$, $p < 0.001$; Fig. 5D).

The levels of intracellular ROS production were similar across all groups on day 1 (Fig. 5E). Significantly lower fluorescence intensities of ROS were detected in sperm stored in the presence of 10–80 mM concentration of taurine compared to the control or other treatments on day 3 ($p < 0.05$, $p < 0.01$; Fig. 5E). Similarly, ROS production was effectively controlled in the diluents with 10–80 mM taurine on days 5 and 7 ($p < 0.05$, $p < 0.01$, $p < 0.001$; Fig. 5E).

The overall results showed that liquid boar semen containing 10–40 mM taurine exhibited high ALP activity, motility, and viability of spermatozoa as well as low ROS production during the storage period. Therefore, to evaluate the fertilizing capacity of the spermatozoa, the relative mRNA expressions of *ODF2*, *ZPBP2*, *AKAP3*, and *AKAP4* were examined in the control without taurine and 40 mM taurine-supplemented samples during the storage period (Fig. 6). *ODF2* expression

was significantly lower in the control group compared to the 40 mM taurine group on days 5 and 7 ($p < 0.05$; Fig. 6A). The expression of ZPBP2 was similar between the experimental groups (Fig. 6B). On day 5 and 7, expression of AKAP3 was significantly downregulated in the control samples compared with the 40 mM taurine group ($p < 0.05$; Fig. 6C). The AKAP4 expression showed downregulation during storage, but significantly lower expressions were seen in the control compared to sperm stored with 40 mM taurine on days 3 and 5 ($p < 0.05$; Fig. 6D).

DISCUSSION

Liquid preservation is a technique used to store boar semen, where diluted semen is kept at 15°C–20°C for several days before it is used for AI [26]. The storage duration varies according to the composition of the extender used, 2 to 3 days in short-term extenders and five or more days in long-term extenders [27]. Usually, boar spermatozoa experience several changes including reduced motility, viability, permeability of the membrane, and DNA damage during the storage period [28]. ALP is an important enzyme present in seminal plasma [29], and the activity of ALP in seminal plasma is commonly used as a marker for evaluating the condition of accessory glands, sperm metabolic functions, and plasma membrane integrity [30].

Previous studies have indicated that ALP activity serves as a marker to detect the capacitated spermatozoa in vitro [31], and identification of true ejaculations in rhinos [32]. In our study, we analyzed the ALP activity in fresh and capacitated spermatozoa and observed a significant reduction in ALP activity in the capacitated spermatozoa compared to the fresh samples. (Fig. 1A). Similar findings were observed by [31] with altered values, which can be attributed to the different incubation media and experimental conditions. They also found evidence that adding 1.2–2.5 IU/mL ALP to the capacitation media decreased the fertilization ability of boar spermatozoa




Fig. 6. Comparison of relative mRNA expression associated with fertility capacity between sperm stored in Beltsville thawing solution (BTS) in the absence (W/O) or presence of 40 mM taurine. The mRNA was extracted from sperm stored on days 1, 3, 5, and 7, respectively, and then subjected to RT-PCR using target primers ODF2 (A), ZPBP2 (B), AKAP3 (C), and AKAP4 (D). Values are expressed as mean \pm SEM. The superscript denotes significance at * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. ODF2, outer dense fiber of sperm tails proteins 2; ZPBP2, zona pellucida binding protein 2; AKAP, A-kinase anchor protein; RT-PCR, real-time polymerase chain reaction.

in a dose-dependent pattern. Similarly, we evidenced that the addition of 0–5 IU/mL ALP in the incubation media reduced the motility, viability, and acrosome integrity in a dose-dependent pattern (Figs. 1B, 1C, and 1D). ALP plays a critical role in spermatozoa by regulating phosphate metabolism, particularly by hydrolyzing phosphate groups from molecules such as ATP and inorganic pyrophosphate, which are essential for energy production and cellular signaling [33]. This enzymatic activity is crucial for maintaining sperm motility and viability, as well as supporting membrane stability and acrosome integrity during capacitation and fertilization. Interestingly, the addition of inorganic pyrophosphatase PPA1 to the culture medium was observed to lower the rates of both fertilized and polyspermic zygotes in boar spermatozoa [34].

Taurine is present in animals in its free form, exhibiting diverse biological effects such as neutralizing free radicals, modulating reproductive functions, enhancing immunity, and improving antioxidant capacity [35,36]. It serves as a vital amino acid peptide antioxidant in the epididymis and reproductive system. Besides its antioxidant properties, it also reduces cell apoptosis and modulates mitochondrial functions [14]. It additionally regulates membrane permeability to positive ions by specifically modulating Ca^{2+} flux across the membrane. This contributes to maintaining the phospholipid membrane integrity, lowers intracellular free radical levels, and enhances the activation of key antioxidant enzymes [37,38]. ALP is a dimeric metalloenzyme. In other words, it consists of two subunits and requires metal ions to function properly [39]. Usually, ALP activity is affected by the presence of specific metal ions, notably magnesium (Mg^{2+}) and zinc (Zn^{2+}), which are critical for its enzymatic function [40]. Interestingly, taurine has been found to activate ALP even in the absence of these two critical metal ions, due to its indirect antioxidant effect by helping to mitigate the harmful impact of ROS by neutralizing cytotoxic aldehydes, which are the final products of peroxidation cascade reactions [41].

In our study, beginning on day 3 of semen storage, we observed that a taurine concentration of 100 mM decreased sperm motility, whereas concentrations ranging from 10 to 80 mM increased motility, with 40 mM showing the highest effect (Fig. 5C). The motility exhibited a trend of initial increase followed by a decline. This could be attributed to high taurine concentrations altering the extender's osmotic pressure [19] and the pH, affecting sperm membrane permeability, which can disrupt the sperm membrane structure, and reduce progressive motility [42]. Additionally, high taurine levels may induce toxicity, damaging sperm and causing excessive activation of antioxidant enzymes and mitochondria, thereby influencing sperm physiology [43]. The inhibitory impact of 100 mM taurine on motility paralleled changes observed in sperm viability and acrosome integrity across treatment groups.

In our study, taurine exhibited a significant impact on both pH stability and ALP activity in boar spermatozoa. Both of these factors are very important for preserving the quality of spermatozoa throughout the storage period. Taurine supplementation, particularly at concentrations of 10–40 mM, helped maintain stable pH levels in the storage medium, which correlated with preserved ALP activity (Figs. 5A and 5B). High concentrations of taurine (80–100 mM) and the absence of taurine notably impacted the pH of the medium through the storage period, resulting in to decrease. This pH reduction was associated with a marked decrease in ALP activity (Fig. 5A), which negatively affected sperm quality. Consequently, these observations indicate that taurine plays a vital role in regulating pH and influencing ALP activity, thus identifying its potential for maintaining sperm quality and fertilization ability during the preservation of liquid semen.

REFERENCES

1. Karunakaran M, Chakurkar EB, Ratnakaran U, Naik PK, Mondal M, Mondal A, et al.

- Characteristics of boar semen preserved at liquid state. *J Appl Anim Res.* 2017;45:217-20. <https://doi.org/10.1080/09712119.2016.1150848>
2. Bryła M, Trzcińska M. Quality and fertilizing capacity of boar spermatozoa during liquid storage in extender supplemented with different antibiotics. *Anim Reprod Sci.* 2015;163:157-63. <https://doi.org/10.1016/j.anireprosci.2015.11.005>
 3. Iyer SK, Daron HH, Aull JL. Purification and properties of alkaline phosphatase from boar seminal plasma. *J Reprod Fertil.* 1988;82:657-64. <https://doi.org/10.1530/jrf.0.0820657>
 4. Turner RMO, McDonnell SM. Alkaline phosphatase in stallion semen: characterization and clinical applications. *Theriogenology.* 2003;60:1-10. [https://doi.org/10.1016/S0093-691X\(02\)00956-1](https://doi.org/10.1016/S0093-691X(02)00956-1)
 5. Einarsson S, Gustafsson B, Settergren I. Alkaline phosphatase activity of epididymal contents in boars with normal or reduced spermatogenesis. *Andrologia.* 1976;8:25-8. <https://doi.org/10.1111/j.1439-0272.1976.tb01640.x>
 6. Soucek DA, Vary JC. Some properties of acid and alkaline phosphates from boar sperm plasma membranes. *Biol Reprod.* 1984;31:687-93. <https://doi.org/10.1095/biolreprod31.4.687>
 7. Pesch S, Bergmann M, Bostedt H. Determination of some enzymes and macro- and microelements in stallion seminal plasma and their correlations to semen quality. *Theriogenology.* 2006;66:307-13. <https://doi.org/10.1016/j.theriogenology.2005.11.015>
 8. Turner RM, Sertich PL. Use of alkaline phosphatase activity as a diagnostic tool in stallions with azoospermia and oligospermia. *Anim Reprod Sci.* 2001;68:315-65.
 9. Marquez B, Suarez SS. Bovine sperm hyperactivation is promoted by alkaline-stimulated Ca²⁺ influx. *Biol Reprod.* 2007;76:660-5. <https://doi.org/10.1095/biolreprod.106.055038>
 10. Nishigaki T, Jos O, González-Cota AL, Romero F, Treviño CL, Darszon A. Intracellular pH in sperm physiology. *Biochem Biophys Res Commun.* 2014;450:1149-58. <https://doi.org/10.1016/j.bbrc.2014.05.100>
 11. Aksu EH. Current semen extenders for bulls. *Vet J Kastamonu Univ.* 2023;2:34-40. <https://doi.org/10.61262/vetjku.1383952>
 12. Miyazaki T, Honda A, Ikegami T, Matsuzaki Y. The role of taurine on skeletal muscle cell differentiation. In: El Idrissi A, L'Amoreaux W, editors. *Taurine 8: volume 2: nutrition and metabolism, protective role, and role in reproduction, development, and differentiation.* Springer; 2013. p. 321-8.
 13. Das J, Roy A, Sil PC. Mechanism of the protective action of taurine in toxin and drug induced organ pathophysiology and diabetic complications: a review. *Food Funct.* 2012;3:1251-64. <https://doi.org/10.1039/C2FO30117B>
 14. Baliou S, Adamaki M, Ioannou P, Pappa A, Panayiotidis MI, Spandidos DA, et al. Protective role of taurine against oxidative stress (review). *Mol Med Rep.* 2021;24:605. <https://doi.org/10.3892/mmr.2021.12242>
 15. Štefkov K, Procházkov J, Pacherník J. Alkaline phosphatase in stem cells. *Stem Cells Int.* 2015;2015:628368. <https://doi.org/10.1155/2015/628368>
 16. Stapleton PP, O'Flaherty L, Redmond HP, Bouchier-Hayes DJ. Review: host defense-a role for the amino acid taurine? *J Parenter Enteral Nutr.* 1998;22:42-8. <https://doi.org/10.1177/014860719802200142>
 17. Cheng CH, Guo ZX, Wang AL. The protective effects of taurine on oxidative stress, cytoplasmic free-Ca²⁺ and apoptosis of pufferfish (*Takifugu obscurus*) under low temperature stress. *Fish Shellfish Immunol.* 2018;77:457-64. <https://doi.org/10.1016/j.fsi.2018.04.022>
 18. Li H, Zhang XG, Fang Q, Liu Q, Du RR, Yang GS, et al. Supplemental effect of different levels of taurine in Modena on boar semen quality during liquid preservation at 17°C. *Anim Sci*

- J. 2017;88:1692-9. <https://doi.org/10.1111/asj.12865>
19. Zhang L, Wang Y, Sohail T, Kang Y, Niu H, Sun X, et al. Effects of taurine on sperm quality during room temperature storage in Hu sheep. *Animals*. 2021;11:2725. <https://doi.org/10.3390/ani11092725>
 20. Bayomy MFF, Hassab El-Nabi SE, El Kassas TA, Attia ZI, Saeed AM, Taha HSA, et al. Extender supplementation with glutathione (GSH) and taurine improves in vitro sperm quality and antioxidant status of New Zealand rabbits during chilled storage for up to 72 hours. *Vet Med Int*. 2023;2023:8339591. <https://doi.org/10.1155/2023/8339591>
 21. Atte PO, Abdulrashid M, Daudu OM, Zahraddeen D. Effects of taurine and season on semen characteristics and sperm morphology of West African Dwarf Rams. *Niger J Anim Sci Technol*. 2022;5:71-9.
 22. Perumal P, Vupru K, Rajkhowa C. Effect of addition of taurine on the liquid storage (5°C) of mithun (*Bos frontalis*) semen. *Vet Med Int*. 2013;2013:165348. <https://doi.org/10.1155/2013/165348>
 23. Pursel VG, Johnson LA. Frozen boar spermatozoa: methods of thawing pellets. *J Anim Sci*. 1976;42:927-31. <https://doi.org/10.2527/jas1976.424927x>
 24. Yi YJ, Sung DY, Millette C, Sutovsky M, Kennedy C, Sutovsky P, et al. Sperm GIRK2-containing K⁺ inward rectifying channels participate in sperm capacitation and fertilization. *Syst Biol Reprod Med*. 2011;57:296-308. <https://doi.org/10.3109/19396368.2011.631685>
 25. Berger T. Pisum sativum agglutinin used as an acrosomal stain of porcine and caprine sperm. *Theriogenology*. 1990;33:689-95. [https://doi.org/10.1016/0093-691X\(90\)90546-6](https://doi.org/10.1016/0093-691X(90)90546-6)
 26. Yeste M. State-of-the-art of boar sperm preservation in liquid and frozen state. *Anim Reprod*. 2017;14:69-81. <https://doi.org/10.21451/1984-3143-AR895>
 27. Karageorgiou MA, Tsousis G, Boscicos CM, Tzika ED, Tassis PD, Tsakmakidis IA. A comparative study of boar semen extenders with different proposed preservation times and their effect on semen quality and fertility. *Acta Vet Brno*. 2016;85:23-31. <https://doi.org/10.2754/avb201685010023>
 28. Trzcińska M, Bryła M, Smoraż Z. Effect of liquid storage on membrane integrity and mitochondrial activity: a new diagnostic method of evaluating boar sperm quality. *J Anim Feed Sci*. 2008;17:372-80. <https://doi.org/10.22358/jafs/66625/2008>
 29. Bucci D, Giaretta E, Spinaci M, Rizzato G, Isani G, Mislei B, et al. Characterization of alkaline phosphatase activity in seminal plasma and in fresh and frozen-thawed stallion spermatozoa. *Theriogenology*. 2016;85:288-95.e2. <https://doi.org/10.1016/j.theriogenology.2015.09.007>
 30. Collodel G, Nerucci F, Signorini C, Iacoponi F, Moretti E. Associations between biochemical components of human semen with seminal conditions. *Syst Biol Reprod Med*. 2019;65:155-63. <https://doi.org/10.1080/19396368.2018.1548668>
 31. Bucci D, Isani G, Giaretta E, Spinaci M, Tamanini C, Ferlizza E, et al. Alkaline phosphatase in boar sperm function. *Andrology*. 2014;2:100-6. <https://doi.org/10.1111/j.2047-2927.2013.00159.x>
 32. Roth TL, Stoops MA, Robeck TR, Ball RL, Wolfe BA, Finnegan MV, et al. Alkaline phosphatase as an indicator of true ejaculation in the rhinoceros. *Theriogenology*. 2010;74:1701-6. <https://doi.org/10.1016/j.theriogenology.2010.05.021>
 33. Terkeltaub RA. Inorganic pyrophosphate generation and disposition in pathophysiology. *Am J Physiol Cell Physiol*. 2001;281:C1-11. <https://doi.org/10.1152/ajpcell.2001.281.1.C1>
 34. Yi YJ, Sutovsky M, Kennedy C, Sutovsky P. Identification of the inorganic pyrophosphate metabolizing, ATP substituting pathway in mammalian spermatozoa. *PLOS ONE*. 2012;7:e34524. <https://doi.org/10.1371/journal.pone.0034524>

35. De la Puerta C, Arrieta FJ, Balsa JA, Botella-Carretero JI, Zamarrón I, Vázquez C. Taurine and glucose metabolism: a review. *Nutr Hosp.* 2010;25:910-9. <https://doi.org/10.3305/nh.2010.25.6.4815>
36. Seidel U, Huebbe P, Rimbach G. Taurine: a regulator of cellular redox homeostasis and skeletal muscle function. *Mol Nutr Food Res.* 2019;63:1800569. <https://doi.org/10.1002/mnfr.201800569>
37. Schaffer SW, Jong CJ, Ramila KC, Azuma J. Physiological roles of taurine in heart and muscle. *J Biomed Sci.* 2010;17:S2. <https://doi.org/10.1186/1423-0127-17-S1-S2>
38. Huxtable RJ. Physiological actions of taurine. *Physiol Rev.* 1992;72:101-63. <https://doi.org/10.1152/physrev.1992.72.1.101>
39. Sharma U, Pal D, Prasad R. Alkaline phosphatase: an overview. *Indian J Clin Biochem.* 2014;29:269-78. <https://doi.org/10.1007/s12291-013-0408-y>
40. Katz AK, Glusker JP. Roles of zinc and magnesium ions in enzymes. *Adv Mol Struct Res.* 1998;4:227-79. [https://doi.org/10.1016/S1087-3295\(98\)80008-X](https://doi.org/10.1016/S1087-3295(98)80008-X)
41. Schaffer S, Azuma J, Takahashi K, Mozaffari M. Why is taurine cytoprotective? In: Lombardini JB, Schaffer SW, Azuma J, editors. *Taurine 5: beginning the 21st century. Advances in experimental medicine and biology.* Springer; 2003. p. 307-21.
42. Khan MIR, Ijaz A. Effects of osmotic pressure on motility, plasma membrane integrity and viability in fresh and frozen-thawed buffalo spermatozoa. *Animal.* 2008;2:548-53. <https://doi.org/10.1017/S1751731108001596>
43. Li Y, Peng Q, Shang J, Dong W, Wu S, Guo X, et al. The role of taurine in male reproduction: physiology, pathology and toxicology. *Front Endocrinol.* 2023;14:1017886. <https://doi.org/10.3389/fendo.2023.1017886>