

Effect of microencapsulated organic acids-essential oils blend and protease on performance and gut health of broilers under nutritional challenges


Pattaveekan Preesong¹, Preeda Lertwatcharasarakul²,
Koonphol Pongmanee¹, Akaradet Seemacharoensri³,
Glenmer Bathan Tactacan³, Chanporn Chaosap⁴, Yuwares Ruangpanit^{1*}

¹Department of Animal Science, Faculty of Agriculture at Kamphaeng Saen, Kasetsart University, Kamphaeng Saen Campus, Nakhon Pathom 73140, Thailand

²Department of Pathology, Faculty of Veterinary Medicine, Kasetsart University, Bangkok 10900, Thailand

³Jefo Nutrition Inc., Saint-Hyacinthe, Quebec J2S 7B6, Canada

⁴Department of Agricultural Education, School of Industrial Education and Technology, King Mongkut's Institute of Technology Ladkrabang, Bangkok 10520, Thailand



Received: Apr 4, 2024

Revised: Aug 29, 2024

Accepted: Nov 28, 2024

*Corresponding author

Yuwares Ruangpanit
Department of Animal Science,
Faculty of Agriculture at Kamphaeng
Saen, Kasetsart University, Kamphaeng
Saen Campus, Nakhon Pathom 73140,
Thailand.

Tel: +66-34351892

E-mail: yowares.s@ku.th

Copyright © 2025 Korean Society of
Animal Science and Technology.
This is an Open Access article
distributed under the terms of the
Creative Commons Attribution
Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted
non-commercial use, distribution, and
reproduction in any medium, provided
the original work is properly cited.

ORCID

Pattaveekan Preesong
<https://orcid.org/0009-0007-8937-7320>
Preeda Lertwatcharasarakul
<https://orcid.org/0000-0002-4023-4341>

Abstract

This study examined the effects of microencapsulated organic acids and essential oils (EOA) combined with a protease (PRO) supplement on the growth performance and gut health of broilers subjected to nutritional challenges through a diet high in wheat and corn distillers dried grains with solubles (DDGS). The treatments were: 1) corn and soybean meal-based diet with high levels of wheat and corn DDGS (WD); 2) WD+microencapsulated organic acids and essential oils at 300 mg/kg (EOA); 3) WD+PRO at 125 mg/kg (PRO); and 4) WD+EOA at 300 mg/kg+PRO at 125 mg/kg (EOA+PRO). Body weight gain, feed intake and mortality rate did not differ among treatments ($p > 0.05$). However, feed conversion ratio from day 1-35 was lower in the EOA+PRO group than in the WD group ($p < 0.05$). The EOA+PRO group had a lower jejunal crypt depth (CD) and a higher villus height/crypt depth (VH/CD) ratio than the other groups ($p < 0.01$). The putrescine level was higher in the WD group than in the other groups ($p < 0.05$). On day 35, the EOA and EOA+PRO groups had higher claudin-1 mRNA expression than the WD and PRO groups ($p < 0.01$). Occludin mRNA expression was higher in the EOA and PRO groups than in the WD group ($p < 0.01$). In summary, the combination of EOA and PRO improved feed efficiency and gut health in broilers fed a high wheat and corn DDGS diet. This was demonstrated by decreased jejunal CD, increased VH/CD ratio, increased mRNA expression of claudin-1 at the tight junction and decreased putrescine content in the hindgut, suggesting an indirect effect on pathogenic bacteria.

Keywords: Fumaric acid, Thymol, Alkaline serine endopeptidase, Tight junction proteins, Amine

Koonphol Pongmanee
<https://orcid.org/0000-0003-0827-0702>
 Akaradet Seemacharoensri
<https://orcid.org/0000-0003-3835-8975>
 Glenmer Bathan Tactacan
<https://orcid.org/0000-0003-3320-3560>
 Chanporn Chaosap
<https://orcid.org/0000-0002-3816-1884>
 Yuwares Ruangpanit
<https://orcid.org/0000-0003-0258-6300>

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This research was funded by Jefe Nutrition Inc., Saint-Hyacinthe, Quebec, Canada (The grant number is KST_Br_1904).

Acknowledgements

The animal experiment and laboratory facilities were supported by the Department of Animal Science, Faculty of Agriculture at Kamphaeng Saen, the Kamphaeng Saen Veterinary Diagnosis Center, Faculty of Veterinary Medicine, and Phytochemistry Research and Analysis Unit, Central Laboratory and Greenhouse Complex, Kasetsart University, Kamphaeng Saen Campus, Nakhon Pathom, 73140, Thailand.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Pongmanee K, Seemacharoensri A, Tactacan GB, Ruangpanit Y.
 Data curation: Ruangpanit Y.
 Formal analysis: Preesong P, Ruangpanit Y.
 Methodology: Preesong P, Lertwacharasarakul P.
 Software: Ruangpanit Y.
 Validation: Chaosap C, Ruangpanit Y.
 Investigation: Preesong P, Ruangpanit Y.
 Writing - original draft: Preesong P, Chaosap C.
 Writing - review & editing: Preesong P, Lertwacharasarakul P, Pongmanee K, Seemacharoensri A, Tactacan GB, Chaosap C, Ruangpanit Y.

Ethics approval and consent to participate

The experimental protocol was approved by the Animal Care and Use Committee of Kasetsart University (protocol number: ACKU62-AQK-012).

INTRODUCTION

Antibiotic growth promoters (AGPs) have traditionally been used in the poultry industry to improve growth, feed efficiency, and gut physiology [1]. However, increasing concerns about antibiotic-resistant microorganisms have led to global efforts to reduce or ban the use of AGPs in livestock production [2]. As a result, regulatory authorities have introduced restrictions and guidelines to promote responsible antibiotic use, which has led to the exploration of alternative strategies to improve poultry health and performance.

To address these challenges, various feed additives are being explored as alternatives to AGPs. Probiotics, prebiotics, organic acids and essential oils have shown promise. Organic acids improve broiler health by supporting immunological function, pancreatic enzyme activity and gut microbiota balance [3–6]. Essential oils containing compounds such as thymol, carvacrol and eugenol provide benefits such as improved immune function and a reduction in pathogenic bacteria [7–9]. The combination of essential oils with organic acids (EOA) can further improve gut health and performance compared to single supplements [10–12].

Alternative feed ingredients such as wheat and corn distillers dried grains with solubles (DDGS) are commonly used in poultry feed. Corn DDGS, a by-product of ethanol production, provides protein and energy, but may have lower protein quality due to high levels of non-starch polysaccharides (NSP) and lower amino acid digestibility [13]. High NSP content in DDGS may promote colonization with *Clostridium perfringens*, especially under conditions of necrotic enteritis (NE) [14]. Similarly, wheat contains arabinoxylan, an NSP that increases gut viscosity, leading to reduced nutrient absorption and microbial imbalances, e.g. with *Escherichia coli* and *Salmonella* spp. [15–19]. In addition, poor protein digestibility associated with high NSP content can lead to microbial fermentation of nitrogen metabolites, which impairs the intestinal barrier, increases tight junction (TJ) permeability and impairs broiler growth [20,21].

Protease (PRO) enzymes contribute significantly to minimizing undigested proteins, maximizing amino acid availability, reducing dietary protein requirements, supporting weight gain and feed efficiency, reducing proteolytic fermentation, reducing biogenic amines, and improving gut integrity [22–24]. Consequently, there is considerable interest in the market to utilize undigested proteins through the use of exogenous enzymes such as PRO. This approach facilitates the formulation of balanced diets with reduced protein levels, which ultimately leads to cost savings in feed production [25,26].

Microencapsulation is an important technique to deliver bioactive compounds into the gastrointestinal tract [27,28]. It ensures the stability and targeted release of these compounds in the hindgut, where pathogenic bacteria are most prevalent [29]. Without microencapsulation, organic acids may dissociate in the upper gastrointestinal tract and essential oils may be absorbed before they reach the hindgut, reducing their efficacy. Microencapsulated organic acids and essential oils show significantly increased bactericidal and bacteriostatic activity compared to unprotected forms [27,28,30].

The hypothesis of this study is that microencapsulated EOA in combination with PRO can improve the growth and gut health of broilers fed a high wheat and corn DDGS diet, which serves as a nutritional model challenging avian gut resilience. The broiler chickens in the current study were raised under AGP-free programs. Chemical coccidiostats, which are not classified as veterinary medicinal products and can be used as feed additives according to the EU regulation [31], were used to control coccidiosis. The microencapsulated organic acids are fumaric, malic, sorbic and citric acids, and the essential oils are vanillin, eugenol and thymol, all encapsulated in hydrogenated vegetable fat. The PRO used is an alkaline serine endopeptidase derived from the fermentation of

Streptomyces. The aim is to investigate the effects of this microencapsulated EOA in combination with PRO on the growth and gut health of broiler chickens fed a high wheat-corn DDGS diet. Gut health was assessed by analyzing gut morphology, microbial metabolites in the cecum and mRNA expression of TJ proteins, which are critical for maintaining intestinal integrity.

MATERIALS AND METHODS

Bird Husbandry and experimental design

The experimental protocol was approved by the Animal Care and Use Committee of Kasetsart University (protocol number: ACKU62-AQK-012). A total of 1,400 male Ross 308 broiler chicks (Panus Pokphand) were reared in 56 pens (1.5 × 2.0 m). All birds were randomly assigned to 4 treatments using a completely randomized design. There were 14 replicates with 25 birds per replicate in each treatment. The dietary treatments were: 1) corn and soybean meal-based diet with high levels of wheat and corn DDGS (WD); 2) WD+microencapsulated organic acids and essential oils at 300 mg/kg (EOA); 3) WD+PRO at 125 mg/kg (PRO); and 4) WD+EOA at 300 mg/kg+PRO at 125 mg/kg (EOA+PRO). The EOA contained a combination of fumaric acid, sorbic acid, malic acid and citric acid with vanillin, eugenol, and thymol microencapsulated in hydrogenated vegetable fat. The PRO enzyme was an alkaline serine endopeptidase with PRO activity of 1.10 U/g. Both are commercially available products provided by Jefe Nutrition. During the trial, the birds had unlimited access to water and feed. The ambient temperature was 32°C for the first three days, then steadily dropped to 25°C on day 14. The light settings were 23 hours of light and 1 hour of darkness during the experiment.

Experimental diets

The main ingredients of the WD group were corn and soybean meal. In the starter, grower, and finisher diets, 20%, 25%, and 30% wheat replaced corn as the energy source, and 10%, 12.5%, and 15% corn DDGS replaced soybean meal as the protein source. All experimental diets were formulated following the strain recommendations [32]. The diets were mixed with a horizontal mixer and pelleted at 80°C according to the manufacturer's instructions (Bangkok Animal Research Center). All experimental diets were analyzed for crude protein, ether extract, crude fiber, gross energy, calcium and phosphorus according to AOAC guidelines [33]. The details of the diet composition are listed in Table 1.

Data recording

The body weight of all birds and the feed intake per pen were recorded on days 1, 7, 14, 28, and 35. Feed intake (FI), feed conversion ratio (FCR), and body weight gain (BWG) were calculated for each bird and each replicate. Mortality was recorded daily, and the weight of dead birds was recorded to calculate the adjusted FCR.

Sampling

On days 14 and 35, one bird was randomly selected from each replicate (a total of 14 birds per treatment), its body weight (BW) was measured and it was then humanely sacrificed by stunning and bleeding. The mid jejunum was removed for intestinal morphological examination. The intestinal mucosa was scraped with a sterile glass slide. Intestinal mucosa samples were immediately frozen in liquid nitrogen and stored at -80°C for subsequent mRNA expression analysis of TJ proteins. Cecal content samples from 35-day-old birds were collected and stored in a freezer at -20°C to analyze ammonia, biogenic amines and volatile fatty acids (VFA) in the ceca.

Table 1. Ingredients, calculated and analyzed nutrient composition of experimental diets¹⁾

| Ingredient (%) | Starter diet (d 0–10) | Grower diet (d 11–24) | Finisher diet (d 25–35) |
|--------------------------------------|--------------------------|--------------------------|----------------------------|
| Corn (7.9% CP) | 30.30 | 26.85 | 25.40 |
| Wheat (13% CP) | 20.00 | 25.00 | 30.00 |
| Soybean meal (48.5% CP) | 29.60 | 24.72 | 17.87 |
| Corn DDGS (27% CP) ²⁾ | 10.00 | 12.50 | 15.00 |
| Soybean oil | 4.44 | 5.83 | 6.85 |
| Mono-dicalcium phosphate | 1.74 | 1.50 | 1.25 |
| Calcium carbonate | 1.29 | 1.19 | 1.13 |
| Pellet binder ³⁾ | 0.30 | 0.30 | 0.30 |
| Salt | 0.07 | 0.08 | 0.06 |
| Broiler vit/min premix ⁴⁾ | 0.20 | 0.20 | 0.20 |
| DL-Methionine | 0.34 | 0.29 | 0.27 |
| L-Lysine HCl | 0.44 | 0.42 | 0.47 |
| L-Threonine | 0.19 | 0.16 | 0.16 |
| Sodium bicarbonate | 0.36 | 0.32 | 0.33 |
| Choline chloride (60%) | 0.38 | 0.36 | 0.35 |
| Coccidiostat (Cygro) | 0.05 | 0.05 | 0.05 |
| L-Isoleucine | 0.10 | 0.07 | 0.09 |
| L-Arginine (98%) | 0.14 | 0.13 | 0.18 |
| L-Valine | 0.06 | 0.03 | 0.04 |
| ME for poultry (Kcal/kg) | 3,000 | 3,100 | 3,200 |
| Crude protein (%) | 23.0 | 21.5 | 19.5 |
| Crude fat (%) | 7.17 | 8.61 | 9.74 |
| Crude fiber (%) | 3.48 | 3.14 | 3.03 |
| Digestible ⁶⁾ lysine (%) | 1.47 | 1.34 | 1.22 |
| Digestible methionine (%) | 0.68 | 0.62 | 0.57 |
| Digestible methionine+cysteine (%) | 1.07 | 0.99 | 0.91 |
| Digestible tryptophan (%) | 0.23 | 0.21 | 0.18 |
| Digestible isoleucine (%) | 0.86 | 0.78 | 0.71 |
| Digestible threonine (%) | 0.86 | 0.77 | 0.69 |
| Digestible valine (%) | 0.96 | 0.87 | 0.78 |
| Digestible arginine (%) | 1.37 | 1.23 | 1.10 |
| Calcium (%) | 0.96 | 0.87 | 0.79 |
| Total phosphorus (%) | 0.81 | 0.75 | 0.68 |
| Available phosphorus (%) | 0.48 | 0.44 | 0.40 |
| Choline (mg/kg) | 1,700 | 1,600 | 1,550 |
| Sodium (%) | 0.16 | 0.16 | 0.16 |
| Analyzed nutrient | | | |
| GE (Kcal/kg) | 4,675 | 4,700 | 4,798 |
| Crude protein (%) | 21.3 | 19.7 | 17.7 |
| Crude fat (%) | 2.7 | 2.9 | 2.8 |
| Crude fiber (%) | 7.3 | 8.7 | 9.4 |
| Ash (%) | 5.8 | 5.3 | 4.8 |
| Calcium (%) | 1.0 | 1.0 | 0.9 |
| Phosphorus (%) | 0.8 | 0.8 | 0.7 |

¹⁾Experimental diet: 1) corn-soybean meal basal diet with wheat and corn distillers dried grains with solubles (WD); 2) WD+microencapsulated organic acids-essential oils blend at 300 mg/kg (EOA); 3) WD+PRO at 125 mg/kg (PRO); 4) WD+microencapsulated organic acids-essential oils blend at 300 mg/kg+PRO at 125 mg/kg (EOA+PRO).

²⁾Corn distillers dried grains with solubles.

³⁾Pellet binder from Pelex Dry, Bentoli.

⁴⁾Broiler vit/min premix provided per kilograms of diet: vitamin A (all-trans retinol) 1,200 IU; vitamin D₃ (cholecalciferol) 2,400 IU; vitamin E (l- α -tocopherol) 60 mg; vitamin K 240 mg; vitamin B₁ 300 mg; vitamin B₂ 800 mg; vitamin B₆ 400 mg; vitamin B₁₂ 2 mg; niacin 5,000 mg; pantothenic acid 1,500 mg; biotin 40 mg; folic 200 mg; Cu (copper sulfate) 1,500 mg; Fe (ferrous sulfate) 4,000 mg; Mn (manganese sulfate) 10,000 mg; Zn (zinc sulfate) 10,000 mg; I (iodide) 100 mg; Se (Selenate) 100 mg.

⁵⁾Coccidiostat from Cygro, Zoetis.

⁶⁾Apparent ileal digestible amino acids.

CP, crude protein; DDGS, distillers dried grains with solubles; HCl, hydrochloric acid; ME, metabolizable energy; GE, gross energy; WD, wheat and corn DDGS; EOA, essential oils with organic acids; PRO, protease; IU, international unit.

Gene expression of intestinal barrier tight junction proteins

RNA isolation and cDNA synthesis

After extraction from frozen jejunum mucosal samples using the GenUP™ total RNA kit (Biotechrabbit GmbH), RNA quantity and quality were determined using a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific) at 260 and 280 nm. Subsequently, 1 µg of RNA was used to synthesize the first strand of cDNA using a cDNA synthesis kit (Biotechrabbit GmbH), and the resulting cDNA was stored at −20 °C for subsequent analysis.

Real-time PCR

Expression of the claudin-1, Zonula Occludens-1 (ZO-1) and occludin genes was determined by real time PCR using the specific primers listed in Table 2 [34,35]. Rigorous testing ensured primer efficiency and linearity. Each reaction was performed in triplicate for each gene and sample. Gene expression was normalized using glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and TATA-binding protein (TBP) as reference genes, according to the methodology described by Taylor et al. [36].

Intestinal morphology

Intestinal morphology examinations were performed according to Iji et al. [37]. A 1 cm sample of the jejunum (between the terminal loop of the duodenum and Meckel's diverticulum) was excised and immediately fixed in 10% formalin. The fixed samples were dehydrated in ethanol, cleared in xylene, and embedded in paraffin. Two sections, each 7 µm thick, were mounted on microscope slides and stained with alcian blue, hematoxylin, and eosin. The stained sections were examined under a light microscope at 40x magnification using an Olympus CX33 microscope equipped with an Olympus DP22 digital camera and DP2-SAL imaging software (Olympus). Villus height (VH), measured from the base transition zone between villus and crypt to the apex, crypt depth (CD), measured from the base of the villi to the bottom of the glands, and villus width (VW), measured from the left villus crypt junction to the right of the villus crypt junction, were quantified. VH/CD ratio was determined by measuring 9 randomly selected villi and their corresponding crypts.

Microbial metabolites in the cecal content

Volatile fatty acid analysis

VFA were analyzed by gas chromatography according to Thanh et al. [38]. In brief, 200 mg of

Table 2. Nucleotide sequences of primers for quantitative real-time PCR assay

| Gene | Primer sequences | GenBank accession number |
|-----------|--------------------------------|--------------------------|
| Claudin-1 | FP 5'-AAGGTGTACGACTCGCTGCT-3' | NM_001013611.2 |
| | RP 5'-CAGCAACAAACACCAACC-3' | |
| ZO-1 | FP 5'-AAGTGGGAAGAATGCCAAA-3' | XM_015278975.2 |
| | RP 5'-GGTCCTGGATCCCGTATCT-3' | |
| Occludin | FP 5'-ACGGCAAAGCCAACATCTAC-3' | NM_205128.1 |
| | RP 5'-ATCCGCCACGTTCTTCAC-3' | |
| GAPDH | FP 5'-CAACCCCAATGTCTCTGTT-3' | NM_204305.1 |
| | RP 5'-TCAGCAGCAGCCTTCACTAC-3' | |
| TBP | FP 5'-GTCCACGGTGAATCTTGTT-3' | NM_205103.1 |
| | RP 5'-GCGCAGTAGTACGTGGTTCTC-3' | |

Source of primer: ZO-1, occludin, GAPDH, and TBP [34], claudin-1 [35].

PCR, polymerase chain reaction; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; ZO-1, zonula occludens 1; TBP, TATA-binding protein. FP, forward primer; RP, reverse primer.

cecal content was mixed with distilled water in a 1:1 ratio (w/v) and centrifuged at 16,300×g at 4°C for 20 min. Then, 100 µL of the supernatant was transferred and mixed with 100 µL of 24% metaphosphoric acid in 1.5 M sulfuric acid, stirred for 5 min, and allowed to stand overnight at 4°C. The mixture was then centrifuged at 8,944×g for 5 min at 4°C. The supernatant was mixed with an equal volume of 3 mM crotonic acid and used as an internal standard. Subsequently, 1 µL of the prepared sample was injected and separated by gas chromatography using a CP-Wax 52 CB (50 m × 0.32 mm) column (Agilent Technologies). Helium (2 mL/min) was used as the mobile phase, and the injector and detector temperatures were 250°C and 280°C, respectively. The column temperature was set to 200°C. External standards with 3 mM acetic acid, propionic acid, butyric acid, and 1.5 mM crotonic acid were used to identify the peaks.

Ammonia analysis

The frozen cecal content was analyzed according to Meyer et al. [39]. In brief, 500 mL of 100 mM 3-(N-morpholino) propanesulfonic acid was added to 250 mg of cecal content. The sample was centrifuged at 4°C and 12,888×g for 20 min. Then, 250 µL of the supernatant was mixed with 25 µL of Carrez Clarification Reagent Kit (Sigma-Aldrich) and centrifuged at 4°C and 12,888×g for 10 min. Ammonia was analyzed according to the method described by Weatherburn [40].

Amine analysis

Amine analysis of the extracted cecal contents was performed as described by Saarinen [41]. A 500 µL aliquot of 0.4 M perchloric acid was used to deprotonate 250 mg of the frozen sample. The derivatization reaction of the amine in the extracted sample was carried out with dansyl chloride as described by Eerola et al. [42]. The derivative solution was filtered using a nylon membrane filter with a pore size of 0.22 µm. Subsequently, 10 µL of the sample was injected into an ODS2 column (4.0 × 250 m; Waters) using a 717 plus autosampler at 40°C. Peaks were detected at 254 nm using a 2998 Photodiode Array Detector (Waters) and analyzed using Empower Software Build 2154 (Waters). HPLC-grade water was used as mobile phase A and HPLC-grade acetonitrile (Fisher Scientific) was used as mobile phase B. The gradient elution was initially 50%, after 25 min 10%, after 35 min 50%, after 40 min 50% at a flow rate of 1 mL/min. Finally, 1-aminoheptane was used as an internal standard.

Putrescine dihydrochloride and cadaverine dihydrochloride were used as external standards and diluted in water to prepare the stock solution. Subsequently, the external standards were diluted with 0.4 M perchloric acid for serial dilution.

Statistical analysis

Percentage mortality data were obtained by square root transformation of $Y + 0.5$ ($Y = \% \text{mortality}$). Relative gene expression was Log-transformed ($\text{Log}_2 \Delta\Delta\text{Cq}$) prior to statistical analysis. All data were tested for normality using the Kolmogorov-Smirnov test before performing statistical analyses. Statistical differences between treatments were analyzed using the GLM procedure from SAS Studio University Edition (SAS Institute). Differences among treatments were determined using Tukey's test for honestly significant differences. Significant values were determined based on a p -value ≤ 0.05 , and trends were reported at $0.05 < p \leq 0.1$.

RESULTS

This experiment was conducted to investigate the effects of EOA in combination with PRO on the growth and gut health of broilers raised without AGPs. It is crucial to challenge intestinal

homeostasis, as in the absence of such challenges, gut-acting growth promoters may have limited effects on performance [43,44]. Therefore, this study employed a nutritional model that challenged avian gut resilience using a diet high in wheat and corn DDGS. The EOA blend was supplemented in a microencapsulated form to ensure the stability and targeted release of these compounds in the hindgut, where pathogenic bacteria are most prevalent. Additionally, PRO was included to assess its potential in improving nutrient utilization, particularly in overcoming the poor digestibility associated with the high NSP content in corn DDGS and wheat. The results of this study are presented below.

Growth performance

In the current study, no effects of the dietary treatments ($p > 0.05$) were observed on BWG, FI and mortality rate (Table 3). On day 8–14, the PRO group had a higher FCR than the EOA+PRO group ($p < 0.01$), while the FCR of the WD and EOA groups did not differ from the others ($p > 0.05$). On day 1–35, the WD group had a higher FCR than the EOA+PRO group ($p < 0.05$), while

Table 3. Growth performance (mean¹⁾) of broiler chickens fed a high wheat and corn DDGS diet²⁾ supplemented with microencapsulated organic acids-essential oils blend and protease

| Trait | WD | EOA | PRO | EOA + PRO | Pooled SEM ³⁾ | p-value |
|-------------------------------|---------------------|---------------------|---------------------|--------------------|--------------------------|---------|
| Body weight gain (g/bird) | | | | | | |
| d 1–7 | 174 | 172 | 174 | 172 | 2.12 | 0.236 |
| d 8–14 | 343 | 346 | 344 | 346 | 2.90 | 0.677 |
| d 15–28 | 1,186 | 1,187 | 1,198 | 1,203 | 6.43 | 0.675 |
| d 29–35 | 585 | 612 | 590 | 595 | 7.21 | 0.563 |
| d 1–35 | 2,286 | 2,303 | 2,308 | 2,317 | 8.59 | 0.721 |
| Feed intake (g/bird) | | | | | | |
| d 1–7 | 178 | 177 | 176 | 175 | 1.68 | 0.152 |
| d 8–14 | 428 | 430 | 432 | 431 | 2.62 | 0.550 |
| d 15–28 | 1,769 | 1,778 | 1,765 | 1,784 | 5.74 | 0.399 |
| d 29–35 | 1,091 | 1,129 | 1,116 | 1,092 | 7.63 | 0.242 |
| d 1–35 | 3,474 | 3,487 | 3,500 | 3,481 | 9.07 | 0.868 |
| Feed conversion ratio (kg/kg) | | | | | | |
| d 1–7 | 1.022 | 1.026 | 1.017 | 1.022 | 0.142 | 0.766 |
| d 8–14 | 1.254 ^{ab} | 1.238 ^{ab} | 1.257 ^a | 1.235 ^b | 0.1397 | 0.007 |
| d 15–28 | 1.487 | 1.496 | 1.483 | 1.492 | 0.1797 | 0.697 |
| d 29–35 | 1.824 | 1.853 | 1.896 | 1.842 | 0.3306 | 0.366 |
| d 1–35 | 1.529 ^a | 1.515 ^{ab} | 1.517 ^{ab} | 1.499 ^b | 0.1643 | 0.037 |
| Mortality (%) | | | | | | |
| d 1–7 | 0.00 | 0.00 | 0.00 | 0.00 | 0.000 | - |
| d 8–14 | 0.29 | 0.00 | 0.29 | 0.30 | 0.572 | 0.801 |
| d 15–28 | 0.00 | 0.30 | 0.31 | 0.00 | 0.5169 | 0.576 |
| d 29–35 | 0.00 | 0.00 | 0.00 | 0.61 | 0.5016 | 0.103 |
| d 1–35 | 0.29 | 0.29 | 0.57 | 0.86 | 0.7065 | 0.676 |

¹⁾Each value represents the mean of 14 replicates.

²⁾Dietary treatments: WD = corn-soybean meal basal diet with wheat and corn distillers dried grains with solubles; EOA = WD + microencapsulated organic acids-essential oils blend at 300 mg/kg; PRO = WD + PRO at 125 mg/kg; and EOA + PRO = WD + microencapsulated organic acids-essential oils blend at 300 mg/kg + PRO at 125 mg/kg.

³⁾Pooled SEM (n = 56).

^{a–c}Within a row, means with different superscripts differ significantly ($p < 0.05$).

DDGS, distillers dried grains with solubles; WD, wheat and corn DDGS; EOA, essential oils with organic acids; PRO, protease.

the FCR of the EOA and PRO groups did not differ significantly from the other groups ($p > 0.05$).

Expression of intestinal barrier tight junction proteins

Figs. 1 and 2 show the effects of the dietary treatments on the mRNA expression of selected intestinal barrier TJ proteins in the jejunum mucosa on days 14 and 35. On day 14, the mRNA




Fig. 1. Zonula occludens-1 (ZO-1) (A), Claudin (B) and Occludin (C) mRNA relative expression in jejunum mucosa of broiler chickens fed a high wheat and corn DDGS diet supplemented with microencapsulated organic acids-essential oils blend and protease at 14 days of age. Each value represents the mean of 14 replicates \pm SEM ($n = 14$), and different letters denote significant p -values > 0.05 and < 0.10 . Dietary treatments: WD = corn-soybean meal basal diet with wheat and corn distillers dried grains with solubles; EOA = WD + microencapsulated organic acids-essential oils blend at 300 mg/kg; PRO = WD + PRO at 125 mg/kg; and EOA + PRO = WD + microencapsulated organic acids-essential oils blend at 300 mg/kg + PRO at 125 mg/kg. DDGS, distillers dried grains with solubles; WD, wheat and corn DDGS; EOA, essential oils with organic acids; PRO, protease.




Fig. 2. Zonula occludens-1 (ZO-1) (A), Claudin (B) and Occludin (C) mRNA relative expression in jejunum mucosa of broiler chickens fed a high wheat and corn DDGS diet supplemented with microencapsulated organic acids-essential oils blend and protease at 35 days of age. Each value represents the mean of 14 replicates \pm SEM (n = 14), and different letters denote significant difference ($p < 0.05$). Dietary treatments: WD = corn-soybean meal basal diet with wheat and corn distillers dried grains with solubles; EOA = WD + microencapsulated organic acids-essential oils blend at 300 mg/kg; PRO = WD + PRO at 125 mg/kg; and EOA + PRO = WD + microencapsulated organic acids-essential oils blend at 300 mg/kg + PRO at 125 mg/kg. DDGS, distillers dried grains with solubles; WD, wheat and corn DDGS; EOA, essential oils with organic acids; PRO, protease.

expression of ZO-1 and occludin did not differ between the four dietary treatments ($p > 0.05$). There was a trend towards higher expression of claudin-1 mRNA in the EOA+PRO group compared to the others ($p = 0.062$). On day 35, the expression of ZO-1 mRNA did not differ significantly between treatments ($p > 0.05$). The EOA and EOA+PRO groups had higher claudin-1 mRNA expression than the WD and PRO groups ($p < 0.01$). Occludin mRNA expression was

higher in the EOA and PRO groups than in the WD group ($p < 0.01$), while the EOA+PRO group had similar expression to the other groups ($p > 0.05$).

Intestinal morphology

On day 14, no significant differences in gut morphology were observed among the four dietary treatments ($p > 0.05$), as indicated in Table 4. On day 35, there were no differences in jejunal VH and VW between treatments ($p > 0.05$). However, the EOA+PRO group had a lower jejunal crypt depth and a higher VH/CD ratio compared to the other treatment groups ($p < 0.01$).

Microbial metabolites

Table 5 illustrates the effects of the dietary treatments on the microbial metabolites in the cecal content on day 35. No significant differences in the ammonia and VFA content were found among the dietary treatments ($p > 0.05$). In terms of biogenic amines, the WD group had a higher putrescine content than the other dietary treatments ($p < 0.05$). In addition, the WD group tended to have a higher cadaverine content than the other dietary treatments ($p < 0.1$).

DISCUSSION

Growth performance

In this study, body weight gain, feed intake, and mortality rate did not differ significantly among the dietary treatments, all of which were based on a basal diet containing a high proportion of wheat and corn DDGS. However, the FCR for the EOA+PRO group was lower than that of the WD group from day 1 to 35, whereas the EOA and PRO groups did not differ substantially from the others. The possible explanation could be the combined effect of EOA and PRO, which could improve the FCR of the birds by stimulating digestive enzyme activity and improving nutrient utilization under the challenging conditions of high wheat and corn DDGS in the diet. Several studies have shown that EOA can stimulate the activity of digestive enzymes and improve feed

Table 4. Jejunal morphology (mean¹⁾) of broiler chickens fed a high wheat and corn DDGS diet²⁾ supplemented with microencapsulated organic acids-essential oils blend and protease

| Trait | WD | EOA | PRO | EOA + PRO | Pooled SEM ³⁾ | p-value |
|---------------------|---------------------|---------------------|---------------------|--------------------|--------------------------|---------|
| Day 14 | | | | | | |
| Villus height (µm) | 572 | 542 | 600 | 545 | 40.29 | 0.159 |
| Villus width (µm) | 103 | 92 | 110 | 107 | 11.01 | 0.123 |
| Crypt depth (µm) | 110 | 107 | 116 | 110 | 11.05 | 0.722 |
| VH/CD ⁴⁾ | 5.1 | 5.2 | 5.3 | 5.2 | 0.57 | 0.965 |
| Day 35 | | | | | | |
| Villus height (µm) | 683.00 | 727.47 | 746.77 | 734.74 | 46.22 | 0.239 |
| Villus width (µm) | 85.08 | 95.99 | 106.19 | 84.86 | 14.21 | 0.117 |
| Crypt depth (µm) | 115.93 ^a | 125.03 ^a | 120.71 ^a | 93.68 ^b | 13.54 | 0.009 |
| VH/CD ⁴⁾ | 5.88 ^b | 6.03 ^b | 6.38 ^b | 7.90 ^a | 0.55 | <0.001 |

¹⁾Each value represents the mean of 14 replicates.
²⁾Dietary treatments: WD = corn-soybean meal basal diet with wheat and corn distillers dried grains with solubles; EOA = WD + microencapsulated organic acids-essential oils blend at 300 mg/kg; PRO = WD + PRO at 125 mg/kg; and EOA + PRO = WD + microencapsulated organic acids-essential oils blend at 300 mg/kg + PRO at 125 mg/kg.
³⁾Pooled SEM (n = 56).
⁴⁾Villus height per crypt depth ratio.
^{a,b}Within a row, means with different superscripts differ significantly ($p < 0.05$).
DDGS, distillers dried grains with solubles; WD, wheat and corn DDGS; EOA, essential oils with organic acids; PRO, protease; VH, villus height; CD, crypt depth.

Table 5. Microbial metabolites (mean¹⁾) in cecal content of broiler chickens fed a high wheat and corn DDGS diet²⁾ supplemented with microencapsulated organic acids-essential oils blend and protease at 35 days of age

| Trait | WD | EOA | PRO | EOA + PRO | Pooled SEM ³⁾ | p-value |
|--|-----------------|-----------------|-----------------|-----------------|--------------------------|---------|
| Ammonia (mg/g wet content) | 6.67 | 6.86 | 6.86 | 6.62 | 0.72 | 0.966 |
| Biogenic amine (µg/g wet content) | | | | | | |
| Putrescine | 84 ^a | 33 ^b | 34 ^b | 31 ^b | 26 | 0.013 |
| Cadaverine | 1576 | 1302 | 1203 | 1180 | 240 | 0.087 |
| Volatile fatty acid (mmol/g wet content) | | | | | | |
| Acetic acid | 69.15 | 73.77 | 69.55 | 78.93 | 12.36 | 0.646 |
| Propionic acid | 4.07 | 4.5 | 4.25 | 3.78 | 0.92 | 0.713 |
| Butyric acid | 4.32 | 4.08 | 4.17 | 4.42 | 1.3 | 0.983 |

¹⁾Each value represents the mean of 14 replicates except volatile fatty acid represents 11 replicates.

²⁾Dietary treatments: WD = corn-soybean meal basal diet with wheat and corn distillers dried grains with solubles; EOA = WD + microencapsulated organic acids-essential oils blend at 300 mg/kg; PRO = WD + PRO at 125 mg/kg; and EOA + PRO = WD + microencapsulated organic acids-essential oils blend at 300 mg/kg + PRO at 125 mg/kg.

³⁾Pooled standard error of mean (n = 56).

^{a,b}Within a column, means with different superscripts differ significantly ($p < 0.05$).

DDGS, distillers dried grains with solubles; WD, wheat and corn DDGS; EOA, essential oils with organic acids; PRO, protease.

efficiency in broiler chickens [45,46]. In addition, administration of a single-component enzyme (serine alkaline endopeptidase) in broilers also improved ADG and FCR in a rye-wheat-soybean meal [18] and corn-soybean meal-canola-based diets [47]. Chowdhury et al. [30] partially confirm the results of this study, showing that broilers fed a diet supplemented with microencapsulated EOA and PRO achieved better FCR than those fed EOA alone. In addition, they found that higher EOA content (300 mg/kg diet) increased FCR regardless of whether PRO was included in the diet or not.

Expression of intestinal barrier tight junction proteins

TJ proteins, including claudins, occludins, ZO-1, and the actin-myosin cytoskeleton, establish connections between layers of epithelial cells in the intestine and form a barrier that separates the lumen contents from the underlying tissue [48,49]. These tight junctions are essential elements of the intestinal epithelial barrier and play a crucial role in maintaining the integrity of the gastrointestinal tract. When this barrier is compromised, luminal antigens such as microbes and toxins can disrupt homeostasis and increase the risk of systemic infection, chronic inflammation, and malabsorption [48,50]. The breakdown of the intestinal barrier has been associated with the pathogenicity of specific gut bacteria, including *Campylobacter jejuni*, *Salmonella enterica* and *Clostridium perfringens* [51]. In this study, the additives EOA, PRO, and EOA+PRO had no effect on ZO-1 mRNA expression. The higher expression of claudin-1 mRNA in the EOA and EOA+PRO groups compared to the WD control group suggests an improvement in gut integrity when the diet is supplemented with these additives. There was no discernible difference between the PRO and the WD control groups, suggesting that the increased claudin-1 mRNA expression in the EOA+PRO group may be due to the effect of EOA rather than PRO. It is possible that claudin-1 mRNA expression was upregulated due to the antibacterial properties of EOA. In addition, Yang et al. [28] observed that the EOA group expressed more claudin-1 mRNA than the antibiotic group or the control group, but there was no significant change in the mRNA expression of occludin or ZO-1. McKnight et al. [52] observed comparable levels of claudin-1 mRNA expression in both the EOA and antibiotic groups, which were higher than those in the control group.

In this study, the EOA and PRO groups showed a higher level of occludin mRNA expression than the WD group, while EOA+PRO was not significantly different from the others. This result

suggests that either the mixture of organic acids and essential oils or the PRO can stimulate occludin by upregulating occludin mRNA expression without a combination effect of EOA and PRO in the EOA+PRO group. The combination of essential oils and organic acids has been shown to be beneficial, e.g. in terms of improved feed efficiency or upregulated mRNA expression of TJ proteins such as claudin-1 and occludin when added to broiler diets [28,43,45,52].

Intestinal morphology

Morphological indicators of intestinal health, such as VH, CD and the VH/CD ratio, provide information about the ability of the intestine to digest and absorb nutrients [53,54]. Higher villi generally indicate a healthier gut, as they provide a larger surface area for nutrient absorption, while shallower crypts are typically associated with a healthier gut, as deeper crypts may indicate increased cell turnover or pathological conditions [54,55]. A higher VH/CD ratio usually reflects a well-functioning and healthy gut, while a lower ratio may indicate problems such as inflammation or impaired nutrient absorption [55]. In addition, a lower VH/CD ratio indicates a reduced number of absorptive cells and an increased number of goblet cells, leading to increased mucin secretion [55,56]. Changes in mucin quantity or composition may impair nutrient uptake or increase energy requirements to maintain homeostasis [55,57]. The addition of EOA to broiler diets has been shown to be an effective strategy to improve gut morphology [45,46,58]. These results could not be confirmed in this study, as supplementation with EOA did not produce any significant effects on intestinal morphology. However, the EOA+PRO group showed increased jejunal VH/CD ratio and decreased CD, suggesting a combination effect of PRO supplementation in combination with EOA on intestinal morphology. The discrepancies between the present study and previous research may be due to differences in dietary formulations, microbial and environmental conditions, methodological approaches, and the synergistic effects of the supplements used.

The possible mechanisms of EOA and PRO that improved the expression of TJ proteins and intestinal morphology under nutritional challenge in this study might be related to toll-like receptors (TLRs), which are part of the innate immune system, recognize pathogens and trigger inflammatory reactions [59]. Excessive activation of TLRs can lead to chronic intestinal inflammation, which damages the intestinal mucosa, disrupts tight junctions and increases intestinal permeability [50,60]. EOA, which contain antimicrobial and anti-inflammatory compounds such as thymol and carvacrol, influence signaling through TLRs by reducing exposure to pathogens and attenuating excessive inflammatory responses. This in turn contributes to the maintenance or improvement of tight junction protein expression and intestinal morphology [45]. While PRO enzymes support gut health by improving protein digestion, which helps maintain tight junction integrity and enhance gut morphology [26]. Efficient protein breakdown prevents excessive stress on TJ proteins and reduces gut inflammation, leading to better gut barrier function and healthier intestinal structure [26,50].

Microbial metabolites

In this study, the lower putrescine levels in the EOA, PRO and EOA+PRO groups compared to the WD group may be due to the suppression of putrefactive proteins and microbes in the gut. Previous studies have shown that the combination of essential oils and organic acids reduces the prevalence of pathogenic bacteria such as *Clostridium perfringens*, *Escherichia coli* and *Salmonella* spp., while beneficial bacteria such as *Lactobacilli* increase [10–12]. This change in microbial composition could explain the lower putrescine levels observed. In addition, the improved protein and amino acid digestibility in birds fed PRO-containing diets may have limited the nutrients available for microbial growth, thereby reducing microbial metabolites [61, 62]. Several studies have also found

a decrease pathogenic microbial populations such as in *Clostridium perfringens*, *Escherichia coli* and *Salmonella* spp. in the ileum of broilers fed diets containing PRO [62–64]. Park and Kim [65] found that the combined effect of essential oils and PRO on reducing ammonia emissions may be due to their role in enhancing nitrogen retention, although this combination did not show a synergistic effect on growth performance or bacterial counts.

Volatile fatty acids are associated with microbial fermentation in the hindgut [66]. Low quality dietary proteins can increase the content of VFA in the cecum. For example, Meyer et al. [39] reported that the addition of feather meal at 5% increased the propionic acid concentration in the ceca of laying hens. The use of corn gluten [67] or DDGS [14] as a protein source in broiler feed increased propionic acid and butyric acid level in the ceca. Yang et al. [28] showed a significant increase in butyric acid with a tendency to increase acetic acid and total short-chain fatty acids in the ileal contents of the EOA-supplemented group compared to the antibiotic group, with no significant difference observed compared to the control group. It was anticipated that dietary treatments or feed additives would modify the microbial substrate, thereby altering VFA levels in the ceca. However, in the present study, no significant effects of dietary treatments on cecal VFA levels were observed. This lack of effect may be attributed to the low inclusion level of the essential oils blend at 300 mg/kg, which might not have been sufficient to induce detectable differences in cecal VFA concentrations. In contrast, a study by Ceylan et al. [68] demonstrated that higher levels of essential oils, at 700 or 1,200 mg/kg, significantly increased cecal acetate, propionate, butyrate, and total short-chain fatty acid concentrations in broilers.

The non-significant differences in VFA levels observed in our study could also be related to the high absorption rate of VFAs in the lower intestinal tract. VFA absorption in the ceca occurs rapidly, reducing existing VFA concentrations and facilitating the renewal of cecal contents [69]. Over 95% of VFAs produced from fermentation are ionized at the prevailing pH of the large intestine and are actively absorbed by Na⁺-coupled monocarboxylate transport proteins (SMCT1). Meanwhile, the non-dissociated form is transported by the H⁺-coupled low-affinity monocarboxylate transporter protein (MCT1) [69,70]. Both transporters function concurrently in poultry to maximize VFA absorption across a wide range of lumen pH levels [71,72].

CONCLUSIONS

In summary, dietary supplementation with a combination of EOA and PRO improved growth performance by improving feed efficiency in broiler chickens fed a high wheat and corn DDGS diet. This improvement was accompanied by better gut health as evidenced by reduced jejunal crypt depth, increased VH/CD ratio and increased mRNA expression of the tight junction protein claudin-1. In addition, both the combined treatment with EOA and PRO and the individual EOA and PRO supplements significantly reduced putrescine levels in the hindgut. Further studies are recommended to better understand the actual mechanism of action of these changes in the gut of broiler chickens.

REFERENCES

1. Bean-Hodgins L, Kiarie EG. Mandated restrictions on the use of medically important antibiotics in broiler chicken production in Canada: implications, emerging challenges, and opportunities for bolstering gastrointestinal function and health — a review. *Can J Anim Sci.* 2021;101:602-29. <https://doi.org/10.1139/cjas-2021-0015>
2. Yang Y, Ashworth AJ, Willett C, Cook K, Upadhyay A, Owens PR, et al. Review of antibiotic

- resistance, ecology, dissemination, and mitigation in U.S. broiler poultry systems. *Front Microbiol.* 2019;10:2639. <https://doi.org/10.3389/fmicb.2019.02639>
3. Ma J, Mahfuz S, Wang J, Piao X. Effect of dietary supplementation with mixed organic acids on immune function, antioxidative characteristics, digestive enzymes activity, and intestinal health in broiler chickens. *Front Nutr.* 2021;8:673316. <https://doi.org/10.3389/fnut.2021.673316>
 4. Miao S, Zhou W, Li H, Zhu M, Dong X, Zou X. Effects of coated sodium butyrate on production performance, egg quality, serum biochemistry, digestive enzyme activity, and intestinal health of laying hens. *Ital J Anim Sci.* 2021;20:1452-61. <https://doi.org/10.1080/1828051X.2021.1960209>
 5. Agboola AF, Omidwura BRO, Odu O, Popoola IO, Iyayi E. Effects of organic acid and probiotic on performance and gut morphology in broiler chickens. *S Afr J Anim Sci.* 2015;45:494-501. <https://doi.org/10.4314/sajas.v45i5.6>
 6. Sun HY, Zhou HB, Liu Y, Wang Y, Zhao C, Xu LM. Comparison of organic acids supplementation on the growth performance, intestinal characteristics and morphology, and cecal microflora in broilers fed corn-soybean meal diet. *Anim Biosci.* 2022;35:1689-97. <https://doi.org/10.5713/ab.21.0448>
 7. Bassolé IHN, Juliani HR. Essential oils in combination and their antimicrobial properties. *Molecules.* 2012;17:3989-4006. <https://doi.org/10.3390/molecules17043989>
 8. Lee KW, Everts H, Beynen AC. Essential oils in broiler nutrition. *Int J Poult Sci.* 2004;3:738-52. <https://doi.org/10.3923/ijps.2004.738.752>
 9. Brenes A, Roura E. Essential oils in poultry nutrition: main effects and modes of action. *Anim Feed Sci Technol.* 2010;158:1-14. <https://doi.org/10.1016/j.anifeedsci.2010.03.007>
 10. El-Dayem GA, Rasheed NM, Ghanem R, Albadrani GM, Abdel-Daim M, Abdel-Latif MA. Essential oils and organic acid blends potentially enhance growth and ameliorate enteritis and inflammatory response after challenge with *Clostridium perfringens* in chickens. *Damanhour J Vet Sci.* 2023;10:19-27. <https://doi.org/10.21608/djvs.2023.230008.1120>
 11. Iqbal H, Rahman A, Khanum S, Arshad M, Badar IH, Asif AR, et al. Effect of essential oil and organic acid on performance, gut health, bacterial count and serological parameters in broiler. *Braz J Poult Sci.* 2021;23:1-10. <https://doi.org/10.1590/1806-9061-2021-1443>
 12. Islam Z, Sultan A, Khan SZ, Khan SB. Effect of organic acids blend, micro-encapsulated phyto-essential oils individually or in combination on growth performance, gut health and nutrients utilization of broilers. *Pak J Zool.* 2022;54:2391-9. <https://doi.org/10.17582/journal.pjz/20210714190714>
 13. Martinez-Amezcuca C, Parsons CM, Singh V, Srinivasan R, Murthy GS. Nutritional characteristics of corn distillers dried grains with solubles as affected by the amounts of grains versus solubles and different processing techniques. *Poult Sci.* 2007;86:2624-30. <https://doi.org/10.3382/ps.2007-00137>
 14. Barekatin MR, Antipatis C, Rodgers N, Walkden-Brown SW, Iji PA, Choct M. Evaluation of high dietary inclusion of distillers dried grains with solubles and supplementation of protease and xylanase in the diets of broiler chickens under necrotic enteritis challenge. *Poult Sci.* 2013;92:1579-94. <https://doi.org/10.3382/ps.2012-02786>
 15. Moradi A, Moradi S, Abdollahi MR. Influence of feed ingredients with pellet-binding properties on physical pellet quality, growth performance, carcass characteristics and nutrient retention in broiler chickens. *Anim Prod Sci.* 2018;59:73-81. <https://doi.org/10.1071/AN17109>
 16. Arabshahi HA, Ghasemi HA, Hajkhodadadi I, Khaltabadi Farahani AH. Effects of multicarbohydrase and butyrate glycerides on productive performance, nutrient digestibility,

- gut morphology, and ileal microbiota in late-phase laying hens fed corn- or wheat-based diets. *Poult Sci.* 2021;100:101066. <https://doi.org/10.1016/j.psj.2021.101066>
17. Gao F, Jiang Y, Zhou GH, Han ZK. The effects of xylanase supplementation on performance, characteristics of the gastrointestinal tract, blood parameters and gut microflora in broilers fed on wheat-based diets. *Anim Feed Sci Technol.* 2008;142:173-84. <https://doi.org/10.1016/j.anifeeds.2007.07.008>
18. Yan F, Dibner JJ, Knight CD, Vazquez-Anon M. Effect of carbohydrase and protease on growth performance and gut health of young broilers fed diets containing rye, wheat, and feather meal. *Poult Sci.* 2017;96:817-28. <https://doi.org/10.3382/ps/pew300>
19. Choct M, Hughes RJ, Wang J, Bedford MR, Morgan AJ, Annison G. Increased small intestinal fermentation is partly responsible for the anti-nutritive activity of non-starch polysaccharides in chickens. *Br Poult Sci.* 1996;37:609-21. <https://doi.org/10.1080/00071669608417891>
20. Bedford MR, Apajalahti JH. The role of feed enzymes in maintaining poultry intestinal health. *J Sci Food Agric.* 2022;102:1759-70. <https://doi.org/10.1002/jsfa.11670>
21. Yokoo K, Yamamoto Y, Suzuki T. Ammonia impairs tight junction barriers by inducing mitochondrial dysfunction in Caco-2 cells. *Faseb J.* 2021;35:e21854. <https://doi.org/10.1096/fj.202100758R>
22. Saleh AA, Dawood MM, Badawi NA, Ebeid TA, Amber KA, Azzam MM. Effect of supplemental serine-protease from *Bacillus licheniformis* on growth performance and physiological change of broiler chickens. *J Appl Anim Res.* 2020;48:86-92. <https://doi.org/10.1080/09712119.2020.1732986>
23. Jabbar A, Tahir M, Alhidary IA, Abdelrahman MA, Albadani H, Khan RU, et al. Impact of microbial protease enzyme and dietary crude protein levels on growth and nutrients digestibility in broilers over 15–28 days. *Animals.* 2021;11:2499. <https://doi.org/10.3390/ani11092499>
24. Sarica S, Yurtseven S, Polat I. Supplementing diets for broilers that are low in crude protein and amino acids with protease. *S Afr J Anim Sci.* 2020;50:521-8. <https://doi.org/10.4314/sajas.v50i4.4>
25. Ding XM, Li DD, Li ZR, Wang JP, Zeng QF, Bai SP, et al. Effects of dietary crude protein levels and exogenous protease on performance, nutrient digestibility, trypsin activity and intestinal morphology in broilers. *Livest Sci.* 2016;193:26-31. <https://doi.org/10.1016/j.livsci.2016.09.002>
26. Duque-Ramírez CF, Javierre JA, Peñuela-Sierra LM, Diaz-Vargas M. Effect of exogenous protease on performance, nutrient digestibility, intestinal histomorphometric, meat quality characteristics, carcass yield in broilers fed low protein diets. *Trop Anim Health Prod.* 2023; 55:190. <https://doi.org/10.1007/s11250-023-03562-y>
27. Yang Y, Wang Q, Diarra MS, Yu H, Hua Y, Gong J. Functional assessment of encapsulated citral for controlling necrotic enteritis in broiler chickens. *Poult Sci.* 2016;95:780-9. <https://doi.org/10.3382/ps/pev375>
28. Yang X, Liu Y, Yan F, Yang C, Yang X. Effects of encapsulated organic acids and essential oils on intestinal barrier, microbial count, and bacterial metabolites in broiler chickens. *Poult Sci.* 2019;98:2858-65. <https://doi.org/10.3382/ps/pez031>
29. Chen H, Ma D, Li Y, Liu Y, Wang Y. Optimization the process of microencapsulation of *Bifidobacterium bifidum* BB01 by Box-Behnken design. *Acta Univ Cibiniensis Ser E Food Technol.* 2016;20:17-28. <https://doi.org/10.1515/auaft-2016-0012>
30. Chowdhury K, Dhara AK, Dey S, Sarkar A, Tactacan GB, Halder S. Dietary supplementation of a microencapsulated essential oil-organic acid blend with and without a protease affects performance, nutrient accretion, cecal microbiota, and circulatory biomarker concentrations in

- male broiler chickens: essential oil-organic acid blend and protease supplementation in broiler chickens. *Lett Anim Biol*. 2021;1:19-32. <https://doi.org/10.62310/liab.v1i1.51>
31. Commission Regulation (EC) No 1831/2003. Regulation (EC) no 1831/2003 of the European Parliament and of the Council of 22 September 2003 on additives for use in animal nutrition (text with EEA relevance) [Internet]. OJ L 268. 2003 [cited 2003 Oct 18]. <http://data.europa.eu/eli/reg/2003/>
 32. Aviagen. Ross 308 broiler nutrition specifications. Aviagen; 2019. Report No.: PO2019-EN.
 33. AOAC (Association of Official Analytical Chemists). Official methods of analysis. AOAC; 2000.
 34. Gilani S, Howarth GS, Natrass G, Kitessa SM, Barekatin R, Forder REA, et al. Gene expression and morphological changes in the intestinal mucosa associated with increased permeability induced by short-term fasting in chickens. *J Anim Physiol Anim Nutr*. 2018;102:e653-61. <https://doi.org/10.1111/jpn.12808>
 35. Chen J, Tellez G, Richards JD, Escobar J. Identification of potential biomarkers for gut barrier failure in broiler chickens. *Front Vet Sci*. 2015;2:1-10. <https://doi.org/10.3389/fvets.2015.00014>
 36. Taylor SC, Nadeau K, Abbasi M, Lachance C, Nguyen M, Fenrich J. The ultimate qPCR experiment: producing publication quality, reproducible data the first time. *Trends Biotechnol*. 2019;37:761-74. <https://doi.org/10.1016/j.tibtech.2018.12.002>
 37. Iji PA, Hughes RJ, Choct M, Tivey DR. Intestinal structure and function of broiler chickens on wheat-based diets supplemented with a microbial enzyme. *Asian-Australas J Anim Sci*. 2001;14:54-60. <https://doi.org/10.5713/ajas.2001.54>
 38. Thanh NT, Loh TC, Foo HL, Hair-Bejo M, Azhar BK. Effects of feeding metabolite combinations produced by *Lactobacillus plantarum* on growth performance, faecal microbial population, small intestine villus height and faecal volatile fatty acids in broilers. *Br Poult Sci*. 2009;50:298-306. <https://doi.org/10.1080/00071660902873947>
 39. Meyer B, Bessei W, Vahjen W, Zentek J, Harlander-Matauschek A. Dietary inclusion of feathers affects intestinal microbiota and microbial metabolites in growing Leghorn-type chickens. *Poult Sci*. 2012;91:1506-13. <https://doi.org/10.3382/ps.2011-01786>
 40. Weatherburn MW. Phenol-hypochlorite reaction for determination of ammonia. *Anal Chem*. 1967;39:971-4. <https://doi.org/10.1021/ac60252a045>
 41. Saarinen MT. Determination of biogenic amines as dansyl derivatives in intestinal digesta and feces by reversed phase HPLC. *Chromatographia*. 2002;55:297-300. <https://doi.org/10.1007/BF02491662>
 42. Eerola S, Hinkkanen R, Lindfors E, Hirvi T. Liquid chromatographic determination of biogenic amines in dry sausages. *J AOAC Int*. 1993;76:575-7. <https://doi.org/10.1093/jaoac/76.3.575>
 43. Stefanello C, Rosa DP, Dalmoro YK, Segatto AL, Vieira MS, Moraes ML, et al. Protected blend of organic acids and essential oils improves growth performance, nutrient digestibility, and intestinal health of broiler chickens undergoing an intestinal challenge. *Front Vet Sci*. 2020;6:491. <https://doi.org/10.3389/fvets.2019.00491>
 44. Bedford M. Removal of antibiotic growth promoters from poultry diets: implications and strategies to minimise subsequent problems. *World Poult Sci J*. 2000;56:347-65. <https://doi.org/10.1079/WPS20000024>
 45. Pham VH, Kan L, Huang J, Geng Y, Zhen W, Guo Y, et al. Dietary encapsulated essential oils and organic acids mixture improves gut health in broiler chickens challenged with necrotic enteritis. *J Anim Sci Biotechnol*. 2020;11:18. <https://doi.org/10.1186/s40104-019-0421-y>
 46. Yang X, Xin H, Yang C, Yang X. Impact of essential oils and organic acids on the growth

- performance, digestive functions and immunity of broiler chickens. *Anim Nutr.* 2018;4:388-93. <https://doi.org/10.1016/j.aninu.2018.04.005>
47. Law FL, Zulkifli I, Soleimani AF, Liang JB, Awad EA. The effects of low-protein diets and protease supplementation on broiler chickens in a hot and humid tropical environment. *Asian-Australas J Anim Sci.* 2018;31:1291-300. <https://doi.org/10.5713/ajas.17.0581>
48. An J, Liu Y, Wang Y, Fan R, Hu X, Zhang F, et al. The role of intestinal mucosal barrier in autoimmune disease: a potential target. *Front Immunol.* 2022;13:871713. <https://doi.org/10.3389/fimmu.2022.871713>
49. Suzuki T. Regulation of the intestinal barrier by nutrients: the role of tight junctions. *Anim Sci J.* 2020;91:e13357. <https://doi.org/10.1111/asj.13357>
50. Ducatelle R, Goossens E, Eeckhaut V, Van Immerseel F. Poultry gut health and beyond. *Anim Nutr.* 2023;13:240-8. <https://doi.org/10.1016/j.aninu.2023.03.005>
51. Awad WA, Hess C, Hess M. Enteric pathogens and their toxin-induced disruption of the intestinal barrier through alteration of tight junctions in chickens. *Toxins.* 2017;9:60. <https://doi.org/10.3390/toxins9020060>
52. McKnight LL, Peppler W, Wright DC, Page G, Han Y. A blend of fatty acids, organic acids, and phytochemicals induced changes in intestinal morphology and inflammatory gene expression in coccidiosis-vaccinated broiler chickens. *Poult Sci.* 2019;98:4901-8. <https://doi.org/10.3382/ps/pez241>
53. Montagne L, Pluske JR, Hampson DJ. A review of interactions between dietary fibre and the intestinal mucosa, and their consequences on digestive health in young non-ruminant animals. *Anim Feed Sci Technol.* 2003;108:95-117. [https://doi.org/10.1016/S0377-8401\(03\)00163-9](https://doi.org/10.1016/S0377-8401(03)00163-9)
54. Zang JJ, Piao XS, Huang DS, Wang JJ, Ma X, Ma YX. Effects of feed particle size and feed form on growth performance, nutrient metabolizability and intestinal morphology in broiler chickens. *Asian-Australas J Anim Sci.* 2009;22:107-12. <https://doi.org/10.5713/ajas.2009.80352>
55. Alagbe EO, Schulze H, Adeola O. Growth performance, nutrient digestibility, intestinal morphology, cecal mucosal cytokines and serum antioxidant responses of broiler chickens to dietary enzymatically treated yeast and coccidia challenge. *J Anim Sci Biotechnol.* 2023;14:57. <https://doi.org/10.1186/s40104-023-00846-z>
56. Hamed S, Rezaian M, Shomali T. Histological changes of small intestinal mucosa of cocks due to sunflower meal single feeding. *Am J Anim Vet Sci.* 2011;6:171-5. <https://doi.org/10.3844/ajavsp.2011.171.175>
57. Faderl M, Noti M, Corazza N, Mueller C. Keeping bugs in check: the mucus layer as a critical component in maintaining intestinal homeostasis. *IUBMB Life.* 2015;67:275-85. <https://doi.org/10.1002/iub.1374>
58. Zhang S, Shen YR, Wu S, Xiao YQ, He Q, Shi SR. The dietary combination of essential oils and organic acids reduces *Salmonella enteritidis* in challenged chicks. *Poult Sci.* 2019;98:6349-55. <https://doi.org/10.3382/ps/pez457>
59. Arancibia SA, Beltrán CJ, Aguirre IM, Silva P, Peralta AL, Malinarich F, et al. Toll-like receptors are key participants in innate immune responses. *Biol Res.* 2007;40:97-112. <https://doi.org/10.4067/S0716-97602007000200001>
60. Gilbert MS, Ijssennagger N, Kies AK, van Mil SWC. Protein fermentation in the gut; implications for intestinal dysfunction in humans, pigs, and poultry. *Am J Physiol Gastrointest Liver Physiol.* 2018;315:G159-70. <https://doi.org/10.1152/ajpgi.00319.2017>
61. Borda-Molina D, Zuber T, Siegert W, Camarinha-Silva A, Feuerstein D, Rodehutschord M. Effects of protease and phytase supplements on small intestinal microbiota and amino acid

- digestibility in broiler chickens. *Poult Sci.* 2019;98:2906-18. <https://doi.org/10.3382/ps/pez038>
62. Chowdhury MAK, Dhara AK, Dey S, Sarkar A, Haldar S, Tactacan GB. Protease complex reduces potentially pathogenic microbial populations in the ileum while optimizing performance of broiler chickens. *Anim Vet Sci.* 2021;9:16-23. <https://doi.org/10.11648/j.avs.20210901.13>
63. Giannenas I, Bonos E, Anestis V, Filioussis G, Papanastasiou DK, Bartzanas T, et al. Effects of protease addition and replacement of soybean meal by corn gluten meal on the growth of broilers and on the environmental performances of a broiler production system in Greece. *PLOS ONE.* 2017;12:e0169511. <https://doi.org/10.1371/journal.pone.0169511>
64. Kamel NF, Naela, Ragaa M, El-Banna RA, Mohamed FF. Effects of a monocomponent protease on performance parameters and protein digestibility in broiler chickens. *Agric Agric Sci Procedia.* 2015;6:216-25. <https://doi.org/10.1016/j.aaspro.2015.08.062>
65. Park JH, Kim IH. Effects of a protease and essential oils on growth performance, blood cell profiles, nutrient retention, ileal microbiota, excreta gas emission, and breast meat quality in broiler chicks. *Poult Sci.* 2018;97:2854-60. <https://doi.org/10.3382/ps/pey151>
66. Cao BH, Karasawa Y, Guo YM. Effects of green tea polyphenols and fructo-oligosaccharides in semi-purified diets on broilers' performance and caecal microflora and their metabolites. *Asian-Australas J Anim Sci.* 2005;18:85-9. <https://doi.org/10.5713/ajas.2005.85>
67. Qaisrani SN, van Krimpen MM, Verstegen MWA, Hendriks WH, Kwakkel RP. Effects of three major protein sources on performance, gut morphology and fermentation characteristics in broilers. *Br Poult Sci.* 2020;61:43-50. <https://doi.org/10.1080/00071668.2019.1671958>
68. Ceylan N, Yenice E, Yavaş İ, Çenesiz AA, Toprak NN, Çiftçi İ. Comparative effects of medium-chain fatty acids or phytobiotics-based feed additives on performance, caecum microbiota, volatile fatty acid production and intestinal morphology of broilers. *Vet Med Sci.* 2023;9:2719-30. <https://doi.org/10.1002/vms3.1249>
69. Moran ET Jr, Bedford MR. Large intestinal dynamics differ between fowl and swine: anatomical modifications, microbial collaboration, and digestive advantages from fibrolytic enzymes. *Anim Nutr.* 2022;11:160-70. <https://doi.org/10.1016/j.aninu.2022.07.004>
70. Engelhardt WV, Rechkemmer G. Absorption of inorganic ions and short-chain fatty acids in the colon of mammals. In: Gilles-Baillien M, Gilles R, editors. *Intestinal transport. Fundamental and comparative aspects.* Springer; 1983. p. 26-45.
71. Holtug K, McEwan GTA, Skadhauge E. Effects of propionate on the acid microclimate of hen (*Gallus domesticus*) colonic mucosa. *Comp Biochem Physiol A Physiol.* 1992;103:649-52. [https://doi.org/10.1016/0300-9629\(92\)90160-R](https://doi.org/10.1016/0300-9629(92)90160-R)
72. Herrmann J, Hermes R, Breves G. Transepithelial transport and intraepithelial metabolism of short-chain fatty acids (SCFA) in the porcine proximal colon are influenced by SCFA concentration and luminal pH. *Comp Biochem Physiol A Mol Integr Physiol.* 2011;158:169-76. <https://doi.org/10.1016/j.cbpa.2010.10.018>