


Supplementation effects of feed additive containing eggshell, *Schisandra chinensis* by-products combined with multi-probiotics strains in laying hens

Ah-Ran Lee¹, Sol-Hee Kim², Hangsul Cho², Yan-Qing Wang², Soo-Ki Kim^{2*}

¹Animal Resources Research Center, Konkuk University, Seoul 05029, Korea

²Department of Animal Science and Technology, Konkuk University, Seoul 05029, Korea



Received: Sep 20, 2024

Revised: Nov 30, 2024

Accepted: Dec 3, 2024

*Corresponding author

Soo-Ki Kim

Department of Animal Science and Technology, Konkuk University, Seoul 05029, Korea.

Tel: +82-2-450-3728

E-mail: sookikim@konkuk.ac.kr

Copyright © 2025 Korean Society of Animal Science and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Ah-Ran Lee

<https://orcid.org/0000-0001-7163-0059>

Sol-Hee Kim

<https://orcid.org/0009-0003-0803-4942>

Hangsul Cho

<https://orcid.org/0009-0009-7756-5302>

Yan-Qing Wang

<https://orcid.org/0009-0006-5848-031X>

Soo-Ki Kim

<https://orcid.org/0000-0003-3499-3330>

Competing interests

No potential conflict of interest relevant to this article was reported.

Abstract

Eggshell waste and *Schisandra chinensis* by-products are natural sources rich in beneficial nutrients and bioactive compounds. However, their combined effects with multi-probiotics on poultry productivity and health remain unexplored. This study assessed the immediate effects of a feed additive—eggshell waste (ES), *Schisandra chinensis* by-product (SC), and multi-probiotics (M)—administered for four weeks to aged laying hens before slaughter, evaluating the improvements of laying performance, egg quality, blood characteristics, visceral organs, tibia, and cecal microbiota. A total of 216 Hy-line Brown laying hens (70-week-old) were assigned to four dietary treatments consisting of 9 replicates of 6 birds in a completely randomized design. The combination of eggshell waste, *Schisandra chinensis* by-product, and multi-probiotics (ESM) of feed additive consisted of 40% eggshell, 5% SC, and 10^9 – 10^{11} CFU/g of multi-probiotic strains including *Bacillus subtilis*, *Bacillus licheniformis*, *Saccharomyces cerevisiae*, *Lactobacillus plantarum*, and supplemental nutrient premix. The treatment groups were as follows: corn-soybean meal-based basal diet (control); basal diet + 0.1% ESM; basal diet + 0.2% ESM, basal diet + 0.4% ESM. The total egg productivity rate during the experiment period tended to improve in ESM 0.2%, as compared with the control. The ESM 0.1% group increased egg weight ($p < 0.05$) while ESM 0.1% and ESM 0.2% tended to increase egg mass, compared to the control ($p = 0.051$). However, there was no significant difference in egg weight, feed intake, feed conversion ratio, and egg quality among the treatments. Furthermore, blood characteristics did not differ between the treatments, except for the total cholesterol contents, which was higher in ESM 0.4% treatment than the control ($p < 0.05$). ESM 0.4% supplementation showed a tendency for higher calcium, compared to the control. ESM 0.4% supplementation showed higher bone mineral density (BMD) of the tibia neck than the control ($p < 0.01$). All three ESM groups demonstrated a significant decrease in the abundance of *Bacteroidaceae* ($p < 0.05$), and an increase in the abundance of *Lactobacillaceae* at the family level ($p < 0.01$). In conclusion, ESM fed hens showed beneficial effects on the egg weight, egg mass, BMD of tibia neck, and cecal microbiota in laying hens.

Keywords: Eggshell waste, *Schisandra chinensis* by-product, Probiotic, Laying hen

Funding sources

This study was supported by the Korea Institute of Planning and Evaluation for Technology in Food, Agriculture and Forestry (IPET) through Technology Commercialization Support Program funded by Ministry of Agriculture, Food and Rural Affairs (MAFRA) (821035-03), Seoul, Korea. And this paper was supported by Konkuk University Researcher Fund in 2023.

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Kim SH, Kim SK.
Data curation: Kim SH, Kim SK.
Formal analysis: Lee AR, Kim SH.
Methodology: Kim SH, Cho H, Wang YQ.
Software: Lee AR, Kim SH, Cho H.
Validation: Lee AR.
Investigation: Lee AR, Kim SH.
Writing - original draft: Lee AR, Kim SK.
Writing - review & editing: Lee AR, Kim SH, Cho H, Wang YQ, Kim SK.

Ethics approval and consent to participate

The experimental protocol used in this study was approved by the Animal Care and Use Committee (IACUC) of Konkuk University, Korea (approval number: KU22233).

INTRODUCTION

While antibiotics have been continuously used to improve animal productivity, the overuse of antibiotics in animal, the environment, and human may continue [1,2]. This can lead to antibiotic resistance and side effects that make the proper treatment of disease impossible [3]. A feed additive needs to be developed that has an effect on animal growth that can replace that of antibiotics [4]. Interest in safe animal products and demand for antibiotic-free animal production is increasing [4,5]. With the expectation of improved growth performance, high quality and safe animal products, and disease prevention, the demand is rapidly increasing among researchers and consumers for multifunctional feed additives that combine animal food by-products, phytochemicals, and probiotics [5–7].

Eggshell waste is considered a potential calcium alternative in livestock production, and is produced in the order of 50,000 tonnes [8]. However, improper disposal of eggshell waste leads to the formation of ammonia, hydrogen sulfate, foul odor, and environmental pollution [9]. During eggshell calcification, approximately (5–6) g of calcium carbonate are deposited in the shell. The mineral composition of the eggshell includes Ca^{2+} , P^- , Na^+ , K^+ , HCO_3^- , and Mg^{2+} , all of which are essential minerals that meet the nutritional requirements for the growth and development of both laying hens and broiler [10,11]. The use of eggshell wastes in feed could contribute to the environmental safety, economic efficiency, productivity, and egg quality of laying hens.

Probiotics are live microorganisms, and have been used extensively as feed additives in the livestock industry [5,12]. The most common probiotics are *Lactobacillus*, *Bacillus*, *Bifidobacterium*, *Saccharomyces*, and *Enterococcus*; these improve the balance of gut microbes and prevent pathogen colonization, thereby improving growth, FCR, and health. These bacteria produce antimicrobial substances, such as organic acids and bacteriocins, to inhibit pathogenic microorganisms [12,13]. Numerous studies have been reported on the effect of the combination of probiotics and phytochemicals on growth performance, immune response, and gut microbiota in chickens [14–16]. Hidayat et al. [6] observed that the combination of probiotic *Lactobacillus acidophilus* (1.2 mL/day) and 4% phytochemicals (bay leaves, onion peels, and garlic peels) improved ileal histomorphology, ileal protein digestibility, and FCR. Lee et al. [17] reported that the *Artemisia Annua* fermented with *Lactobacillus plantarum* improved the Haugh unit value and prevented lipid oxidation of egg for 3 weeks storage, compared to the control and non-fermentation group, which suggesting higher antioxidant activity in the FA group.

Phytochemical substances are derived from plants, such as herbs, spices, and oleoresins, and are rich in bioactive compounds. They have been used as feed additives to improve animal productivity [18,19]. Among the various phytochemicals, *Schisandra chinensis* is well known as a high polyphenolic compound that has antioxidant, antimicrobial, antiviral, anticancer, and anti-inflammatory effects, and produces a substantial amount of the *S. chinensis* pomace [20]. *Schisandra chinensis* pomace contains higher levels of fiber, polyphenols, lignans, vitamins, and minerals than *S. chinensis* fruit, due to the concentration of these compounds during processing of the *S. chinensis* [21,22]. In addition, several studies have reported the effect of *S. chinensis* and pomace supplementation on improving the antioxidant activity, immunity in laying hens and physicochemical properties, and meat color stability in broilers [23–26].

Many studies and applications have explored the use of probiotics and phytochemicals as feed additives to improve productivity. However, few studies have investigated the effects of eggshells, *S. chinensis* by-products, and multi-probiotics on the laying performance and health of laying hens. Each of the feed additives—eggshell, *S. chinensis* by-products, and probiotics—has a distinctive nutritional value and a range of metabolites with beneficial physiological activities. The combined

nutritional and functional benefits of these additives are hypothesized to positively influence the productivity, blood profile, and gut health of laying hens. Therefore, this study aims to investigate the synergistic effects of these novel additives to enhance the productivity, egg quality, blood characteristics, visceral organs, tibia properties, and gut microbiota.

MATERIALS AND METHODS

Preparation of feed additive

The eggshell (ES) was produced and supplied by Poonglim Food. Briefly, ES membranes were removed by washing with water, followed by heating at 150°C for 12 h, and then the ES were crushed to a particle size of 1–5 mm using a hammer mill (SM-D3, Wilhelm Siefert GmbH). *Schisandra chinensis* by-products (SC) were obtained from a juice factory, Omija Valley, sun-dried for 24–48 h, and stored at 4°C, until use. Table 1 shows the source and composition of the feed additive, Biocalcium® (Hanong). The dietary supplement, combination of eggshell waste, *Schisandra chinensis* by-product, and multi-probiotics (ESM) consisted of 40% Eggshell, 5% by-products of *S. chinensis*, and 10^9 – 10^{11} CFU/g of Multi-probiotic strains, including *Bacillus subtilis*, *Bacillus licheniformis*, *Saccharomyces cerevisiae*, *Lactobacillus plantarum* (isolated from ES and SC), and supplemental nutrient premix with phytase.

Experimental animals and design

A total of 216 Hy-line Brown hens at 70 weeks of age were assigned to four dietary treatment groups: basal diet (control); basal diet+0.1% ESM; basal diet+0.2% ESM, basal diet+0.4% ESM. Each treatment consisted of nine replicates, with six birds each. All hens were housed in three-

Table 1. Ingredient and composition of feed additives (Biocalcium®)

Item	
Ingredient (%)	
Eggshell	40
<i>Schisandra chinensis</i> by-product	5
<i>Bacillus subtilis</i> powder (10^{11} CFU/g)	2
<i>Bacillus licheniformis</i> powder (10^{11} CFU/g)	2
<i>Saccharomyces cerevisiae</i> powder (10^{10} CFU/g)	2
<i>Bacillus licheniformis</i> SK4279 culture (10^9 CFU/mL)	0.1
<i>Bacillus subtilis</i> SK4282 culture (10^9 CFU/mL)	0.1
<i>Lactobacillus plantarum</i> SK4288 culture (10^9 CFU/mL)	0.1
Corn gluten meal	15.4
Glucose	4
Yeast culture	10
Angelica	0.1
Biotin	0.1
Vitamin A, D ₃ , E	4
Lysine	5
Methionine	5
Ginsenoside	0.1
Phytase	5
Total	100

tier battery cage with two birds in each cage (43 × 45 × 42 cm, length × width × height). The basal diet used in this experiment was formulated with nutrient levels that meet the requirements of the 2017 Korean Poultry Feeding Standard (Table 2). The appropriate amount of ESM was added to the basal diet, and mixed for 5 min using a feed mixer (DKM 350SU, Daekwang). After a 2-week adaptation period to the basal diet, the experimental diets were fed for 4 weeks of the experimental period. Food and drinking water were provided *ad libitum* throughout the entire experimental period. An automatic lighting controller was used to maintain a 16 h of light and 8 h dark period, and the temperature was maintained at (22 ± 3)°C. At the end of the experiment, hens were fasted for 18 h, prior to sampling. One bird per replicate was randomly selected and euthanized with carbon dioxide for evaluation of the blood, organ, and tibia characteristics.

Egg productivity

The number of eggs laid by birds in each replicate was recorded daily at 10 am, and expressed as the percentage of egg production. The hen-day egg production rate (EPR) is calculated by dividing

Table 2. Ingredients and chemical compositions of the basal diet

Items	Amount (%)
Ingredient (%)	
Corn	56.93
Dried distillers' grains with solubles	15.00
Soybean meal (crude protein, 45%)	5.54
Wheat gluten	4.12
Rapeseed meal	2.96
Sesame oil meal	2.04
Beef tallow	0.48
Limestone	11.49
Monocalcium phosphate	0.51
Methionine	0.16
Lysine sulfate	0.30
Threonine	0.02
NaCl	0.24
Choline chloride	0.02
Vitamin Premix ¹⁾	0.07
Mineral Premix ²⁾	0.12
Total	100.00
Calculated chemical composition	
Crude protein (%)	15.00
Crude fat (%)	3.82
Crude fiber (%)	2.73
Crude ash (%)	12.86
Calcium (%)	4.20
Available phosphorus (%)	0.53
AMEn (kcal/kg)	2,700

¹⁾Vitamin mixture provided the following nutrients per kg of diet: vitamin A, 20,000 IU; vitamin D₃, 4,600 IU; vitamin E, 40 mg; vitamin K₃, 4 mg; vitamin B₁, 3.6 mg; vitamin B₂, 8 mg; vitamin B₆, 5.8 mg; vitamin B₁₂, 0.04 mg.

²⁾Mineral mixture provided the following nutrients per kg of diet: Fe, 70 mg; Cu, 7.5 mg; Zn, 60 mg; Mn, 80 mg; I, 1 mg; Co, 0.1 mg; Se, 0.2 mg.

AMEn, nitrogen corrected apparent metabolizable energy.

the total number of eggs collected by the number of live hens daily in each replicate [27]. The total number of eggs produced in a day was weighed collectively for each replicate, and used to estimate the average egg weight (AEW). Daily egg mass was calculated by multiplying the EPR by the AEW. Feed intake (FI) was measured weekly once per replicate, weighing the amount of feed distributed and that of residual and scattered feed. The Feed conversion ratio (FCR) was calculated from the FI and daily egg mass [28].

Egg quality

Twenty-seven eggs per treatment (3 eggs per replicate) were randomly selected after each week, and analyzed for their quality on the same day of collection. Egg quality characteristics, including Haugh unit, albumen height, yolk color, eggshell weight, eggshell strength, and eggshell thickness were determined using an automatic egg analyzer (Digital egg tester DET6000, NABEL). The Haugh unit (HU) was calculated using the following equation: $HU = 100 \times \log [H + 7.57 - (1.7 \times W^{0.37})]$, where H is the albumen height (mm), and W is the egg weight (g) [29].

Blood sampling and analysis

At the end of the experiment, one bird (75 weeks of age) per group of replicates was randomly selected, and euthanized by CO₂ injection. After euthanasia, approximately 8 mL of blood was collected by cardiac puncture. The collected blood was kept refrigerated in a clot activator tube (CAT). Serum was separated from the blood sample in the CAT tube by centrifugation at 1,500 for 10 min using a centrifuge (HA-1000-3, Hanil Science Medical). The separated serum was stored at -20°C for observation of the biochemical properties. Serum concentrations of aspartate aminotransferase (AST), alanine aminotransferase (ALT), blood urea nitrogen (BUN), triglycerides (TG), lactate dehydrogenase (LDH), total cholesterol (TC), high-density lipoprotein (HDL; mg/dL), HDL (% total), low-density lipoprotein (LDL)+very low-density lipoprotein (VLDL), glucose, total protein (TP), albumin, creatinine, and calcium were determined by automated clinical chemistry analyzer (FUJI DRICHEM 7000i, Fujifilm). HDL (%) was expressed as the ratio of HDL to TC content, and LDL+VLDL was calculated by subtracting HDL from TC [30].

Organ weight and intestinal length

The weight of the visceral organs was determined from the weight of the liver and spleen. This was expressed as a weight ratio per 100 g of live body weight using an electronic balance (EL4002, Mettler Toledo). The intestine was divided into four sections (duodenum, jejunum, ileum, and cecum). The duodenum was measured from the pancreatic loop, the jejunum from the end of the pancreatic loop to the Meckel's diverticulum, the ileum from the Meckel's diverticulum to the ileocecal junction, and the cecum as the average of the right and left cecal lengths. The lengths of the four intestinal segments were measured, and expressed as the ratio of the length per 100 g of live body weight.

Tibia characteristics

At the end of the experiment, one bird (75 weeks old) per group of replicates was randomly selected to collect the left tibia, after the removal of non-bone tissues (fat, tendon, and muscle). The tibiae were individually sealed in plastic bags to minimize moisture loss, and stored at 4°C for one day. Tibia length and width were measured using a micrometer caliper, and the weight was recorded. Tibia strength was determined from a 3-point flexural test (ASAE Standards S459, 2001) using an Instron Universal Testing Machine (Model 3342) with a 50 kg load range and a crosshead speed of 200 mm/min; the tibia was supported on a 4 cm span [31].

Bone mineral density

Bone mineral density (BMD) of all the collected tibiae was analyzed by quantitative computed tomography (QCT) at the College of Veterinary Medicine, Konkuk University (Korea, Seoul). Three positions of each tibia including the neck (section of the mastoid arthrodesis), 1/3 of the proximal portion, and 2/3 of the distal portion, were scanned using a CT scanner (LightSpeed Plus, GE HealthCare).

Cecal microbiota

Three birds were randomly selected per treatment, and for each bird, approximately 1 g of the chicken ceca contents was collected, and quenched with liquid nitrogen. PCR conditions, DNA extraction, bioinformatics, and NGS sequencing analysis were performed according to a previously described method [32]. Briefly, a PowerSoil DNA Isolation Kit (Mobio Laboratories) was first used to isolate genomic DNA. The V3–V4 region of the bacterial 16S rRNA gene was then amplified using 341F and 785R primers. Sequencing was performed on the Illumina Miseq platform using the commercial service of Macrogen (Seoul, South Korea). Amplicon sequence variants (ASVs), Chao1, Shannon, and Gini–Simpson indices were checked to compare alpha diversity. Principal coordinate analysis (PCoA) and unweighted pair–group mean average (UPGMA) analysis based on the UniFrac distance matrix were used.

Statistical analysis

Data was analyzed in a completely randomized design with 4 treatments using the PROC GLM procedures of SAS 9.4 (SAS Institute). The replicate group (9 hens each) was the experimental unit for the analysis of performance data. Egg quality traits were statistically analyzed each week, using the number of eggs as the experimental unit. For blood parameters, organ weight, intestinal length, bone quality measurements, and cecal microbiota, the individual bird was used as the experimental unit. Significant differences between the treatments were determined using Duncan’s multiple range test at $p < 0.05$. Significance level $0.05 \leq p < 0.10$ was indicated as a trend. Data are presented as the least squares mean and SEM.

RESULTS

Egg productivity

Table 3 shows the effect of ESM on laying performance. Laying performance tended to increase in the ESM 0.2% group, compared to the control group ($p = 0.08$). ESM 0.1% supplementation had a higher egg weight, compared to the control ($p < 0.05$). ESM 0.1% and ESM 0.2% tended

Table 3. Supplementary effect of ESM on the laying performance in laying hens

Item	Treatment ¹⁾				SEM	p-value
	CON	ESM 0.1%	ESM 0.2%	ESM 0.4%		
Egg production ratio (%)	87.35	88.96	90.48	87.37	0.998	0.080
Average egg weight (g)	61.49 ^a	63.46 ^b	62.07 ^{ab}	62.43 ^{ab}	0.500	0.043
Daily egg mass (g/hen/day)	53.72 ^a	56.53 ^b	56.16 ^b	54.54 ^{ab}	0.822	0.051
Feed intake (g/hen/day)	131.97	136.84	131.85	135.39	2.299	0.310
FCR (g feed/g egg)	2.47	2.45	2.37	2.50	0.060	0.405

¹⁾CON, control, basal diet; ESM 0.1%, basal diet + 0.1% ESM; ESM 0.2%, basal diet + 0.2% ESM; ESM 0.4%, basal diet + 0.4% ESM.

^{a–b} Means with the different superscript in the same row differ significantly ($p < 0.05$).

ESM, combination of eggshell waste, *Schisandra chinensis* by-product, and multi-probiotics; FCR, feed conversion ratio.

to increase egg mass, compared to control ($p = 0.051$). There were no differences between the treatments in feed intake and FCR during the experimental period.

Egg quality

Table 4 presents the egg quality characteristics. The Laying hens fed ESM 0.4% group had the highest value for egg yolk color, while the ESM 0.2% group had the lowest value for egg yolk color ($p < 0.05$). There were no differences in the Haugh units, albumen height, and eggshell characteristics between the treatments during the experimental period.

Blood characteristics

Table 5 shows the effect of ESM on the blood biochemical parameters of the layers. The TC content was significantly higher in the ESM 0.4% group than in the control group ($p < 0.05$). There

Table 4. Supplementary effect of ESM on the egg quality in laying hens

Item	Treatment ¹⁾				SEM	p-value
	CON	ESM 0.1%	ESM 0.2%	ESM 0.4%		
Haugh units	87.35	88.44	88.90	87.77	0.453	0.635
Albumen height (mm)	7.71	8.00	8.00	7.88	0.074	0.484
Egg yolk color	8.22 ^{ab}	8.21 ^{ab}	8.05 ^b	8.29 ^a	0.027	0.012
Eggshell weight (g)	5.83	5.99	5.99	5.83	0.031	0.064
Eggshell breaking strength (kg/cm)	4.71	4.60	4.65	4.36	0.002	0.157
Eggshell thickness (mm)	0.44	0.43	0.43	0.42	0.017	0.077

¹⁾CON, control, basal diet; ESM 0.1%, basal diet + 0.1% ESM; ESM 0.2%, basal diet + 0.2% ESM; ESM 0.4%, basal diet + 0.4% ESM.

^{a-b} Means with the different superscript in the same row differ significantly ($p < 0.05$).

ESM, combination of eggshell waste, *Schisandra chinensis* by-product, and multi-probiotics.

Table 5. Supplementary effect of ESM on the blood characteristics in laying hens

Items	Treatment ¹⁾				SEM	p-value
	CON	ESM 0.1%	ESM 0.2%	ESM 0.4%		
AST (U/L)	206.88	202.56	183.00	194.33	16.464	0.759
ALT (U/L)	4.88	5.11	5.00	4.25	0.267	0.149
BUN (mg/dL)	2.16	2.00	2.14	2.04	0.087	0.512
TG (mg/dL)	472.75	566.56	651.00	951.89	170.996	0.254
LDH (mg/dL)	2325.25	2217.33	1735.56	2528.33	330.968	0.394
TC (mg/dL)	56.63 ^b	61.67 ^{ab}	65.00 ^{ab}	89.89 ^a	8.368	0.043
HDL (mg/dL)	22.38	23.56	23.89	27.89	2.272	0.371
HDL (%)	41.91	38.61	41.14	35.31	4.525	0.740
LDL+VLDL (mg/dL)	34.25	38.11	41.11	62.00	8.686	0.135
Glucose (mg/dL)	225.25	242.56	242.67	227.00	11.006	0.544
TP (g/dL)	4.83	4.56	4.74	5.01	0.203	0.472
Albumin (g/dL)	1.68	1.64	1.48	1.96	0.161	0.228
Creatinine (mg/dL)	0.21	0.24	0.21	0.24	0.026	0.665
Calcium (mg/dL)	13.28	15.59	15.20	17.53	1.021	0.059

¹⁾CON, control, basal diet; ESM 0.1%, basal diet + 0.1% ESM; ESM 0.2%, basal diet + 0.2% ESM; ESM 0.4%, basal diet + 0.4% ESM.

^{a-b} Means with the different superscript in the same row differ significantly ($p < 0.05$).

ESM, combination of eggshell waste, *Schisandra chinensis* by-product, and multi-probiotics; AST, aspartate aminotransferase; ALT, alanine aminotransferase; BUN, blood urea nitrogen; TG, triglyceride; LDH, lactate dehydrogenase; TC, total cholesterol; HDL, high-density lipoprotein; LDL, low-density lipoprotein; VLDL, very-low-density lipoprotein; TP, total protein.

was a tendency for ESM 0.4% to have a higher calcium content, compared to the control ($p = 0.059$).

Organ weight and intestinal length

Table 6 shows the organ weight and intestinal length of laying hens. There were no significant differences between treatments in the relative organ weight and intestinal length.

Tibia characteristics

Table 7 shows the tibia characteristics and BMD of laying hens. There was no significant effect of ESM treatment on the bone weight, length, width, and bone breaking strength. However, there were significant differences in the tibia BMD between treatments. The proximal tibia of laying hens fed ESM 0.4% had higher BMD, compared to the control group and the ESM 0.2% group ($p < 0.01$). In addition, the ESM 0.4% group had higher total tibia BMD, compared to the control group ($p < 0.05$).

Cecal microbiota

Table 8 shows the alpha diversity indices (ASVs, Chao1, Shannon, and Gini–Simpson) for the cecal

Table 6. Supplementary effect of ESM on the organ weight and intestinal length in laying hens

Item	Treatment ¹⁾				SEM	p-value
	CON	ESM 0.1%	ESM 0.2%	ESM 0.4%		
Visceral organ weight (g/100 g BW)						
Liver	1.78	1.73	1.88	1.82	0.043	0.250
Spleen	0.11	0.11	0.14	0.12	0.005	0.351
Intestinal length (cm/100 g BW)						
Duodenum	1.25	1.28	1.25	1.50	0.051	0.515
Jejunum	3.02	3.01	3.26	2.71	0.088	0.241
Ileum	2.66	2.72	2.91	2.62	0.077	0.408
Ceca	0.66	0.67	0.73	0.68	0.018	0.267

¹⁾CON, control, basal diet; ESM 0.1%, basal diet + 0.1% ESM; ESM 0.2%, basal diet + 0.2% ESM; ESM 0.4%, basal diet + 0.4% ESM.

ESM, combination of eggshell waste, *Schisandra chinensis* by-product, and multi-probiotics; BW, body weight.

Table 7. Supplementary effect of ESM on the tibia bone quality traits and BMD in laying hens

Item	Treatment ¹⁾				SEM	p-value
	CON	ESM 0.1%	ESM 0.2%	ESM 0.4%		
Bone weight (g)	13.91	13.91	14.30	13.37	0.029	0.525
Bone length (cm)	12.28	12.29	12.09	12.21	0.053	0.432
Bone width (mm)	8.76	8.91	8.92	8.87	0.066	0.524
Bone breaking strength	16.24	17.97	17.55	19.17	0.768	0.731
BMD (mg/cm ³)						
Tibia neck	264.53 ^c	342.32 ^{ab}	306.53 ^{bc}	385.33 ^a	28.250	0.004
1/3 tibia	340.09	412.08	396.27	435.06	31.447	0.171
2/3 tibia	313.86	375.12	353.92	391.05	31.965	0.380
Total	306.16 ^b	376.50 ^{ab}	352.24 ^{ab}	403.81 ^a	26.824	0.044

¹⁾CON, control, basal diet; ESM 0.1%, basal diet + 0.1% ESM; ESM 0.2%, basal diet + 0.2% ESM; ESM 0.4%, basal diet + 0.4% ESM.

^{a,b}Means with the different superscript in the same row differ significantly ($p < 0.05$).

ESM, combination of eggshell waste, *Schisandra chinensis* by-product, and multi-probiotics; BMD, bone mineral density.

Table 8. Supplementary effect of ESM on alpha diversity of cecum

Item	Treatment ¹⁾				SEM	p-value
	CON	ESM 0.1%	ESM 0.2%	ESM 0.4%		
ASVs	475.67 ^{ab}	427.00 ^a	442.00 ^a	499.00 ^b	14.48	0.030
Chao 1	483.50 ^{ab}	428.05 ^c	447.78 ^{ac}	507.79 ^b	15.38	0.025
Shannon	7.15	7.02	6.93	7.29	0.086	0.079
Gini-Simpson	0.981	0.982	0.979	0.985	0.002	0.596

¹⁾CON, control, basal diet; ESM 0.1%, basal diet + 0.1% ESM; ESM 0.2%, basal diet + 0.2% ESM; ESM 0.4%, basal diet + 0.4% ESM.

^{a,b}Means with the different superscript in the same row differ significantly ($p < 0.05$).

ESM, combination of eggshell waste, *Schisandra chinensis* by-product, and multi-probiotics; ASVs, amplicon sequence variants.

microbiota of laying hens. Supplementation with ESM 0.4% showed higher alpha diversity (ASVs and Chao1) than ESM 0.1% and ESM 0.2% ($p < 0.05$), but ESM 0.4% was not significantly different from the control. Shannon and Gini-Simpson were not significantly different between the groups. Fig. 1 shows the result of the PCoA (beta diversity) and phylogenetic tree analysis representing the similarity of the microbial community. The results demonstrated that microbial communities in the ESM supplementation exhibited distinct clustering patterns compared to the control. Specifically, the ESM group samples were clearly differentiated by their microbial composition, indicating that the feed additive had a significant effect on the gut microbiota. Fig. 2 shows the cecal microbiota. *Firmicutes* (71.7%) was the most abundant phylum in the cecal microbiome, followed by *Bacteroidetes* (23.8%) as the second most abundant phylum (Fig. 2A). At the family level, the abundance of *Bacteroidaceae* was significantly higher in the control groups than in the other ESM groups ($p < 0.05$). *Lactobacillaceae* was significantly more abundant in the ESM groups than in the control group. ESM 0.1 showed the highest abundance of *Lactobacillaceae* (25.63%) ($p < 0.01$).

DISCUSSION

Egg productivity

Interest in environment-friendly feed additives is increasing, and there is considerable research into the effects of probiotics and agricultural by-products on laying hen productivity and health. In this




Fig. 1. Beta-diversity analysis of cecal microbiota (A). Phylogenetic tree (B). CON, control, basal diet; ESM 0.1%, basal diet + 0.1% ESM; ESM 0.2%, basal diet + 0.2% ESM; ESM 0.4%, basal diet + 0.4% ESM.




Fig. 2. Relative abundances of the cecal microbiota at the phylum level (A). Relative abundances of the cecal microbiota at the family level (B). Abundances at the family level of *Bacteroidaceae* (C) and *Lactobacillaceae* (D). CON, control, basal diet; ESM 0.1%, basal diet + 0.1% ESM; ESM 0.2%, basal diet + 0.2% ESM; ESM 0.4%, basal diet + 0.4% ESM. ^{a,b}Means with the different superscript in the column differ significantly ($p < 0.05$, $n = 3$).

study, the four-week duration of the feeding trial was conducted to evaluate the immediate effects of the feed additive on laying performance in aged laying hens before slaughter. This period was designed to capture short-term impacts on productivity, egg quality, physiological changes, and economic benefits. Aged hens were selected to evaluate their potential for sustained productivity, providing insights into the practical and economic benefits of using feed additives to enhance performance in older birds.

In this study, supplementation with ESM improved egg weight and egg mass during the experimental period. Lee et al. [33] reported that supplementation with eggshell coarse (ESC) improved the egg weight, egg mass, and FCR, compared to other calcium source treatments. Similarly, eggshell meal supplementation increased the average egg weight, egg mass, and FCR, compared to bone meal treatment, or the inclusion of eggshell meal and bone meal [34]. The main composition of eggshell is calcium. Eggshells also have high protein concentrations, due to the egg membranes. Appropriate calcium supplementation can produce stronger eggshells and help to reduce the production of soft-shelled or shell-less eggs, thus improving the laying performance and FCR [35,36].

Multi-probiotics are known to contain bioactive compounds and secondary metabolites, and have been used as a potential feed additive [12,13,37]. Many studies have reported that supplementation with multi-probiotics improved egg productivity, egg weight, egg mass, and FCR. Ma et al. [23] reported that either 1% *Ligustrum lucidum* or *Schisandra chinensis* supplementation

improved egg production and FCR to laying hens (57 weeks of age). In contrast, body weight and FCR in layer chicks were not affected by either 1% *Ligustrum lucidum* or *S. chinensis* treatment [38]. Some studies reported that *S. chinensis* and probiotics made no significant difference in laying performance [39,40]. The discrepancy in outcomes may be attributed to the impact of various factors on productivity, including age, diet, fermentation method, and farm environment.

The findings of this study indicate that ESM 0.1% and 0.2% are associated with an increase in egg mass. The observed increase in egg mass associated with ESM may be attributed to various physiological mechanisms. The bioactive compounds in ESM may improve gut health or enhance nutrient absorption, thereby increasing the effective use of nutrients for egg formation. However, it is important to acknowledge some limitations of this study. The research was conducted within a specific farm environment, which may limit the generalizability of the results. Further studies in diverse farm settings are needed to confirm these findings, and the long-term effects of ESM supplementation at different stages of the laying cycle should also be investigated. Such additional research would provide a more comprehensive understanding of the effects of ESM on egg production.

Egg quality

Eggshell powder contains protein and minerals (Ca, Fe, Mg, Zn, Cu, and Se), and the balanced mineral content of the diet can influence egg quality. Among the minerals, calcium plays an important role in eggshell formation and increases eggshell strength [10,41,42]. Several studies have shown that eggshell powder with high calcium content can improve eggshell quality. Lee et al. [33] showed that supplementation with eggshell coarse improved egg weight among dietary treatments. The oyster shell or eggshell coarse group had a higher albumen height than the cockle shell group, and egg yolk color was the highest in laying hens fed eggshell fine. Kismiati et al. [43] observed that a 7.5% eggshell flour group or mixture of 5% eggshell flour and 2.5% limestone increased eggshell weight. This suggests that increasing the concentration of ESM improves egg quality.

In contrast to previous studies, this study found no significant differences between the ESM groups. This lack of effect might be related to several factors, including the possibility that the amounts of eggshell powder in the ESM treatments were insufficient, or that the trial duration was not long enough to observe measurable changes in egg quality. Additionally, the specific physical and chemical properties of the ESM used in our study may differ from those in previous studies, potentially influencing the outcomes.

Blood characteristics

The general health of laying hens could be assessed by blood analysis. ALT, AST, and LDH are commonly used biomarkers of liver damage in laying hens [44]. In this study, ESM did not affect the levels of ALT, AST, and LDH in laying hens, suggesting that the ESM diet did not adversely affect liver health.

Albumin is a protein produced by the liver, and albumin concentrations can indicate liver and kidney function. BUN and creatinine in blood tests can indicate kidney function and health [45,46]. These are nitrogenous end products of metabolism. A high ratio of BUN to creatinine leads to reduced filtration by the kidneys, due to reduced blood flow to the kidneys [47]. In this study, albumin, total protein, BUN, and creatinine were not affected by ESM supplementation, suggesting that the ESM treatments did not have a detrimental effect on protein metabolism.

The range of serum TC levels can vary depending on factors such as age, diet, and genetics. Several studies reported serum total cholesterol ranges of (107.29–116.67) mg/dL [26], (103.8–157.8) mg/dL [33], and (157.81–170.53) mg/dL [24] in laying hens, although there

was no significant difference between the treatments. In this study, although total cholesterol and LDL+VLDL levels were lower than in other studies, these may not represent a general standard. Therefore, in the present study, no hens died during the experiment, suggesting that the ESM diet was non-toxic, metabolically stable, and had no adverse effects on the health of laying hens.

Organ weight and length

Changes in the structure and size of organs are related to their development, including gut immunity and digestive function, and can be used to assess their health status [48]. In general, as the size of an organ increases, the energy required to maintain it increases, reducing the energy available for productivity [49]. In addition, the spleen is small, and is an important lymphoid organ in the immune system. However, infections, liver and blood diseases, and a rapid immune response can lead to an enlarged spleen [50]. The liver is a large organ responsible for toxin removal, digestion, metabolism, and immunity. An increase in liver size is a sign of health problems, such as fatty liver disease, hepatitis, and cancer [51,52].

Kim et al. [24] observed that 2% *S. chinensis* supplementation showed the lowest liver weight and abdominal fat, but *S. chinensis* treatments had no effect on spleen weight. Supplementation of whole hatchery waste meal including eggshell showed no significant difference in abdominal fat and internal organs (liver, lung, heart, and gizzard) in broiler [53]. This indicated that eggshell powder might not be affect the organ characteristics.

In contrast, our study revealed that ESM supplementation did not significantly impact organ characteristics. These findings suggest that ESM supplementation does not negatively affect organ characteristics, indicating its safety with respect to organ health. Additionally, it is possible that the bioavailability or the specific components of ESM were insufficient to elicit measurable changes in organ characteristics under the conditions tested. Further research could explore different dosages or durations of ESM to determine whether any conditions might reveal potential benefits or effects on organ characteristics.

Tibia characteristics

Recently, there has been increasing interest in improving BMD and bone quality in laying hens. The bones of laying hens play an important role in mobility, productivity, and overall health. Calcium is an essential component of bone, and this influences bone quality and breaking strength [54,55].

Several studies have reported the effect of eggshell powder supplementation on tibia bone characteristics and BMD. Lee et al. [33] observed that supplementation with oyster shell or coarse eggshell particles showed higher BMD in the proximal, distal, and total tibia. Kismati et al. [43] found that 5% eggshell flour supplementation had the highest calcium rate in the tibia, while eggshell flour had no effect on the tibia length and weight. Similar to previous studies, this study showed that when ESM 0.4% was fed to laying hens, total and tibial neck BMD were improved. Eggshell powder is known to have a high calcium content, so supplementation with eggshell may have an effect on BMD increase and bone quality in laying hens.

Cecal microbiota

The gut microbiota plays a critical role in maintaining overall health and influencing digestive system health, immunity, and resistance to pathogens. In this study, analysis of alpha diversity metrics, including ASVs and the Chao1 index, showed that supplementation with ESM 0.4% increased microbial diversity in the cecum compared to ESM 0.1% and ESM 0.2%, although not significantly compared to control. ASVs provide high-resolution insights into the composition of microbial communities, and highlight the diversity and possible functional roles of the microbiota

[56]. Similarly, the increased Chao1 index indicates richer species diversity [57], suggesting a more complex and potentially resilient ecosystem under ESM 0.4% treatment.

PCoA is used to determine the beta diversity analysis. PCoA plays a critical role in assessing variation in species composition across samples, and provides valuable insight into the effects of dietary interventions on microbial community structure [58]. In this study, PCoA showed that ESM groups had a more similar composition of cecal microbiota, compared to the control. This suggests that ESM groups influence the composition of the gut microbiome.

Furthermore, *Firmicutes* and *Bacteroidetes* were the most abundant strains in the cecum of laying hens in the study presented. ESM supplementation increased the relative abundance of the *Lactobacillaceae* family of *Firmicutes*, and decreased the *Bacteroidaceae* family of *Bacteroidetes*. This suggested that ESM supplementation increased the *Lactobacillaceae* family, while decreasing the composition of *Bacteroidaceae* in the cecum. Ren et al. [16] reported that combinations of phytobiotics and probiotics increased the lactobacilli and decreased ESBL-producing *E. coli* in the gut of young broiler chickens. These lactic acid bacteria are known to have many beneficial effects, including stimulating the immune system, producing lactic acid, inhibiting the growth of pathogens, and contributing to the overall health of laying hens [58–60]. Therefore, it is proposed that the ESM intervention alters the structure of the cecal microbiota and increases its diversity in the gut. ESM supplementation may be a promising strategy to enhance gut health by improving the balance of gut microbiota.

An improvement in gut microbiota is closely linked to enhanced immunity, digestive efficiency, and overall poultry productivity [61,62]. A growing body of evidence indicates that modulation of the gut microbiota can exert a beneficial influence on a number of key aspects of poultry production, including growth, feed efficiency, immune function, and disease resistance [63]. [64] reported that fermented plant product interventions can improve productivity, egg mass, Haugh unit, gut health, and alter the cecal microbial community in laying hens. Therefore. These findings emphasize the crucial role of gut microbiota in supporting poultry health and productivity.

CONCLUSION

Supplementation with ESM resulted in significant increases in egg weight, egg mass, tibial BMD, and cecal microbiota diversity. In addition, ESM did not affect blood characteristics or visceral organ properties, suggesting that it does not adversely affect the overall health of laying hens. Notably, ESM has not previously been studied as a feed additive for poultry, which may highlight its novel application. The observed improvements in egg weight, bone health, and microbial diversity underscore the potential value of ESM as a beneficial feed additive to improve egg performance and gut health in laying hens.

REFERENCES

1. Barton MD. Antibiotic use in animal feed and its impact on human health. *Nutr Res Rev.* 2000;13:279-99. <https://doi.org/10.1079/095442200108729106>
2. Chowdhury R, Haque MN, Islam KMS, Khaleduzzaman ABM. A review on antibiotics in an animal feed. *Bangl J Anim Sci.* 2009;38:22-32. <https://doi.org/10.3329/bjas.v38i1-2.9909>
3. Wegener HC. Antibiotics in animal feed and their role in resistance development. *Curr Opin Microbiol.* 2003;6:439-45. <https://doi.org/10.1016/j.mib.2003.09.009>
4. Han D, Ren T, Yang Y, Li Z, Du X, Zhang C, et al. Application and substitution of antibiotics in animal feeding. *Med Weter.* 2024;80:5-11. <https://doi.org/10.21521/mw.6830>

5. Arsène MMJ, Davares AKL, Andreevna SL, Vladimirovich EA, Carime BZ, Marouf R, et al. The use of probiotics in animal feeding for safe production and as potential alternatives to antibiotics. *Vet World*. 2021;14:319-28. <https://doi.org/10.14202/vetworld.2021.319-328>
6. Hidayat R, Yunianto VD, Sukanto B, Sugiharto S. Effect of dietary supplementation of probiotic, phytobiotics or their combination on performance, blood indices and jejunal morphology of laying hens during post peak production. *Online J Anim Feed Res*. 2021;11:8-12. <https://doi.org/10.51227/ojafr.2021.2>
7. Hussein EOS, Ahmed SH, Abudabos AM, Suliman GM, Abd El-Hack ME, Swelum AA, et al. Ameliorative effects of antibiotic-, probiotic- and phytobiotic-supplemented diets on the performance, intestinal health, carcass traits, and meat quality of *Clostridium perfringens*-infected broilers. *Animals*. 2020;10:669. <https://doi.org/10.3390/ani10040669>
8. Das S, Mohanty PK, Mallik BK. Agricultural and pharmaceutical applications of eggshells: a comprehensive review of eggshell waste value-added products. *J Pharm Negat Results*. 2022;13:3979-84.
9. Sivakumar A, Srividhya S, Prakash R, Padma S. Properties of biocomposites from waste eggshell as fillers. *Glob Nest J*. 2023;25:109-15. <https://doi.org/10.30955/gnj.005095>
10. Waheed M, Yousaf M, Shehzad A, Inam-Ur-Raheem M, Khan MKI, Khan MR, et al. Channelling eggshell waste to valuable and utilizable products: a comprehensive review. *Trends Food Sci Technol*. 2020;106:78-90. <https://doi.org/10.1016/j.tifs.2020.10.009>
11. Osonwa UE, Okoye CS, Abali SO, Uwaezuoke OJ, Adikwu MU. Egg shell powder as a potential direct compression excipient in tablet formulation. *West Afr J Pharm*. 2017;28:107-18. <https://doi.org/10.60787/wapcp-28-1-144>
12. Wang Y, Sun J, Zhong H, Li N, Xu H, Zhu Q, et al. Effect of probiotics on the meat flavour and gut microbiota of chicken. *Sci Rep*. 2017;7:6400. <https://doi.org/10.1038/s41598-017-06677-z>
13. Zhang Q, Cho S, Kibria S, Kim IH. Dietary *Bacillus subtilis*- and *Clostridium butyricum*-based probiotics supplement improves growth and meat quality, and alters microbiota in the excreta of broiler chickens. *Can J Anim Sci*. 2023;104:200-6. <https://doi.org/10.1139/cjas-2023-0076>
14. Perić L, Milošević N, Žikić D, Bjedov S, Cvetković D, Markov S, et al. Effects of probiotic and phytogenic products on performance, gut morphology and cecal microflora of broiler chickens. *Arch Anim Breed*. 2010;53:350-9. <https://doi.org/10.5194/aab-53-350-2010>
15. Lee JS, Kim MJ, Park SH, Lee SB, Wang T, Jung US, et al. Effects of dietary mixture of garlic (*Allium sativum*), coriander (*Coriandrum sativum*) and probiotics on immune responses and caecal counts in young laying hens. *J Anim Physiol Anim Nutr*. 2017;101:e122-32. <https://doi.org/10.1111/jpn.12573>
16. Ren H, Vahjen W, Dadi T, Saliu EM, Boroojeni FG, Zentek J. Synergistic effects of probiotics and phytobiotics on the intestinal microbiota in young broiler chicken. *Microorganisms*. 2019;7:684. <https://doi.org/10.3390/microorganisms7120684>
17. Lee AR, Niu KM, Lee WD, Kothari D, Kim SK. Comparison of the dietary supplementation of *Lactobacillus plantarum*, and fermented and non-fermented *Artemisia annua* on the performance, egg quality, serum cholesterol, and egg yolk-oxidative stability during storage in laying hens. *Braz J Poult Sci*. 2019;21:eRBCA-2018-0903. <https://doi.org/10.1590/1806-9061-2018-0903>
18. Shahrajabian MH, Cheng Q, Sun W. Application of herbal plants in organic poultry nutrition and production. *Curr Nutr Food Sci*. 2022;18:629-41. <https://doi.org/10.2174/1573401318666220308155156>

19. Shehata AA, Yalçın S, Latorre JD, Basiouni S, Attia YA, Abd El-Wahab A, et al. Probiotics, prebiotics, and phytogetic substances for optimizing gut health in poultry. *Microorganisms*. 2022;10:395. <https://doi.org/10.3390/microorganisms10020395>
20. Yang S, Yuan C. *Schisandra chinensis*: a comprehensive review on its phytochemicals and biological activities. *Arab J Chem*. 2021;14:103310. <https://doi.org/10.1016/j.arabjc.2021.103310>
21. Ji R, Wang Z, Kuang H. Extraction, purification, structural characterization, and biological activity of polysaccharides from *Schisandra chinensis*: a review. *Int J Biol Macromol*. 2024;271:132590. <https://doi.org/10.1016/j.ijbiomac.2024.132590>
22. Kopustinskiene DM, Bernatoniene J. Antioxidant effects of *Schisandra chinensis* fruits and their active constituents. *Antioxidants*. 2021;10:620. <https://doi.org/10.3390/antiox10040620>
23. Ma D, Shan A, Chen Z, Du J, Song K, Li J, et al. Effect of *Ligustrum lucidum* and *Schisandra chinensis* on the egg production, antioxidant status and immunity of laying hens during heat stress. *Arch Anim Nutr*. 2005;59:439-47. <https://doi.org/10.1080/17450390500353499>
24. Kim YJ, Chung TH, Choi IH. Influence of supplemental *Schisandra chinensis* powder on growth performance, serum cholesterol, and meat quality of broilers. *Acta Agric Scand A Anim Sci*. 2013;63:175-82. <https://doi.org/10.1080/09064702.2013.861861>
25. Kang HM, Park EJ, Park SY, Hwang DY, Lee JC, Kim M, et al. Production of lignan-rich eggs as functional food by supplementing *Schisandra chinensis* by-product in laying hens. *J Life Sci*. 2024;34:18-27. <https://doi.org/10.5352/JLS.2024.34.1.18>
26. Moon SG, Lee SK, Lee WD, Niu KM, Hwang WU, Oh JS, et al. Effect of dietary supplementation of a phytogetic blend containing *Schisandra chinensis*, *Pinus densiflora*, and *Allium tuberosum* on productivity, egg quality, and health parameters in laying hens. *Anim Biosci*. 2021;34:285-94. <https://doi.org/10.5713/ajas.20.0552>
27. Xu CL, Ji C, Ma Q, Hao K, Jin ZY, Li K. Effects of a dried *Bacillus subtilis* culture on egg quality. *Poult Sci*. 2006;85:364-8. <https://doi.org/10.1093/ps/85.2.364>
28. Bonekamp RPRT, Lemme A, Wijtten PJA, Sparla JKWM. Effects of amino acids on egg number and egg mass of brown (heavy breed) and white (light breed) laying hens. *Poult Sci*. 2010;89:522-9. <https://doi.org/10.3382/ps.2009-00342>
29. Haugh RR. The Haugh unit for measuring egg quality. *United States Egg Poult Mag*. 1937; 43:552-5.
30. Aryal M, Poudel A, Satyal B, Gyawali P, Pokheral BR, Raut BK, et al. Evaluation of non-HDL-c and total cholesterol: HDL-c ratio as cumulative marker of cardiovascular risk in diabetes mellitus. *Kathmandu Univ Med J*. 2010;8:398-404. <https://doi.org/10.3126/kumj.v8i4.6239>
31. Ndazigaruye G, Kim DH, Kang CW, Kang KR, Joo YJ, Lee SR, et al. Effects of low-protein diets and exogenous protease on growth performance, carcass traits, intestinal morphology, cecal volatile fatty acids and serum parameters in broilers. *Animals*. 2019;9:226. <https://doi.org/10.3390/ani9050226>
32. Niu KM, Khosravi S, Kothari D, Lee WD, Lim JM, Lee BJ, et al. Effects of dietary multi-strain probiotics supplementation in a low fishmeal diet on growth performance, nutrient utilization, proximate composition, immune parameters, and gut microbiota of juvenile olive flounder (*Paralichthys olivaceus*). *Fish Shellfish Immunol*. 2019;93:258-68. <https://doi.org/10.1016/j.fsi.2019.07.056>
33. Lee WD, Kothari D, Niu KM, Lim JM, Park DH, Ko J, et al. Superiority of coarse eggshell as a calcium source over limestone, cockle shell, oyster shell, and fine eggshell in old laying hens. *Sci Rep*. 2021;11:13225. <https://doi.org/10.1038/s41598-021-92589-y>
34. Okpanachi U, Yusuf KA, Ikubaje MK, Okpanachi GCA. Effects of egg shell meal on the

- performance and haematology of layers and their egg quality. *Afr J Sci Technol Innov Dev*. 2021;13:89-96. <https://doi.org/10.1080/20421338.2020.1838111>
35. Olgun O, Yildiz AÖ, Cufadar Y. The effects of eggshell and oyster shell supplemental as calcium sources on performance, eggshell quality and mineral excretion in laying hens. *Indian J Anim Res*. 2015;49:205-9. <https://doi.org/10.5958/0976-0555.2015.00056.4>
 36. Wang J, Wang W, Qi G, Cui C, Wu S, Zhang H, et al. Effects of dietary *Bacillus subtilis* supplementation and calcium levels on performance and eggshell quality of laying hens in the late phase of production. *Poult Sci*. 2021;100:100970. <https://doi.org/10.1016/j.psj.2020.12.067>
 37. Kim D, Min Y, Yang J, Heo Y, Kim M, Hur CG, et al. Multi-probiotic *Lactobacillus* supplementation improves liver function and reduces cholesterol levels in Jeju native pigs. *Animals*. 2021;11:2309. <https://doi.org/10.3390/ani11082309>
 38. Ma D, Liu Y, Liu S, Li Q, Shan A. Influence of *Ligustrum lucidum* and *Schisandra chinensis* fruits on antioxidative metabolism and immunological parameters of layer chicks. *Asian-Australas J Anim Sci*. 2007;20:1438-43. <https://doi.org/10.5713/ajas.2007.1438>
 39. Baghban-Kanani P, Hosseintabar-Ghasemabad B, Azimi-Youvalari S, Seidavi A, Ragni M, Laudadio V, et al. Effects of using *Artemisia annua* leaves, probiotic blend, and organic acids on performance, egg quality, blood biochemistry, and antioxidant status of laying hens. *J Poult Sci*. 2019;56:120-7. <https://doi.org/10.2141/jpsa.0180050>
 40. Afsari M, Mohebbifar A, Torki M. Effects of dietary inclusion of olive pulp supplemented with probiotics on productive performance, egg quality and blood parameters of laying hens. *Ann Res Rev Biol*. 2013;4:198-211. <https://doi.org/10.9734/ARRB/2014/5212>
 41. King'ori A. A review of the uses of poultry eggshells and shell membranes. *Int J Poult Sci*. 2011;10:908-12. <https://doi.org/10.3923/ijps.2011.908.912>
 42. Hamideh F, Akbar A. Application of eggshell wastes as valuable and utilizable products: a review. *Res Agric Eng*. 2018;64:104-14. <https://doi.org/10.17221/6/2017-RAE>
 43. Kismiati S, Yuwanta T, Zuprizal Z, Supadmo S, Atmomarsono U. Calcium deposition in egg due to substitution of limestone by eggshell flour in feed of laying hens. *J Indones Trop Anim Agric*. 2018;43:257-64. <https://doi.org/10.14710/jitaa.43.3.257-264>
 44. Ozer J, Ratner M, Shaw M, Bailey W, Schomaker S. The current state of serum biomarkers of hepatotoxicity. *Toxicology*. 2008;245:194-205. <https://doi.org/10.1016/j.tox.2007.11.021>
 45. Hosten AO. BUN and creatinine. In: Walker HK, Hall WD, Hurst JW, editors. *Clinical methods: the history, physical, and laboratory examinations*. Butterworths; 1990.
 46. Salazar JH. Overview of urea and creatinine. *Lab Med*. 2014;45:e19-20. <https://doi.org/10.1309/LM920SBNZPJRGUT>
 47. Uchino S, Bellomo R, Goldsmith D. The meaning of the blood urea nitrogen/creatinine ratio in acute kidney injury. *Clin Kidney J*. 2012;5:187-91. <https://doi.org/10.1093/ckj/sfs013>
 48. Yegani M, Korver DR. Factors affecting intestinal health in poultry. *Poult Sci*. 2008;87:2052-63. <https://doi.org/10.3382/ps.2008-00091>
 49. Celi P, Cowieson AJ, Fru-Nji F, Steinert RE, Kluenter AM, Verlhac V. Gastrointestinal functionality in animal nutrition and health: new opportunities for sustainable animal production. *Anim Feed Sci Technol*. 2017;234:88-100. <https://doi.org/10.1016/j.anifeedsci.2017.09.012>
 50. William BM, Corazza GR. Hyposplenism: a comprehensive review. Part I: basic concepts and causes. *Hematology*. 2007;12:1-13. <https://doi.org/10.1080/10245330600938422>
 51. Whitehead CC. Nutritional and metabolic aspects of fatty liver disease in poultry. *Vet Q*. 1979;1:150-7. <https://doi.org/10.1080/01652176.1979.9693738>
 52. Julian RJ. Production and growth related disorders and other metabolic diseases of poultry – a

- review. *Vet J.* 2005;169:350-69. <https://doi.org/10.1016/j.tvjl.2004.04.015>
53. Abiola SS, Radebe NE, Westhuizen C, Umesiobi DO. Whole hatchery waste meal as alternative protein and calcium sources in broiler diets. *Arch Zootec.* 2012;61:229-34. <https://doi.org/10.4321/S0004-05922012000200007>
 54. Olgun O, Aygun A. Nutritional factors affecting the breaking strength of bone in laying hens. *Worlds Poult Sci J.* 2016;72:821-32. <https://doi.org/10.1017/S0043933916000696>
 55. Zhao SC, Teng XQ, Xu DL, Chi X, Ge M, Xu SW. Influences of low level of dietary calcium on bone characters in laying hens. *Poult Sci.* 2020;99:7084-91. <https://doi.org/10.1016/j.psj.2020.08.057>
 56. Tom WA, Judy JV, Kononoff PJ, Fernando SC. Influence of empirically derived filtering parameters, amplicon sequence variant, and operational taxonomic unit pipelines on assessing rumen microbial diversity. *J Dairy Sci.* 2024;107:9209-34. <https://doi.org/10.3168/jds.2023-24479>
 57. Kers JG, Saccenti E. The power of microbiome studies: some considerations on which alpha and beta metrics to use and how to report results. *Front Microbiol.* 2022;12:796025. <https://doi.org/10.3389/fmicb.2021.796025>
 58. Wang W, Jia H, Zhang H, Wang J, Lv H, Wu S, et al. Supplemental plant extracts from flos lonicerae in combination with baikal skullcap attenuate intestinal disruption and modulate gut microbiota in laying hens challenged by *Salmonella pullorum*. *Front Microbiol.* 2019;10:1681. <https://doi.org/10.3389/fmicb.2019.01681>
 59. Khan S, Moore RJ, Stanley D, Chousalkar KK. The gut microbiota of laying hens and its manipulation with prebiotics and probiotics to enhance gut health and food safety. *Appl Environ Microbiol.* 2020;86:e00600-20. <https://doi.org/10.1128/AEM.00600-20>
 60. Lv J, Guo L, Chen B, Hao K, Ma H, Liu Y, et al. Effects of different probiotic fermented feeds on production performance and intestinal health of laying hens. *Poult Sci.* 2022;101:101570. <https://doi.org/10.1016/j.psj.2021.101570>
 61. Kogut MH. The effect of microbiome modulation on the intestinal health of poultry. *Anim Feed Sci Technol.* 2019;250:32-40. <https://doi.org/10.1016/j.anifeedsci.2018.10.008>
 62. Jha R, Das R, Oak S, Mishra P. Probiotics (direct-fed microbials) in poultry nutrition and their effects on nutrient utilization, growth and laying performance, and gut health: a systematic review. *Animals.* 2020;10:1863. <https://doi.org/10.3390/ani10101863>
 63. Yadav S, Jha R. Strategies to modulate the intestinal microbiota and their effects on nutrient utilization, performance, and health of poultry. *J Anim Sci Biotechnol.* 2019;10:2. <https://doi.org/10.1186/s40104-018-0310-9>
 64. Tian Y, Li G, Zhang S, Zeng T, Chen L, Tao Z, et al. Dietary supplementation with fermented plant product modulates production performance, egg quality, intestinal mucosal barrier, and cecal microbiota in laying hens. *Front Microbiol.* 2022;13:955115. <https://doi.org/10.3389/fmicb.2022.955115>