

Effects of phytobiotics on intestinal barrier function and gut microbiome in weaned piglets challenged with enterotoxigenic *Escherichia coli*

Gi Beom Keum^{1#}, Hyunok Doo^{1#}, Jinok Kwak^{1#}, Xingmin Sun², Jinho Cho³, Hyeun Bum Kim^{1*}



¹Department of Animal Biotechnology, Dankook University, Cheonan 31116, Korea

²Department of Molecular Medicine, Morsani College of Medicine, University of South Florida, Tampa, FL 33612, USA

³Department of Animal Science, Chungbuk National University, Cheongju 28644, Korea

Received: Apr 24, 2025

Revised: May 8, 2025

Accepted: May 8, 2025

#These authors contributed equally to this work.

*Corresponding author

Hyeun Bum Kim

Department of Animal Biotechnology,
Dankook University, Cheonan 31116,
Korea.

Tel: +82-41-550-3653

E-mail: hbkim@dankook.ac.kr

Copyright © 2026 Korean Society of Animal Science and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Gi Beom Keum

<https://orcid.org/0000-0001-6006-9577>

Hyunok Doo

<https://orcid.org/0000-0003-4329-4128>

Jinok Kwak

<https://orcid.org/0000-0003-1217-3569>

Xingmin Sun

<https://orcid.org/0000-0002-8900-9161>

Jinho Cho

<https://orcid.org/0000-0001-7151-0778>

Hyeun Bum Kim

<https://orcid.org/0000-0003-1366-6090>

Abstract

Weaned piglets are highly susceptible to infections caused by enterotoxigenic *Escherichia coli* (ETEC), and phytobiotic supplementation has been explored as a potential strategy to prevent or mitigate such infections during the weaning period. However, comprehensive studies on the specific effects of phytobiotics on pig gut health and microbiota composition remains limited. Therefore, this study aimed to investigate the effects of various phytogenic feed additives (PFAs) on intestinal barrier function and gut microbiota composition in weaned piglets challenged with ETEC. A total of 63 weaned piglets ([Yorkshire × Landrace] × Duroc), 28 days old with an initial body weight of 8.03 ± 0.43 kg, were used in a 21-day trial. Piglets were randomly assigned to one of seven treatment groups: NC, negative control (basal diets without *E. coli* challenge); PC, positive control (basal diets + *E. coli* challenge); T1, PC + 0.04 % P1; T2, PC + 0.01 % P2; T3, PC + 0.10 % P3; T4, PC + 0.04 % P4; T5, PC + 0.10 % P5. The five phytobiotic materials were as follows: a bitter citrus extract rich in flavonoids (P1), a microencapsulated blend of thymol and carvacrol (P2), a composite of P1 and P2 (P3), a grape polyphenol-based mixture containing green tea and hops (P4), and a fenugreek seed powder rich in saponins (P5). Piglets in challenged groups were orally inoculated with ETEC at a concentration of 1.2×10^{10} CFU/mL from days 8 to 10. The results showed that PFA supplementation upregulated the expression of tight junction and mucin-related genes including ZO-1 and MUC3. Based on 16S rRNA gene sequencing, distinct microbial shifts were observed depending on the specific PFA composition. The relative abundances of beneficial genera such as *Lactobacillus*, *Olsenella*, and *Collinsella* increased following PFA supplementation. These genera are known to be associated with short-chain fatty acid production, improved gut health, and increased resistance to pathogens. Additionally, increases in *Acinetobacter* and *Medioterraneibacter*, which may be involved in the metabolism of plant-derived compounds, were observed. Collectively, our results suggest that PFAs may enhance gut health and contribute to the stabilization of the intestinal microbiota in weaned piglets.

Keywords: Piglet, Gut microbiome, Phytobiotics, Gut health

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

The present research was supported by the research fund of Dankook University in 2025.

Acknowledgements

Not applicable.

Availability of data and material

The datasets analyzed for this study can be found in the National Center for Biotechnology Information (NCBI) database under Sequence Read Archive (SRA) with the BioProject accession number PRJNA1195693.

Authors' contributions

Conceptualization: Keum GB, Cho J, Kim HB. Data curation: Doo H, Kwak J. Formal analysis: Keum GB. Validation: Sun X. Investigation: Keum GB, Doo H, Kwak J. Writing - original draft: Keum GB, Doo H, Kwak J. Writing - review & editing: Keum GB, Doo H, Kwak J, Sun X, Cho J, Kim HB.

Ethics approval and consent to participate

The protocol for this study was reviewed and approved by the Institutional Animal Care and Use Committee of Chungbuk National University, Cheongju, Korea (approval no. CBNUA-1618-21-02).

INTRODUCTION

The weaning process represents one of the most significant stressors in swine production, particularly affecting piglets at 3–4 weeks of age, when their gastrointestinal and immune systems are still immature. During this critical transition period, weaned piglets undergo substantial physiological changes including disruptions in gut integrity, alterations in villus architecture and mucosal permeability, and shifts in gut microbiota composition [1]. These changes primarily driven by dietary transitions increase the susceptibility of piglets to infectious diseases [2,3]. *Escherichia coli* is considered one of the major enteric pathogens infecting weaned piglets, and infection can lead to diseases such as post-weaning diarrhea (PWD). PWD is primarily caused by F4 or F18 adhesin-type *E. coli* strains. Infected pigs typically show severe diarrhea resulting in increased mortality, growth retardation, and economic losses [4,5].

Phytobiotic supplementation has been explored as a potential strategy to prevent or mitigate diseases caused by pathogenic bacteria during the weaning period. Phytobiotics are natural bioactive compounds derived from various plants that support animal health, promote overall growth, and provide protection against infectious diseases [6]. To date, more than 5,000 phytobiotics have been identified from diverse sources such as herbs, essential oils, and agricultural byproducts. Phytobiotics can be administered in various forms, including dried materials, powders, extracts, or solid formulations. Phytobiotics are generally classified into four categories based on their origin and processing characteristics: 1. Herbs (flowering, non-woody, and non-perennial plants); 2. Spices (plants with strong aromas or flavors); 3. Essential oils (volatile lipophilic compounds); 4. Oleoresins (extracts derived from non-aqueous solutions).

The efficacy of phytobiotics as feed additives (PFAs) for pigs has been extensively studied. Numerous reports have demonstrated that dietary inclusion of PFAs improved growth performance in pigs, which was largely attributed to enhanced nutrient digestibility and improved intestinal morphology [7,8]. In addition to their direct effects on intestinal tissues, phytobiotics can modulate gut microbiota composition, with different compounds exerting distinct effects. For example, carvacrol, a phenolic compound found in black pepper and thyme, showed antimicrobial activity comparable to that of conventional antibiotics by reducing bacterial load and suppressing microbial activity in the gastrointestinal tract [9]. Its mechanisms of action include disruption of bacterial cell wall integrity, inhibition of nucleic acid and protein synthesis, and compromise of membrane permeability [10,11]. These antimicrobial effects have been associated with reductions in *Salmonella* and *E. coli* counts in pig feces, along with decreased incidence of diarrhea [12,13]. Conversely, certain phytobiotics promote beneficial microbial populations. Essential oil blends derived from oregano, anise, and citrus peel have been shown to increase the abundance of lactic acid bacteria, thereby enhancing intestinal fermentation capacity [14]. Similarly, supplementation with coix seed has been reported to significantly increased the abundance of *Lactobacillus* and *Bacteroides* in the gastrointestinal tract of weaning piglets. This effect is likely due to its rich composition of starch, oil, polysaccharides, and proteins [15].

Collectively, these findings suggest that phytobiotics not only enhance intestinal integrity and function but also serve as fermentation substrates for beneficial gut microbiota, thereby contributing to host health. As a result, there is growing interest in their application as alternatives to antibiotics, particularly during the weaning period, a time when immune competence is still developing and pigs are highly vulnerable to enteric infections. Despite their potential, comprehensive studies examining the specific effects of phytobiotics on gut health and microbial composition remain limited. Therefore, the present study aimed to investigate the effects of dietary supplementation with phytobiotic compounds on intestinal immunity and gut microbiota composition in weaned

piglets challenged with pathogenic *E. coli*.

MATERIALS AND METHODS

Phytobiotics used in the experiment

Five phytobiotic materials, labeled P1 through P5, were used in both *in vitro* and *in vivo* experiments. All materials were procured from Eugene-Bio. The compositions of the phytobiotic treatments were as follows: P1: bitter citrus extract containing 25%–27% naringin and 11%–15% neohesperidin (BioFlavex GC, HTBA); P2: microencapsulated blend containing 7% thymol and 7% carvacrol (Avipower 2, VetAgro SpA, Reggio Children); P3: mixture containing 40% P1 + 10% P2 + 50% excipient; P4: premixture of grape seed, grape marc extract, green tea, and hops containing 10% flavonoids (AntaOx Flavosyn, DR. Eckel GmbH); P5: fenugreek seed powder containing 12% saponin (Fenugreek Seed Powder, P&D Export).

In-vitro evaluation of cytotoxicity and inflammatory response of phytobiotics in RAW 264.7 cell

The Raw 264.7 murine macrophage cell line (Cat. No AC28116) was obtained from the Korean Collection for Type Cultures (KCTC). Cells were seeded at a density of 1×10^5 cells/well in 500 μL of culture medium in 24-well plates (Corning). Dulbecco's Modified Eagle's Medium (DMEM; Gibco) supplemented with 10% fetal bovine serum (FBS; Gibco) and 1% antibiotics (100 U/mL penicillin and 100 $\mu\text{g}/\text{mL}$ streptomycin) was used for cultivation. Cells were incubated at 37°C in a humidified atmosphere containing 5% CO₂ for 24 hours. Following the initial incubation, 100 μL of each phytobiotic was added to the wells, and cells were then incubated for an additional 24 hours under the same conditions. Triton X-100 was used as a positive control.

To measure the expression levels of TNF- α , NF- κ B (p50), and NF- κ B (p65), total RNA was extracted using the NucleoSpin[®] RNA kit (MACHEREY-NAGEL) after washing the cells twice with 500 μL of 1× DPBS. RNA concentration and purity were measured using a Colibri Microvolume Spectrometer (Titertek Berthold). Subsequently, the RNA was reverse transcribed into complementary DNA (cDNA) using the AccuPower[®] RT PreMix (Bioneer) according to the manufacturer's protocol.

Quantitative real-time polymerase chain reaction (qRT-PCR) was conducted using the CFX Connect[™] Real-Time System (Bio-Rad) to quantify gene expression levels. The qRT-PCR cycling conditions were as follows: initial denaturation at 95°C for 30 seconds, followed by 40 cycles of denaturation at 95°C for 10 seconds and annealing at 60°C for 10 seconds. The reaction concluded with a final step at 65°C for 5 seconds, followed by 95°C. Expression levels of genes associated with immune and inflammatory responses were normalized to the housekeeping gene glyceraldehyde 3-phosphate dehydrogenase (GAPDH), and relative expressions were compared across experimental groups. Primer sequences used in this study are listed in Table 1.

Animals, experimental design and management

The animal experiment was approved by the Institutional Animal Care and Use Committee of Chungbuk National University, Cheongju, Korea (Approval No. CBNUA-1618-21-02). A total of 63 weaned piglets ([Yorkshire × Landrace] × Duroc), 28 days of age with an initial average body weight (BW) of 8.03 ± 0.43 kg, were used in a three-week (21-day) experiment. Piglets were randomly allocated to seven treatment groups based on their initial body weight and *E. coli* challenge status. Each treatment group consisted of nine replicate cages, with one castrated piglet housed per cage. All piglets were housed in individual stainless steel metabolic cages (45 × 55 × 45

Table 1. Primer sequences used for qRT-PCR analysis of inflammatory response gene expression in RAW 264.7 cells

Gene	Primer sequence	Product size (BP)
GAPDH	F 5' TGA TGA CAT CAA GAA GGT GGT GAA G 3' R 5' TCC TTG GAG GCC ATG TGG GCC AT 3'	240
TNF- α	F 5' ATG AGC ACA GAA AGC ATG ATC 3' R 5' TAC AGG CTT GTC ACT CGA ATT 3'	276
NF- κ B (p50)	F 5' GGA GGC ATG TTC GGT AGT GG 3' R 5' CCC TGC GTT GGA TTT CGT G 3'	135
NF- κ B (p65)	F 5' AGG CTT CTG GGC CTT ATG TG 3' R 5' TGC TTC TCT CGC CAG GAA TAC 3'	111

qRT-PCR, quantitative real-time polymerase chain reaction; GAPDH, glyceraldehyde 3-phosphate dehydrogenase.

cm) under optimized environmental conditions. The experimental treatment groups were as follows: NC, negative control (basal diets without *E. coli* challenge); PC, positive control (basal diets + *E. coli* challenge); T1, PC + 0.04 % P1; T2, PC + 0.01 % P2; T3, PC + 0.10 % P3; T4, PC + 0.04 % P4; T5, PC + 0.10 % P5. The basal diet was formulated to meet the nutritional requirements of weaned piglets. The ingredient composition and nutrient contents of the diets are provided in Table 2.

Throughout the 21-day experimental period, piglets had ad libitum access to water, and feed was provided twice daily at 08:30 and 17:30. The feed was mixed with water in a 1:1 ratio immediately before feeding. The *E. coli* challenge was administered orally from days 8 to 10 by delivering 10 mL of nutrient broth containing *E. coli* at a concentration of 1.2×10^{10} CFU/mL.

Measurement of gut epithelial barrier gene expression in mucosa using quantitative real-time polymerase chain reaction

At the end of the experiment (day 21), piglets were euthanized using carbon dioxide gas followed by exsanguination. Intestinal tissue samples were collected from the ileum and colon, at least 10 cm distal to the cecum. The collected samples were rinsed with phosphate-buffered saline (PBS), and mucosal tissues were carefully scraped using sterile scalpel blades. All samples were immediately stored at -80°C for subsequent analysis.

Total RNA was extracted from the mucosal samples using the NucleoSpin[®] RNA kit (MACHEREY-NAGEL) according to the manufacturer's instructions. RNA concentration and purity were assessed using a Colibri Microvolume Spectrometer (Titertek Berthold). cDNA was synthesized from the extracted RNA using the AccuPower[®] RT PreMix (Bioneer) following the manufacturer's protocol.

qRT-PCR was performed using the CFX Connect[™] Real-Time System (Bio-Rad) under the following thermal cycling conditions: initial denaturation at 95°C for 30 seconds, followed by 40 cycles of denaturation at 95°C for 10 seconds and annealing at 55°C for 10 seconds. The reaction concluded with an extension step at 55°C for 5 seconds and a final extension at 95°C .

Gene expression levels of the tight junction protein ZO-1 and mucins (MUC1, MUC2, and MUC3) in the ileal and colonic mucosa were analyzed. Primer sequences used were listed in Table 3. Expression levels were normalized to the housekeeping gene GAPDH and compared across treatment groups.

Fecal sampling and DNA extraction

Fecal samples were collected from three randomly selected piglets per treatment group before the *E. coli* challenge (day 8) and at the end of the experiment (day 21). Fecal samples were collected directly from the rectum of each pig using sterile gloves resulting in a total of 42 samples from 21

Table 2. Compositions of the weaning piglet diets (as-feed basis)

Item	Basal diet
Ingredients (%)	
Corn, Yellow Dent	34.43
Extruded corn	15.00
Lactose	10.00
Dehulled soybean meal (51% CP)	13.50
Soy protein concentrate (65% CP)	10.00
Plasma powder	6.00
Whey	5.00
Soy oil	2.20
MCP	1.26
Limestone	1.40
L-Lysine-HCl (78%)	0.06
DL-Methionine (50%)	0.15
Choline chloride (25%)	0.10
Vitamin premix	0.25
Trace mineral premix	0.25
Salt	0.40
Total	100
Calculated nutrients content	
ME (kcal/kg)	3,433
CP (%)	20.76
Ca (%)	0.82
P (%)	0.65
Lys (%)	1.35
Met (%)	0.39

MCP, monocalcium phosphate monohydrate; ME, metabolizable energy; CP, crude protein; Ca, calcium; P, phosphorus; Lys, lysine; Met: methionine.

Table 3. Primer sequences used for qRT-PCR analysis of gut epithelial barrier gene expression

Gene	Primer sequence	Product size (BP)
GAPDH	F 5' GTA GAG GCA GGG ATG ATG TTC T 3' R 5' CTT TGG TAT CGT GGA AGG ACT C 3'	132
ZO-1	F 5' CCG CCT CCT GAG TTT GAT AG 3' R 5' CAG CTT TAG GCA CTG TGC TG 3'	97
MUC1	F 5' GTG CCG ACG AAA GAA CTG 3' R 5' TGC CAG GTT CGA GTA AGA G 3'	187
MUC2	F 5' CTT CTA GAT GGG TGT GTC TC 3' R 5' GTG GTA GTT GGT GGT GTA 3'	149
MUC3	F 5' CCG GAC CTC AAT GAC AAC ACT 3' R 5' ACC ACG ATG CTG CCA TTC CT 3'	146

qRT-PCR, quantitative real-time polymerase chain reaction; GAPDH, glyceraldehyde 3-phosphate dehydrogenase.

piglets. All fecal samples were immediately stored at -80 °C until further analysis.

For microbial community analysis, total DNA was extracted from 200 mg of feces using the QIAamp Fast DNA Stool Mini Kit (QIAGEN) according to the manufacturer's instructions. The concentration and purity of the extracted DNA were measured using a Colibri Microvolume

Spectrometer (Titertek Berthold). DNA purity with an optical density (OD) ratio of 260/280 between 1.8 and 2.0 was considered to be of high purity and acceptable for downstream applications.

16S rRNA gene sequencing preparation

For amplicon sequencing of the V5–V6 hypervariable regions of the 16S rRNA gene, PCR was performed using primers 799F-mod6 (5'-CMGGATTAGATACCCKG-3') and 1114R (5'-GGTTGCCTCGTTGC-3') [16]. Each 50 µL PCR reaction contained KOD One™ PCR Master Mix -Blue- (TOYOBO), 10 pmol of each primer, and 5 ng/µL of template DNA. The PCR cycling conditions were as follows: initial denaturation at 98°C for 3 minutes, followed by 25 cycles of denaturation at 98°C for 10 seconds, annealing at 57°C for 5 seconds, and extension at 68°C for 1 second, with a final extension at 72°C for 5 minutes. The amplified PCR products were purified using the Wizard® SV Gel and PCR Clean-Up System kit (Promega). Barcoded 16S rRNA gene amplicons were then sequenced on the Illumina MiSeq platform by BRD Korea.

16S rRNA gene sequence analysis

Raw 16S rRNA gene sequencing data were analyzed using the Quantitative Insights into Microbial Ecology 2 (QIIME2) software package [17]. Quality filtering was performed based on a PHRED quality score threshold of 27 to remove low-quality reads and sequences with ambiguous base calls, thereby minimizing the influence of random sequencing errors. The deblur plugin was used to trim sequences to a uniform length of 300 bp, after which amplicon sequence variants (ASVs) were inferred to represent true biological sequences.

For phylogenetic diversity analysis, multiple sequence alignment was conducted using the multiple alignment using fast Fourier transform (MAFFT) pipeline. Alpha diversity metrics were calculated to evaluate species richness and evenness within individual samples. Beta diversity was analyzed to compare microbial community composition between groups, using both weighted (quantitative) and unweighted (qualitative) UniFrac distance metrics. Differences in community structure were visualized through principal coordinate analysis (PCoA) plots. Taxonomic classification of ASVs was performed using a naïve Bayesian classifier trained on the Ribosomal Database Project (RDP) reference database, version 19.

Statistical analysis

Statistical analyses were performed using GraphPad Prism version 8.00 for Windows (GraphPad Software). Significant differences in experimental parameters among treatment groups were assessed using the Kruskal-Wallis H test. Dunn's multiple comparison test was used to determine pairwise differences between treatment groups. To evaluate differences in microbial community structure among experimental groups, the analysis of similarities (ANOSIM) method was applied.

RESULTS

Cytotoxicity and inflammatory responses of RAW 264.7 cells to phytobiotic treatment

The effects of phytobiotics on the expression of immune and inflammation-related markers including TNF- α and NF- κ B were evaluated in RAW 264.7 cells (Fig. 1). Treatment with Triton X-100 (PC) significantly upregulated TNF- α expression compared to the NC. Notably, TNF- α expression in the P4 treatment group was lower than that in the NC group and significantly reduced compared to the PC group. In terms of NF- κ B expression, both p50 and p65 subunits showed decreased levels in the PC group relative to the NC group. The P1, P2, P3, and P5




Fig. 1. Evaluation of cytotoxic effects of phytobiotics by quantifying the relative mRNA expression levels of inflammation-related genes. (A) relative TNF- α mRNA expression (B) relative NF- κ B p50 mRNA expression (C) relative NF- κ B p65 mRNA expression. Expression levels were compared using the $2^{-\Delta\Delta CT}$ method. ^{a,b}Bars with different letters indicate statistically significant differences between groups ($p < 0.05$). NC, negative control (DMEM); PC, positive control (Triton X-100); P1: bitter citrus extract containing 25%-27% naringin and 11%-15% neohesperidin; P2: microencapsulated blend containing 7% thymol and 7% carvacrol; P3: mixture containing 40% P1 + 10% P2 + 50% excipient; P4: premixture of grape seed, grape marc extract, green tea, and hops containing 10% flavonoids; P5: fenugreek seed powder containing 12% saponin. DMEM, Dulbecco's Modified Eagle's Medium.

treatment groups exhibited expression levels generally comparable to or slightly higher than those in the NC group. In contrast, the P4 group showed NF- κ B p50 and p65 subunit expression levels that were comparable to or slightly lower than the NC group.

Comparison of intestinal integrity among the piglets

The *E. coli* challenge and phytobiotic supplementation resulted in significant changes in the expression of tight junction and mucin genes in the ileal and colonic mucosa of weaned piglets (Fig. 2).

In the ileum, the PC group showed relatively lower expression of ZO-1, MUC2, and MUC3




Fig. 2. mRNA expression levels of tight junction and mucin gene in the ileum (A) and colon (B). Expression levels were compared using the $2^{-\Delta\Delta CT}$ method. NC, negative control, (basal diets without *Escherichia coli* challenge); PC, positive control (basal diets + *E. coli* challenge); T1, PC + 0.04 % P1; T2, PC + 0.01 % P2; T3, PC + 0.10 % P3; T4, PC + 0.04 % P4; T5, PC + 0.10 % P5. $^{**}p < 0.01$, $^{***}p < 0.001$.

genes compared to the NC group. In contrast, piglets in the T1 and T5 treatment groups exhibited significantly higher expression of ZO-1 and MUC3 genes compared to the PC group. In the colon, gene expression levels in the PC group were comparable to those in the NC group. However, the T2 group showed significantly higher expression of ZO-1, while both T1 and T2 groups demonstrated significantly elevated MUC3 expression compared to the NC group.

Microbial diversity

From the 42 samples, a total of 5,101,284 raw 16S rRNA gene sequence reads were obtained. After quality filtering, approximately 52% of the reads (2,662,821 reads in total) were retained for downstream analysis with per-sample read counts ranging from 13,738 to 154,970. These high-quality reads were used for microbial community analysis of the weaning piglets across experimental treatments.

Alpha diversity was assessed using Observed Features, Chao1, Shannon, and Simpson indices. No significant differences were observed in alpha diversity before (day 8) and after (day 21) the experiment regardless of the *E. coli* challenge or phytobiotic supplementation (Table 4).

Beta diversity was analyzed using the ANOSIM to compare the weighted and unweighted UniFrac distances. PCoA was used to visualize group clustering (Fig. 3). The ANOSIM results based on unweighted UniFrac distances showed an *R*-value of 0.8402, indicating a distinct shift in the microbial community structure between pre-experiment (day 8) and post-experiment (day 21) samples. However, the PCoA plots based on both weighted and unweighted UniFrac distances showed that the microbial communities from pre-experiment (day 8) and post-experiment (day 21) samples were not clearly separated, indicating substantial overlap in community composition despite the observed structural differences.

Microbial compositions

We investigated the fecal microbial community composition of weaning piglets before and after the *E. coli* challenge and phytobiotic supplementation. Taxonomic assignment of ASVs was performed using the RDP database. At the phylum level, 14 phyla were identified (Fig. 4A). Bactealota was the most dominant phylum in all groups, constituting 43.85%–72.06% on day 8 and 60.24%–74.12% on day 21. Pseudomonadota significantly decreased from day 8 to day 21 in the NC (11.88% to 3.11%), PC (32.27% to 4.95%), T1 (18.41% to 0.76%), T2 (7.81% to 0.65%), and T4 (10.34% to 0.19%) groups. However, it increased in the T3 (5.55% to 7.49%) and T5 (6.47% to 15.67%)

Table 4. Alpha diversity indices of the gut microbiome of piglets before and after *Escherichia coli* challenge and phytobiotic supplementation

	NC	PC	T1	T2	T3	T4	T5	SEM	p-value
Day 8									
Observed features	557.0	431.0	440.3	534.0	314.3	537.7	607.7	37.588	0.079
Chao1	564.0	435.4	444.8	540.6	316.2	544.3	615.9	38.399	0.079
Shannon	4.290	3.844	3.900	4.557	4.011	4.443	4.468	0.111	0.333
Simpson	0.960	0.938	0.943	0.974	0.955	0.964	0.961	0.005	0.493
Day 21									
Observed features	393.7	363.0	346.7	333.7	415.0	410.3	457.0	16.406	0.321
Chao1	397.6	367.0	351.2	336.8	419.5	413.8	462.2	16.561	0.374
Shannon	4.490	4.223	4.147	4.331	4.368	4.341	4.404	0.043	0.827
Simpson	0.972	0.965	0.950	0.969	0.968	0.969	0.958	0.003	0.816

NC, negative control; PC, positive control.




Fig. 3. Principal coordinate analysis (PCoA) plots based on (A) weighted and (B) unweighted UniFrac distance metrics. Different shapes indicate treatment groups as follows: NC, negative control (basal diets without *Escherichia coli* challenge); PC, positive control (basal diets with *E. coli* challenge); T1, PC + 0.04% P1; T2, PC + 0.01% P2; T3, PC + 0.10% P3; T4, PC + 0.04% P4; T5, PC + 0.10% P5.




Fig. 4. Taxonomic classification of 16S rRNA gene sequences at the phylum, family, and genus levels in piglets before and after *Escherichia coli* challenge and phytobiotics supplementation. (A) Phylum level on day 8, (B) Family level on day 8, (C) Genus level on day 8, (D) Phylum level on day 21, (E) Family level on day 21 (F) and (G) Genus level on day 21. Bar plots represent the relative abundance of bacterial taxa. NC, negative control (basal diets without *E. coli* challenge); PC, positive control (basal diets + *E. coli* challenge); T1, PC + 0.04% P1; T2, PC + 0.01% P2; T3, PC + 0.10% P3; T4, PC + 0.04% P4; T5, PC + 0.10% P5.

groups. Additionally, Actinomycetota significantly increased in the T3 group from 1.43% to 11.86%.

At the family level, 71 families were identified, with 12 predominant families identified and the others categorized as “Others” (Fig. 4B). On day 21, the most abundant families across all groups were Oscillospiraceae, Lachnospiraceae, Prevotellaceae, and Clostridiaceae_1. Prevotellaceae

significantly increased in all groups, rising from 1.2%–5.14% on day 8 to 6.22%–19.52% on day 21. In contrast, Enterobacteriaceae, which ranged from 2.03% to 21% on day 8, decreased to only 0.01%–0.65% by day 21. Moraxellaceae significantly increased in the T3 (from 0.002% to 6.27%) and T5 (from 0.64% to 12.55%) groups.

At the genus level, microbial profiling revealed increased relative abundances of *Prevotella* and *Clostridium sensu stricto* in all groups by day 21, while *Hydrogeniclostridium* and *Treponema* decreased (Fig. 4C). *Lactobacillus* showed a decrease in relative abundance from day 8 to day 21 in most groups, except for the T4 group, where it increased from 3.65% to 8.24%. Similarly, *Limosilactobacillus* increased in the NC group (from 1.57% to 3%). However, it decreased in the *E. coli* challenge groups (PC, T1, T2, T3, and T5), while increasing from 0.85% to 3.04% in the T4 group. *Acinetobacter* was predominant only in the T3 (11.79%) and T5 (21.01%) groups on day 21. Within the phylum Actinomycetota, the genera *Olsenella* and *Collinsella* were also identified.

Comparative analysis of genera on day 21 showed that the relative abundance of *Lactobacillus* was higher in the T4 and T5 groups compared to the NC and PC groups (Fig. 5A), with a significant increase in T4 ($p < 0.05$). *Mediterraneibacter* abundance was significantly lower in the PC group compared to the NC group, but it was significantly enriched in the T3 group ($p < 0.05$; Fig. 5B).

Linear discriminant analysis effect size (LEfSe) was conducted with a p-value cutoff of 0.1 and a




Fig. 5. Differential abundance of bacteria among all groups on day 21. Bar plots showing the different relative abundance of a genera *Lactobacillus* (A) and *Mediterraneibacter* (B) among the weaned piglet groups. ^{a-c}Different letters represent statistical differences ($p < 0.05$). Identification of characteristic genera for each dietary group identified by LEfSe (C). The Log_{10} LDA score threshold was set at 4. NC, negative control (basal diets without *Escherichia coli* challenge); PC, positive control (basal diets + *E. coli* challenge); T1, PC + 0.04 % P1; T2, PC + 0.01 % P2; T3, PC + 0.10 % P3; T4, PC + 0.04 % P4; T5, PC + 0.10 % P5.

Log LDA score of 3, and it identified four genera that were significantly enriched in each group on day 21 (Fig. 5C). *Dialister* and *Flintibacter* were characteristic of the T1 group, while *Psychrobacter* was representative of the T2 group. However, *Anaerotigum* was characteristic of the T3 group.

DISCUSSION

In this study, we investigated the cytotoxic and immunomodulatory effects of phytobiotics by analyzing the expression of TNF- α and NF- κ B in RAW 264.7 cells treated with phytobiotics. Overall, the results showed that the tested phytobiotics did not exhibit cytotoxicity and might possess potential for modulating immune responses in weaning piglets.

The mucus layer, primarily composed of mucins such as MUC2 and MUC3, functions as a physical barrier that prevents direct bacterial contact with the epithelial surface, thereby limiting pathogen access and colonization [18,19]. Infection with enterotoxigenic *Escherichia coli* (ETEC) has been shown to compromise epithelial barrier function, resulting in electrolytes and water imbalances and the downregulation of protective mucosal proteins [20]. In our study, the expression of tight junction and mucin genes in the ileum and colon following ETEC challenge revealed that the ileum was more significantly affected. The observed reduction in the expression of tight junction proteins and mucin in the ileum of the PC is likely due to the preferential adhesion of ETEC fimbriae F18 to specific receptors present in the small intestinal epithelium [21,22]. This finding is consistent with previous reports by Gao et al. [23] and Becker et al. [24], which documented decreased expression of occludin and ZO-1 in the jejunum and ileum of pigs infected with ETEC. In this study, phytobiotic supplementation led to the upregulation of mucin and tight junction gene expression, suggesting a potential role in improving epithelial barrier function and maintaining intestinal homeostasis in weaned piglets. These results align with previous findings demonstrating the protective effects of phytobiotics against ETEC-induced intestinal damage. For instance, Liu et al. [25] reported that pigs infected with ETEC and supplemented with phytobiotics exhibited significantly greater villus height and elevated expression of tight junction-associated genes, which contributed to improved intestinal integrity. Similarly, Girard et al. demonstrated that dietary supplementation with chestnut extract rich in tannins reduced the incidence of diarrhea and enhanced growth performance in ETEC-infected pigs, further supporting the anti-inflammatory and barrier-protective properties of phytobiotics [12].

In this study, the diversity of the gut microbiota in weaning piglets generally decreased as weaning progressed, accompanied by a notable shift in microbial community composition. At the family level, a general increase in Lachnospiraceae and Prevotellaceae was observed, while Enterobacteriaceae decreased. These findings are consistent with other studies, which attribute these shifts to the transition from a milk-based diet to solid feed [26–28]. Lachnospiraceae and Prevotellaceae have been widely recognized for their crucial contributions to gut health [29,30]. Both families are prominent producers of short-chain fatty acids (SCFAs) including acetate and butyrate, which play essential roles in maintaining intestinal homeostasis. SCFAs serve as primary energy sources for colonocytes, promote mucosal immunity, and contribute to the regulation of inflammatory responses [31,32]. Moreover, SCFAs strengthen the intestinal barrier by enhancing tight junction integrity and lowering luminal pH, thereby creating an unfavorable environment for pathogenic bacterial colonization [33,34]. The observed increase in Lachnospiraceae and Prevotellaceae during the weaning period may thus reflect not only a microbial adaptation to dietary changes but also a favorable shift toward enhanced mucosal defense and resistance to enteric pathogens.

No significant differences in microbial diversity were observed between the NC group and the

other groups challenged with *E. coli*. Additionally, comparisons of microbial composition revealed minimal differences in community structure between the PC and NC groups. This observation is consistent with previous studies reporting that ETEC exerts only a limited impact on the overall fecal microbial community structure during the post-weaning period [35,36].

It is well established that specific plant secondary metabolites found in phytobiotics can modulate gut bacterial communities by selectively promoting or inhibiting the growth of certain microbial taxa [37]. In our study, distinct shifts in bacterial composition were observed across treatment groups, suggesting compound-specific effects. Notably, the relative abundance of Pseudomonadota decreased more markedly in the T1, T2, and T4 groups compared to the NC and PC groups. In contrast, both Actinomycetota and Pseudomonadota were more prevalent in the T3 and T5 groups, indicating that the phytobiotic blends used in these treatments may favor the proliferation of these phyla.

In the phylum Pseudomonadota of T3 and T5 groups, the majority of the microbial composition was represented by the genus *Acinetobacter*. The genus *Acinetobacter* is typically recognized as an opportunistic pathogen associated with health-related infections [38–40]. However, several studies have reported that *Acinetobacter* species can inhabit the mammalian gut, although the ecological roles of strains other than the commonly studied pathogenic types remain largely unexplored [41]. While the precise mechanisms underlying their presence in the gastrointestinal tract are not fully understood, *Acinetobacter* spp. have been reported to participate in the degradation and metabolism of phytobiotics [42]. This suggests that the increased abundance of *Acinetobacter* observed in our study may be linked to the metabolic activity induced by the specific phytobiotic formulations administered in the T3 and T5 groups.

In the T3 group, the observed increase in the phylum Actinomycetota was attributed to the elevated abundance of the genera *Collinsella* and *Olsenella*. This increase may be associated with carbohydrate fermentation, potentially influenced by the excipients included in the T3 diet. Subramaniam et al. reported that various inactive pharmaceutical excipients, particularly those based on polysaccharides, can serve as fermentable substrates for gut microbes, thereby promoting microbial diversity and abundance [43]. *Collinsella* has been reported to produce SCFAs from both animal- and plant-derived carbohydrates such as lactose, fructose, and starch [44]. Similarly, *Olsenella* species are capable of fermenting carbohydrates and producing SCFAs including acetate, as metabolic by-products [45]. Beyond their SCFA production capacity, increases in *Collinsella* and *Olsenella* have been associated with elevated levels of IL-10, an anti-inflammatory cytokine involved in immune regulation and mucosal homeostasis [46]. These findings suggest that these genera may contribute to maintaining microbial diversity and ecological balance in the gut, thereby limiting pathogen colonization.

Lactobacillus was more abundant in the T4 and T5 groups compared to the NC and PC groups, with significantly higher levels observed in T4. The genus *Lactobacillus* is well known as a beneficial probiotic bacterium [47]. Its increase abundance has been associated with enhanced SCFA production, which can help prevent the invasion of pathogenic bacteria in the gastrointestinal tract while supplying energy to epithelial cells and strengthening gut barrier function [48,49]. Collado et al. [50] demonstrated that *Lactobacillus* can inhibit the adhesion of pathogens, including *E. coli*, to porcine intestinal mucus suggesting its potential role in suppressing *E. coli* colonization.

In addition, *dialister* was more abundant in the T1 group, and *Psychrobacter* was characteristic of the T2 group. Both genera have been reported as commensal gut bacteria commonly found in healthy pigs [51–53]. Yang et al. demonstrated that *Psychrobacter* may function as a probiotic, potentially contributing to increased gut microbial diversity [54]. Furthermore, the genus *Flintibacter* was more abundant in the T1 group. *Flintibacter* has also been shown to produce

butyrate, a key SCFA involved in maintaining gastrointestinal health [55,56]. These findings indicate that the phytobiotics used in this study have the potential to beneficially modulate the gut microbiome of weaning piglets by promoting the growth of commensal and probiotic bacteria.

In summary, our results suggest that dietary supplementation with phytobiotics may enhance immune responses, mitigate inflammatory reactions, and beneficially modulate the gut microbiota in weaned piglets. However, further research is necessary to elucidate the specific roles and functional contributions of the microbial taxa influenced by phytobiotic supplementation, as many of these remain incompletely characterized in the context of gut health. In particular, the variation in microbial responses according to the duration of supplementation and the specific phytogenic compounds used warrants deeper investigation. A more comprehensive understanding of these temporal and compositional dynamics could support the development of optimized dietary strategies aimed at promoting microbial stability and improving host resilience during the critical weaning period.

REFERENCES

1. Saladrigas-García M, D'Angelo M, Ko HL, Nolis P, Ramayo-Caldas Y, Folch JM, et al. Understanding host-microbiota interactions in the commercial piglet around weaning. *Sci Rep.* 2021;11:23488. <https://doi.org/10.1038/s41598-021-02754-6>
2. Kim JC, Hansen CF, Mullan BP, Pluske JR. Nutrition and pathology of weaner pigs: nutritional strategies to support barrier function in the gastrointestinal tract. *Anim Feed Sci Technol.* 2012;173:3-16. <https://doi.org/10.1016/j.anifeedsci.2011.12.022>
3. Campbell JM, Crenshaw JD, Polo J. The biological stress of early weaned piglets. *J Anim Sci Biotechnol.* 2013;4:19. <https://doi.org/10.1186/2049-1891-4-19>
4. Fairbrother JM, Nadeau É, Gyles CL. Escherichia coli in postweaning diarrhea in pigs: an update on bacterial types, pathogenesis, and prevention strategies. *Anim Health Res Rev.* 2005;6:17-39. <https://doi.org/10.1079/AHR2005105>
5. Luise D, Lauridsen C, Bosi P, Trevisi P. Methodology and application of Escherichia coli F4 and F18 encoding infection models in post-weaning pigs. *J Anim Sci Biotechnol.* 2019;10:53. <https://doi.org/10.1186/s40104-019-0352-7>
6. Pandey S, Kim ES, Cho JH, Song M, Doo H, Kim S, et al. Cutting-edge knowledge on the roles of phytobiotics and their proposed modes of action in swine. *Front Vet Sci.* 2023;10:1265689. <https://doi.org/10.3389/fvets.2023.1265689>
7. Wenk C. Herbs and botanicals as feed additives in monogastric animals. *Asian-Australas J Anim Sci.* 2003;16:282-9. <https://doi.org/10.5713/ajas.2003.282>
8. Gheisar MM, Hosseindoust A, Kim IH. Evaluating the effect of microencapsulated blends of organic acids and essential oils in broiler chickens diet. *J Appl Poult Res.* 2015;24:511-9. <https://doi.org/10.3382/japr/pfv063>
9. Manzanilla EG, Perez JF, Martin M, Kamel C, Baucells F, Gasa J. Effect of plant extracts and formic acid on the intestinal equilibrium of early-weaned pigs. *J Anim Sci.* 2004;82:3210-8. <https://doi.org/10.2527/2004.82113210X>
10. Filazi A, Yurdakok-Dikmen B. Nutraceuticals in poultry health and disease. In: Gupta R, Srivastava A, Lall R, editors. *Nutraceuticals in veterinary medicine.* Springer; 2019. p. 661-72.
11. Valenzuela-Grijalva NV, Pinelli-Saavedra A, Muhlia-Almazan A, Domínguez-Díaz D, González-Ríos H. Dietary inclusion effects of phytochemicals as growth promoters in animal production. *J Anim Sci Technol.* 2017;59:8. <https://doi.org/10.1186/s40781-017-0133-9>
12. Girard M, Hu D, Pradervand N, Neuenschwander S, Bee G. Chestnut extract but not sodium

salicylate decreases the severity of diarrhea and enterotoxigenic *Escherichia coli* F4 shedding in artificially infected piglets. PLOS ONE. 2020;15:e0214267. <https://doi.org/10.1371/journal.pone.0214267>

13. Ahmed ST, Hossain ME, Kim GM, Hwang JA, Ji H, Yang CJ. Effects of resveratrol and essential oils on growth performance, immunity, digestibility and fecal microbial shedding in challenged piglets. Asian-Australas J Anim Sci. 2013;26:683-90. <https://doi.org/10.5713/ajas.2012.12683>
14. Kroismayr A, Sehm J, Pfaffl MW, Schedle K, Plitzner C, Windisch W. Effects of avilamycin and essential oils on mRNA expression of apoptotic and inflammatory markers and gut morphology of piglets. Czech J Anim Sci. 2008;53:377-87. <https://doi.org/10.17221/338-CJAS>
15. Li Z, Lin Z, Lu Z, Feng Z, Chen Q, Deng S, et al. Coix seed improves growth performance and productivity in post-weaning pigs by reducing gut pH and modulating gut microbiota. AMB Express. 2019;9:115. <https://doi.org/10.1186/s13568-019-0828-Z>
16. Hanshew AS, Mason CJ, Raffa KF, Currie CR. Minimization of chloroplast contamination in 16S rRNA gene pyrosequencing of insect herbivore bacterial communities. J Microbiol Methods. 2013;95:149-55. <https://doi.org/10.1016/j.mimet.2013.08.007>
17. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol. 2019;37:852-7. <https://doi.org/10.1038/s41587-019-0209-9>
18. Bergstrom KSB, Kissoon-Singh V, Gibson DL, Ma C, Montero M, Sham HP, et al. Muc2 protects against lethal infectious colitis by disassociating pathogenic and commensal bacteria from the colonic mucosa. PLoS Pathog. 2010;6:e1000902. <https://doi.org/10.1371/journal.ppat.1000902>
19. McGuckin MA, Lindén SK, Sutton P, Florin TH. Mucin dynamics and enteric pathogens. Nat Rev Microbiol. 2011;9:265-78. <https://doi.org/10.1038/nrmicro2538>
20. McLamb BL, Gibson AJ, Overman EL, Stahl C, Moeser AJ. Early weaning stress in pigs impairs innate mucosal immune responses to enterotoxigenic *E. coli* challenge and exacerbates intestinal injury and clinical disease. PLOS ONE. 2013;8:e59838. <https://doi.org/10.1371/journal.pone.0059838>
21. Fleckenstein JM, Hardwidge PR, Munson GP, Rasko DA, Sommerfelt H, Steinsland H. Molecular mechanisms of enterotoxigenic *Escherichia coli* infection. Microbes Infect. 2010;12: 89-98. <https://doi.org/10.1016/j.micinf.2009.10.002>
22. Dubreuil JD, Isaacson RE, Schifferli DM. Animal enterotoxigenic *Escherichia coli*. EcoSal Plus. 2016;7:10-1128. <https://doi.org/10.1128/ecosalplus.ESP-0006-2016>
23. Gao Y, Han F, Huang X, Rong Y, Yi H, Wang Y. Changes in gut microbial populations, intestinal morphology, expression of tight junction proteins, and cytokine production between two pig breeds after challenge with *Escherichia coli* K88: a comparative study. J Anim Sci. 2013;91:5614-25. <https://doi.org/10.2527/jas.2013-6528>
24. Becker SL, Li Q, Burrough ER, Kenne D, Sahin O, Gould SA, et al. Effects of an F18 enterotoxigenic *Escherichia coli* challenge on growth performance, immunological status, and gastrointestinal structure of weaned pigs and the potential protective effect of direct-fed microbial blends. J Anim Sci. 2020;98:skaa113. <https://doi.org/10.1093/jas/skaa113>
25. Liu Y, Song M, Che TM, Lee JJ, Bravo D, Maddox CW, et al. Dietary plant extracts modulate gene expression profiles in ileal mucosa of weaned pigs after an *Escherichia coli* infection. J Anim Sci. 2014;92:2050-62. <https://doi.org/10.2527/jas.2013-6422>
26. Hu J, Nie Y, Chen J, Zhang Y, Wang Z, Fan Q, et al. Gradual changes of gut microbiota in weaned miniature piglets. Front Microbiol. 2016;7:1727. <https://doi.org/10.3389/fmicb.2016.01727>

27. Luise D, le Scellour M, Buchet A, Resmond R, Clement C, Rossignol MN, et al. The fecal microbiota of piglets during weaning transition and its association with piglet growth across various farm environments. *PLOS ONE*. 2021;16:e0250655. <https://doi.org/10.1371/journal.pone.0250655>
28. Keum GB, Pandey S, Kim ES, Doo H, Kwak J, Ryu S, et al. Understanding the diversity and roles of the ruminal microbiome. *J Microbiol*. 2024;62:217-30. <https://doi.org/10.1007/s12275-024-00121-4>
29. Guevarra RB, Lee JH, Lee SH, Seok MJ, Kim DW, Kang BN, et al. Piglet gut microbial shifts early in life: causes and effects. *J Anim Sci Biotechnol*. 2019;10:1. <https://doi.org/10.1186/s40104-018-0308-3>
30. Dou S, Gadonna-Widehem P, Rome V, Hamoudi D, Rhazi L, Lakhal L, et al. Characterisation of early-life fecal microbiota in susceptible and healthy pigs to post-weaning diarrhoea. *PLOS ONE*. 2017;12:e0169851. <https://doi.org/10.1371/journal.pone.0169851>
31. Luo Y, Ren W, Smidt H, Wright ADG, Yu B, Schyns G, et al. Dynamic distribution of gut microbiota in pigs at different growth stages: composition and contribution. *Microbiol Spectr*. 2022;10:e00688-21. <https://doi.org/10.1128/spectrum.00688-21>
32. Amat S, Lantz H, Munyaka PM, Willing BP. Prevotella in pigs: the positive and negative associations with production and health. *Microorganisms*. 2020;8:1584. <https://doi.org/10.3390/microorganisms8101584>
33. Zaplana T, Miele S, Tolonen AC. Lachnospiraceae are emerging industrial biocatalysts and biotherapeutics. *Front Bioeng Biotechnol*. 2024;11:1324396. <https://doi.org/10.3389/fbioe.2023.1324396>
34. Machado MG, Sencio V, Trottein F. Short-chain fatty acids as a potential treatment for infections: a closer look at the lungs. *Infect Immun*. 2021;89:e00188-21. <https://doi.org/10.1128/IAI.00188-21>
35. Pollock J, Gally DL, Glendinning L, Tiwari R, Hutchings MR, Houdijk JGM. Analysis of temporal fecal microbiota dynamics in weaner pigs with and without exposure to enterotoxigenic *Escherichia coli* 1,2. *J Anim Sci*. 2018;96:3777-90. <https://doi.org/10.1093/jas/sky260>
36. Jenkins TP, Ács N, Arendrup EW, Swift A, Duzs Á, Chatzigiannidou I, et al. Protecting the piglet gut microbiota against ETEC-mediated post-weaning diarrhoea using specific binding proteins. *npj Biofilms Microbiomes*. 2024;10:42. <https://doi.org/10.1038/s41522-024-00514-8>
37. Santhiravel S, Bekhit AEDA, Mendis E, Jacobs JL, Dunshea FR, Rajapakse N, et al. The impact of plant phytochemicals on the gut microbiota of humans for a balanced life. *Int J Mol Sci*. 2022;23:8124. <https://doi.org/10.3390/ijms23158124>
38. Lee CR, Lee JH, Park M, Park KS, Bae IK, Kim YB, et al. Biology of *Acinetobacter baumannii*: pathogenesis, antibiotic resistance mechanisms, and prospective treatment options. *Front Cell Infect Microbiol*. 2017;7:55. <https://doi.org/10.3389/fcimb.2017.00055>
39. Mancilla-Rojano J, Ochoa SA, Reyes-Grajeda JP, Flores V, Medina-Contreras O, Espinosa-Mazariego K, et al. Molecular epidemiology of *Acinetobacter calcoaceticus*-*Acinetobacter baumannii* complex isolated from children at the Hospital Infantil de México Federico Gómez. *Front Microbiol*. 2020;11:576673. <https://doi.org/10.3389/fmicb.2020.576673>
40. Pandey S, Doo H, Keum GB, Kim ES, Kwak J, Ryu S, et al. Antibiotic resistance in livestock, environment and humans: one health perspective. *J Anim Sci Technol*. 2024;66:266-78. <https://doi.org/10.5187/jast.2023.e129>
41. Glover JS, Browning BD, Ticer TD, Engevik AC, Engevik MA. *Acinetobacter calcoaceticus* is well adapted to withstand intestinal stressors and modulate the gut epithelium. *Front Physiol*. 2022;13:880024. <https://doi.org/10.3389/fphys.2022.880024>

42. Zhang S, Shu J, Xue H, Zhang W, Zhang Y, Liu Y, et al. The gut microbiota in camellia weevils are influenced by plant secondary metabolites and contribute to saponin degradation. *mSystems*. 2020;5:e00692-19. <https://doi.org/10.1128/mSystems.00692-19>
43. Subramaniam S, Kamath S, Ariaee A, Prestidge C, Joyce P. The impact of common pharmaceutical excipients on the gut microbiota. *Expert Opin Drug Deliv*. 2023;20:1297-314. <https://doi.org/10.1080/17425247.2023.2223937>
44. Qin P, Zou Y, Dai Y, Luo G, Zhang X, Xiao L. Characterization a novel butyric acid-producing bacterium *Collinsella aerofaciens* subsp. *Shenzhenensis* subsp. *Nov. Microorganisms*. 2019;7:78. <https://doi.org/10.3390/microorganisms7030078>
45. Wongkuna S, Ghimire S, Janvilisri T, Doerner K, Chankhamhaengdecha S, Scaria J. Taxogenomics description of *Olsenella lakotia* SW165T sp. nov., a new anaerobic bacterium isolated from cecum of feral chicken. *F1000Research*. 2020;9:1103. <https://doi.org/10.12688/f1000research.25823.1>
46. Huang S, Yang L, Wang L, Chen Y, Ding X, Yang F, et al. The effects of octapeptin supplementation on growth performance, serum biochemistry, serum immunity, and gut microbiota in weaned piglets. *Animals*. 2024;14:2546. <https://doi.org/10.3390/ani14172546>
47. Doo H, Kwak J, Keum GB, Ryu S, Choi Y, Kang J, et al. Lactic acid bacteria in Asian fermented foods and their beneficial roles in human health. *Food Sci Biotechnol*. 2024;33:2021-33. <https://doi.org/10.1007/s10068-024-01634-9>
48. Xin J, Zeng D, Wang H, Sun N, Zhao Y, Dan Y, et al. Probiotic *Lactobacillus johnsonii* BS15 promotes growth performance, intestinal immunity, and gut microbiota in piglets. *Probiotics Antimicrob Proteins*. 2020;12:184-93. <https://doi.org/10.1007/s12602-018-9511-Y>
49. Wang T, Teng K, Liu Y, Shi W, Zhang J, Dong E, et al. *Lactobacillus plantarum* PFM 105 promotes intestinal development through modulation of gut microbiota in weaning piglets. *Front Microbiol*. 2019;10:90. <https://doi.org/10.3389/fmicb.2019.00090>
50. Collado MC, Grześkowiak Ł, Salminen S. Probiotic strains and their combination inhibit in vitro adhesion of pathogens to pig intestinal mucosa. *Curr Microbiol*. 2007;55:260-5. <https://doi.org/10.1007/s00284-007-0144-8>
51. Lee JH, Kim S, Kim ES, Keum GB, Doo H, Kwak J, et al. Comparative analysis of the pig gut microbiome associated with the pig growth performance. *J Anim Sci Technol*. 2023;65:856-64. <https://doi.org/10.5187/jast.2022.e122>
52. Nguyen TQ, Martínez-Álvaro M, Lima J, Auffret MD, Rutherford KMD, Simm G, et al. Identification of intestinal and fecal microbial biomarkers using a porcine social stress model. *Front Microbiol*. 2023;14:1197371. <https://doi.org/10.3389/fmicb.2023.1197371>
53. Koh HW, Kim MS, Lee JS, Kim H, Park SJ. Changes in the swine gut microbiota in response to porcine epidemic diarrhea infection. *Microbes Environ*. 2015;30:284-7. <https://doi.org/10.1264/jsme2.ME15046>
54. Yang HL, Sun YZ, Ma RL, Li JS, Huang KP. Probiotic *Psychrobacter* sp. improved the autochthonous microbial diversity along the gastrointestinal tract of grouper *Epinephelus coioides*. *J Aquacult Res Dev*. 2011;1:1-6. <https://doi.org/10.4172/2155-9546.S1-001>
55. Lagkouvardos I, Pukall R, Abt B, Foesel BU, Meier-Kolthoff JP, Kumar N, et al. The mouse intestinal bacterial collection (miBC) provides host-specific insight into cultured diversity and functional potential of the gut microbiota. *Nat Microbiol*. 2016;1:16131. <https://doi.org/10.1038/nmicrobiol.2016.131>
56. Kim H, Chae Y, Cho JH, Song M, Kwak J, Doo H, et al. Understanding the diversity and roles of the canine gut microbiome. *J Anim Sci Biotechnol*. 2025;16:95. <https://doi.org/10.1186/s40104-025-01235-4>