


Effect of polyphosphates on productivity and physiological characteristics in laying hens

Hangsul Cho^{1#}, Ah-Ran Lee^{2#}, Jung-Hyeon Park¹, Yan-Quing Wang¹, Ji-Yun Hwang¹, Yong-Bum Cho³, Soo-Ki Kim^{1,2*}

¹Department of Animal Science and Technology, Konkuk University, Seoul 05029, Korea

²Animal Resources Research Center, Konkuk University, Seoul 05029, Korea

³Department of Animal Resources Science, SahmYook University, Seoul 01795, Korea



Received: Aug 28, 2024

Revised: Dec 12, 2024

Accepted: Jan 9, 2025

#These authors contributed equally to this work.

*Corresponding author

Soo-Ki Kim

Department of Animal Science and Technology, Konkuk University, Seoul 05029, Korea.

Tel: +82-2-450-3728

E-mail: sookikim@konkuk.ac.kr

Copyright © 2026 Korean Society of Animal Science and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Hangsul Cho

<https://orcid.org/0009-0009-7756-5302>

Ah-Ran Lee

<https://orcid.org/0000-0001-7163-0059>

Jung-Hyeon Park

<https://orcid.org/0009-0001-2384-9918>

Yan-Quing Wang

<https://orcid.org/0009-0006-5848-031X>

Ji-Yun Hwang

<https://orcid.org/0009-0005-3936-3188>

Yong-Bum Cho

<https://orcid.org/0009-0001-6008-5275>

Soo-Ki Kim

<https://orcid.org/0000-0003-3499-3330>

Competing interests

No potential conflict of interest relevant to this article was reported.

Abstract

Polyphosphates play a crucial role in enhancing nutrient absorption, energy metabolism, stress response, and cellular function and growth. However, the effects of polyphosphates on the laying performance and gut microbiota of laying hens are not known. This study assessed the effects of polyphosphate supplementation on laying performance, egg quality, blood, organ, tibia and cecal characteristics, and cecal microbiota in laying hens. A total of 100 Lohman Brown laying hens (51 weeks old) were distributed into four dietary treatments in a completely randomized design, each consisting of five replicates with four birds. Treatment groups were designated as corn-soybean meal basal diet (control), basal diet + 0.1% short-chain polyphosphate (P3), basal diet + 0.1% medium-chain polyphosphate (P14), and 0.1% long-chain polyphosphate (P130). Egg productivity rate and egg mass increased in the control and P3 group compared to the P130 group ($p < 0.05$). There was no significant effect of polyphosphate supplementation on egg weight, feed intake, FCR, Haugh unit, eggshell thickness, and eggshell weight. There were no significant differences in tibia strength among the groups. All polyphosphate groups (P3, P14, and P130) showed decreased crypt depth in the ileum and the P130 group showed an increased ratio of villus height to crypt depth (VH/CD) in the ileum ($p < 0.05$). P130 supplementation significantly increased bacterial diversity (ASVs and Chao1 index) in the cecal contents ($p < 0.05$), although there were no significant differences between groups in beta diversity and cecal microbiota composition. In conclusion, supplementation with P130 had a greater effect on the VH/CD ratio and cecal bacterial diversity while not positively affecting egg productivity, suggesting potential benefits for nutrient absorption and gut health in laying hens.

Keywords: Polyphosphate, Egg production, Egg quality, Microbiota, Laying hen

INTRODUCTION

Providing appropriate nutrients and feed additives is crucial for optimizing the health, growth, productivity, and overall well-being of livestock [1]. Novel feed additives have been researched and developed, including functional mineral preparations, natural plant products, and compounds designed

Funding sources

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF), funded by the Ministry of Education (NRF-2022R1F1A1073956).

Acknowledgements

The authors thank to Dr. Toshikazu Shiba for providing polyphosphates.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Cho H, Kim SK.
Data curation: Cho H, Kim SK.
Formal analysis: Cho H, Wang YQ.
Methodology: Cho H, Hwang JY, Cho YB.
Software: Cho H, Park JH.
Validation: Cho H, Lee AR.
Investigation: Cho H, Lee AR.
Writing - original draft: Cho H, Lee AR.
Writing - review & editing: Cho H, Lee AR, Park JH, Wang YQ, Hwang JY, Cho YB, Kim SK.

Ethics approval and consent to participate

The experimental protocol for this study was approved by the Animal Care and Use Committee (IACUC) of Konkuk University, Korea (approval number: KU23227).

to enhance animal health and productivity [2].

Polyphosphates, naturally found in microorganisms, plants, and animals, are known as polyphosphoric acids. They are a class of inorganic compounds composed of multiple orthophosphate units connected by phosphoanhydride bonds [3,4]. These compounds are crucial in various industrial and biological applications due to their unique chemical properties and functional versatility [3]. They contribute to bone formation and regeneration, anti-inflammatory responses, energy metabolism, and buffering and water retention in tissues. Additionally, they exhibit antimicrobial activity and inhibit the growth of pathogenic bacteria [5,6].

The functionality of polyphosphate depends on the molecule's chain length [7]. Short -chain polyphosphates primarily act as an energy carrier, similar to adenosine triphosphate (ATP), which supports cellular metabolism and energy storage [7]. Meanwhile, medium-chain polyphosphates are involved in physiological processes associated with cell growth, differentiation, migration, bone formation, and tissue regeneration [8]. Long -chain polyphosphates play roles in regulating inflammatory responses, blood coagulation, and immune functions [9]. Due to their diverse functional roles based on chain length, polyphosphates demonstrate significant potential as a feed additive in livestock.

Recent studies on polyphosphates supplementation in broilers have been reported [10,11]. The absence of IL-6 and TNF α expression in the intestinal tissues of healthy chickens indicates that polyphosphates did not induce severe inflammation, thereby supporting its potential as a safe dietary intervention. P14 increased the diversity of gut microbiota and decreased the number of *Shigella*, *Salmonella* in the cecum of broilers [10]. Supplementation with long-chain polyphosphates (P130) has improved growth performance, organ characteristics, blood urea nitrogen, and gut bacterial composition in broilers [11]. These studies suggest that medium-chain and long-chain polyphosphates improve productivity, immunity, and a diverse and balanced gut microbiome in broiler chickens.

Although previous studies have demonstrated the potential benefits of polyphosphate supplementation in broilers, there are few studies on how these benefits translate to laying hens, particularly with regard to effects on egg production and quality. Research is needed to determine the specific effects of polyphosphate chain length on productivity and various physiological and in laying hens. Comparing polyphosphate supplementation of various chain lengths could provide insights into the optimal polyphosphate chain length and the benefits of these feed additives for laying hens. Thus, this study aimed to evaluate the effects of supplementation with short, medium, and long-chain polyphosphates on laying performance, egg quality, blood characteristics, visceral organs, tibia characteristics, intestinal histomorphology, and cecal microbiota in laying hens. This research will contribute to more sustainable and efficient poultry production by identifying the most effective types of polyphosphates.

MATERIALS AND METHODS

Preparation of experimental additives

Polyphosphates of varying chain lengths—P3 (short-chain polyphosphate [SCPP]), P14 (medium-chain polyphosphate [MCPP]), and P130 (long-chain polyphosphate [LCPP])—were obtained from RegeneTiss for use in this study.

Experimental animals and design

This study involved one hundred 51-week-old Lohmann Brown laying hens, divided into four groups, each consisting of five replicates with four hens each. The treatment groups received diets

supplemented with 0.1% of P3, P14, or P130, while the control group was given a basic diet devoid of polyphosphate additives. The basal diet used in this experiment was formulated with nutrient levels that meet the requirements of the 2017 Korean Poultry Feeding Standard (Table 1). For the preparation of the polyphosphate-included diets, P3, P14, or P130 was dissolved in one liter of water and mixed with two kilograms of the basic feed. This premix was then thoroughly blended with the main feed using a feed mixer (DKM-350SU, Daekwang) for ten minutes. After a 2-week adaptation period to the basal diet, the experimental diets were then fed for 4 weeks of the experimental period. Feed and water were provided ad libitum, and a lighting schedule of 18 hours of light and 6 hours of darkness was maintained throughout the experiment. The temperature was consistently maintained at $22 \pm 3^\circ\text{C}$. At the end of the experiment, five hens from each treatment group were randomly selected and euthanized with carbon dioxide to assess blood, organ, and tibia characteristics, gut histomorphology, and gut microbiota. The experimental protocol was approved by the Animal Care and Use Committee (IACUC) of Konkuk University (approval number: KU23227).

Table 1. Ingredients and chemical compositions of the basal diet

Items	Amount (%)
Ingredient (%)	
Corn	56.93
Dried distillers' grains with solubles	15.0
Soybean meal (crude protein, 45%)	5.54
Wheat gluten	4.12
Rapeseed meal	2.96
Sesame oil meal	2.04
Beef tallow	0.48
Limestone	11.49
Monocalcium phosphate	0.51
Methionine	0.16
Lysine sulfate	0.30
Threonine	0.02
NaCl	0.24
Choline chloride	0.02
Vitamin Premix ¹⁾	0.07
Mineral Premix ²⁾	0.12
Total	100.0
Calculated chemical composition	
Crude protein (%)	15.00
Crude fat (%)	3.82
Crude fiber (%)	2.73
Crude ash (%)	12.86
Calcium (%)	4.20
Available phosphorus (%)	0.53
AMEn (kcal/kg)	2,700

¹⁾Vitamin mixture provided the following nutrients per kg of diet: vitamin A, 20,000 IU; vitamin D₃, 4600 IU; vitamin E, 40 mg; vitamin K₃, 4 mg; vitamin B₁, 3.6 mg; vitamin B₂, 8 mg; vitamin B₆, 5.8 mg; vitamin B₁₂, 0.04 mg.

²⁾Mineral mixture provided the following nutrients per kg of diet: Fe, 70 mg; Cu, 7.5 mg; Zn, 60 mg; Mn, 80 mg; I, 1 mg; Co, 0.1 mg; Se, 0.2 mg.

AMEn, nitrogen corrected apparent metabolizable energy.

Laying performance

Throughout the experiment, data were collected daily at 10 am regarding the number of eggs, their weight, and any occurrences of damage or deformation. The egg production rate (EPR, %) was calculated weekly by dividing the total number of eggs produced by the initial count of hens and repeated each week. Egg weight (EW) was determined daily by dividing the total weight of the normally produced eggs by their quantity. Egg mass (EM) was derived by multiplying the daily egg weight by the egg production rate. Feed intake (FI) was measured and documented weekly. The feed conversion ratio (FCR) was calculated by dividing the weekly feed intake by the egg mass [12].

Egg quality

Each week, twenty eggs per treatment were collected and evaluated for various parameters including egg weight, Haugh unit, eggshell strength, and eggshell thickness. All egg quality measurements, except for eggshell weight, were performed using a Digital Egg Tester (DET6000, NABEL). The haugh unit (HU) was calculated using the following formula: $HU = 100 \times \text{Log} (H + 7.57 - [1.7 \times W^{0.37}])$. In the aforementioned formula, H denotes the albumen height (mm) and W represents the egg weight (g) [13].

Blood biochemical characteristics

Following euthanasia, cardiac puncture was performed to collect approximately 10 mL of blood from each bird, which was then stored in clot activator tubes (CAT) under refrigeration until analysis. Serum was isolated from the blood samples by centrifugation at 1,500×g for 10 min using a centrifuge (HA-1000-3, Hanil Science Medical) and subsequently stored at -20°C for biochemical analysis. Biochemical parameters such as aspartate aminotransferase (AST), alanine aminotransferase (ALT), blood urea nitrogen (BUN), calcium, total cholesterol (TC), creatinine, glucose, inorganic phosphorus (IP), triglyceride (TG), and total protein (TP) were analyzed using an automatic biochemistry analyzer (CHEM 7000i, Fujifilm) [14].

Organ characteristics

The intestines were segmented into four parts: the duodenum, measured from the pancreatic ring; the jejunum, from the end of the pancreatic ring to Meckel's diverticulum; the ileum, from Meckel's diverticulum to the ileocecal junction; and the cecum, quantified by the average length of the right and left ceca. These segments' lengths were recorded relative to body weight, specifically per 100 g of body weight. The liver and spleen weights were also measured using an electronic scale (EL4002, Mettler Toledo) and recorded relative to 100 g of body weight.

Tibia characteristics

The left tibia was collected from each specimen, with all surrounding tissues removed before analysis. The tibia's breaking strength was determined using a three-point bending test conducted on an Instron Universal Testing Machine (Model 3342). The bone was secured at intervals of 4.0 cm, and force was applied at a crosshead speed of 200 mm/min within a 50 kg load range to determine its strength [15].

Intestinal histomorphology

Samples measuring 2 cm each from the jejunum and ileum were collected and fixed in 10% formalin solutions, respectively. Following fixation, the samples were sent to a slide production company (KP&T) for hematoxylin and eosin staining. The processed slides were analyzed under a microscope (BX43, Olympus) using eXcope X3 software (DIXI Science). Measurements of villus

height (VH) and crypt depth (CD) were taken at five points within each sample, and the averages were calculated. The results are expressed as the ratio of VH to CD.

Population size of bacteria in the cecum

Immediately after dissection, the cecum's contents were preserved on dry ice and transported to the laboratory for microbial enumeration. They were stored in 50 mL conical tubes (Falcon). Various media were used for culturing: Nutrient Agar (NA; Difco) for total bacteria, deMan Rogosa Sharpe (MRS; Difco) for lactobacilli, MacConkey (Difco) for coliform bacteria, and *Shigella* and *Salmonella* (SS; Difco) for detecting *Shigella* and *Salmonella*. Each 1g sample was serially diluted in sterile distilled water and inoculated onto the respective media. Cultures were incubated at 37°C for 24 to 48 hours. Following incubation, colony counts were computed and expressed in Log CFU/g.

Cecal microbiota

Approximately one gram of cecal contents was collected from five birds per treatment and immediately frozen in liquid nitrogen. PCR conditions, DNA extraction, bioinformatics and NGS sequencing analysis were performed according to a previously described method [16]. Briefly, a PowerSoil DNA Isolation Kit (Mobio Laboratories) was first used to isolate genomic DNA. The V3-V4 region of the bacterial 16S rRNA gene was then amplified using 341F and 785R primers. Sequencing was then performed on the Illumina Miseq platform using the commercial service of Macrogen. Amplicon sequence variants (ASVs), Chao1, Shannon and Gini-Simpson indices were checked to compare alpha diversity. Principal coordinate analysis (PCoA) and unweighted pair-group mean average (UPGMA) analysis based on the UniFrac distance matrix were used.

Statistical analysis

The data were subjected to a completely randomized design via the PROC GLM procedure in SAS 9.4 (SAS Institute). Laying performance was evaluated based on replications, and egg quality was assessed on a per-egg basis. Analyses of blood, organs, tibia, and cecal microbiota used individual laying hens as the experimental units. Differences between means were determined using one-way ANOVA with Tukey's test at a significance level of $p < 0.05$. Significance level $0.05 < p < 0.10$ was indicated as trend. Results are presented as mean and standard error of the mean (SEM).

RESULTS AND DISCUSSION

Laying performance

The effects on egg production rate, egg weight, daily egg mass per hen, and FCR are presented in Table 2. The P130 group exhibited a significantly lower egg production rate and egg mass compared to the control and P3 group ($p < 0.05$). The P130 group also tended to have the lowest egg weight ($p = 0.07$). The control's feed intake was lower than that of the P3 group ($p < 0.05$). However, FCR did not significantly differ between groups ($p > 0.05$). Previous studies have demonstrated that sodium long-chain polyphosphate (P130) can significantly reduce FCR in broiler chickens from day 7 to day 21 [11]. Poultry performance results may vary according to factors such as bird breed, age, rearing environment, and supplementation. Phosphorus concentration of each polyphosphate, with a molecular structure of $(\text{NaPO}_3)_n$, can vary due to the different sodium forms, making it challenging to accurately quantify its phosphorus content. Given that polyphosphate was introduced at only 0.1% of the diet, its influence on the overall calcium-to-phosphorus (Ca:P) ratio is minimal and unlikely to affect the balance.

This study suggests that while P130 may negatively affect egg production parameters in laying

Table 2. Effect of supplementation of polyphosphate on laying performance in laying hens

Items	Treatment ¹⁾				SEM	p-value
	Control	P3	P14	P130		
Egg production ratio (%)	96.60 ^a	96.96 ^a	94.99 ^{ab}	92.91 ^b	0.08	0.01
Egg weight (g)	60.4	60.43	60.71	59.55	0.25	0.07
Egg mass (g/d/hen)	58.35 ^a	58.60 ^a	57.67 ^{ab}	55.32 ^b	0.45	0.02
Feed intake (g/d/hen)	106.17 ^b	115.99 ^a	114.46 ^{ab}	111.07 ^{ab}	2.17	0.03
FCR	1.77	1.92	1.88	1.87	0.05	0.71

¹⁾Control, basal diet; P3, basal diet + 0.1% short-chain polyphosphates; P14, basal diet + 0.1% medium-chain polyphosphates; P130, basal diet + 0.1% long-chain polyphosphates.

^{a,b}Means within the same row with different letters significantly differ at $p < 0.05$.

FCR, feed conversion ratio; SEM, standard error of the mean.

hens, its effects on feed efficiency, as indicated by FCR, were not significant in this context. This highlights the need for careful evaluation of the use of P130 in laying hen diets, particularly in light of its potential to reduce overall productivity. The implications of these findings suggest that while P130 has shown benefits in broilers, its use in laying hens may require adjustments in dosage or a reconsideration of its suitability for this purpose. Furthermore, research on the effects of polyphosphate feed additives on animal productivity remains sparse, underscoring the need for further experimental studies and detailed analyses. The limited availability of previous studies makes it challenging to draw definitive conclusions at this time. Future research should focus on identifying the optimal dosage of polyphosphates for laying hens, as well as investigating its long-term effects on laying performance at different stages of the laying cycle.

Egg quality

Table 3 assesses the effects of polyphosphate supplementation in the feed on various egg quality parameters such as weight, Haugh unit, shell -breaking strength, and thickness. The average weight of the eggs tended to be lower in the P130 group compared to the control ($p = 0.07$). The P3 group exhibited significantly greater shell strength compared to the P130 groups ($p < 0.05$). The Haugh unit and eggshell thickness were not significantly different between groups ($p > 0.05$). The findings suggest that different chain lengths of polyphosphate in the feed did not affect the Haugh unit and eggshell thickness. Egg quality is genetically determined by the breed of laying hens and influenced by factors such as environmental conditions, the age of the hens, diet composition, and other variables [17].

Blood biochemical characteristics

Table 4 illustrates the effects of dietary supplementation with polyphosphates of various chain

Table 3. Effect of polyphosphate supplementation on egg quality in laying hens

Items	Treatment ¹⁾				SEM	p-value
	Control	P3	P14	P130		
Egg weight	60.4	60.43	60.71	59.55	0.25	0.07
Haugh unit	88.63	89.34	87.33	87.26	0.51	0.24
Egg breaking strength (kg/cm)	4.56 ^{ab}	4.84 ^a	4.65 ^{ab}	4.5 ^b	0.08	0.03
Eggshell thickness (mm)	0.4	0.41	0.4	0.4	0.05	0.22

¹⁾Control, basal diet; P3, basal diet + 0.1% short-chain polyphosphates; P14, basal diet + 0.1% medium-chain polyphosphates; P130, basal diet + 0.1% long-chain polyphosphates.

^{a,b}Means within the same row with different letters significantly differ at $p < 0.05$.

SEM, standard error of the mean.

Table 4. Effect of polyphosphate supplementation on the blood biochemical characteristics of laying hens

Items	Treatment ¹⁾				SEM	p-value
	Control	P3	P14	P130		
ALT (U/L)	3.8	4.4	3.6	3.2	0.25	0.07
AST (U/L)	147.4	107.6	136.2	122.4	8.61	0.24
BUN (mg/d)	3.2	1.6	2.8	3.6	0.43	0.62
Calcium (mg/d)	26.42	23.01	24.83	21.39	1.09	0.44
Total Cholesterol (mg/d)	68	64.2	73.8	72	2.14	0.22
Creatinine (mg/dL)	0.2	0.06	0.12	0.1	0.03	0.06
Glucose (mg/dL)	220.2	189.2	203.2	200.8	6.39	0.35
IP (mg/dL)	5.16	4.61	5.25	5.58	0.20	0.74
TG (mg/dL)	1,274.4	1,235.4	1,414.2	1,187.2	48.84	0.77
TP (g/dL)	5.36 ^a	4.54 ^b	4.94 ^{ab}	4.56 ^b	0.19	0.03

¹⁾Control, basal diet; P3, basal diet + 0.1% short-chain polyphosphates; P14, basal diet + 0.1% medium-chain polyphosphates; P130, basal diet + 0.1% long-chain polyphosphates.

^{a,b}Means within the same row with different letters significantly differ at $p < 0.05$.

ALT, alanine aminotransferase; AST, aspartate aminotransferase; BUN, blood urea nitrogen; IP, inorganic phosphorus; SEM, standard error of the mean; TG, triglycerides; TP, total protein.

lengths on biochemical blood characteristics. Serum enzyme levels serve as markers of organ or tissue damage [18]. There were no significant differences in ALT, AST, BUN, calcium, total cholesterol, creatinine, glucose, IP, and TG between the groups ($p > 0.05$). TP levels were significantly higher in the control compared to P3 and P130 ($p < 0.05$). However, this variation falls within the range (4.5 to 7.0 g/dL) documented in other studies on healthy laying hens. Adequate serum protein levels are essential for laying hens, facilitating effective utilization of energy and nutrients essential for growth and productivity. These levels are also critical for the normal functioning of biological processes, including immune responses [19,20]. Furthermore, there is a possibility that polyphosphate may influence protein metabolism or utilization, although the precise mechanisms underlying this effect remain unclear. It is hypothesized that polyphosphate may interact with processes such as nutrient absorption, liver protein synthesis, or immune modulation, which highlights the need for further investigation.

Previous poultry studies have also reported that polyphosphate can affect blood parameters, which supports its potential role in modulating physiological processes. Chang et al. [10] reported that P130 increased glucose and triglyceride levels while P3 decreased triglyceride levels in broilers. Moon et al. [11] observed that P130 decreased blood urea nitrogen in broilers. The properties of blood can be influenced by bird breed, environmental factors or different nutritional elements, which affect intermediate metabolic processes [21]. This study indicates that polyphosphate supplementation did not influence mortality or have adverse effects on the health of laying hens.

Organ characteristics

Table 5 reveals the effects of polyphosphate supplementation on the organ characteristics of laying hens. The size and structure of intestines offer valuable insights into how dietary components affect organ function and development in these birds [22]. Longer intestines require more energy for maintenance, which may reduce the energy available for productive activities. Additionally, an extended residence time for digestive enzymes usually leads to an increase in intestinal length. Consequently, shorter intestines can improve nutrient absorption rates and decrease the FCR [23]. There were no significant differences in the relative lengths of the duodenum, jejunum, ileum, and ceca among the groups ($p > 0.05$).

Table 5. Effect of polyphosphate supplementation on organ characteristics in laying hens

Items	Treatment ¹⁾				SEM	p-value
	Control	P3	P14	P130		
Intestinal length (cm/100 g BW)						
Duodenum	1.38	1.46	1.45	1.43	0.02	0.94
Jejunum	3.27	3.34	3.68	3.22	0.10	0.32
Ileum	3.12	3.3	3.27	2.9	0.09	0.46
Ceca	0.88	0.93	0.98	0.85	0.03	0.27
Visceral organ weight (g/100 g BW)						
Liver	2.16 ^{ab}	2.39 ^a	2.14 ^{ab}	1.91 ^b	0.10	0.01
Spleen	0.1	0.09	0.09	0.09	0.003	0.44

¹⁾Control, basal diet; P3, basal diet + 0.1% short-chain polyphosphates; P14, basal diet + 0.1% medium-chain polyphosphates; P130, basal diet + 0.1% long-chain polyphosphates.

^{a,b)}Means within the same row with different letters significantly differ at $p < 0.05$.

BW, body weight; SEM, standard error of the mean.

Regarding liver weight, the P3 group had significantly higher values compared to the P130 group ($p < 0.05$). However, polyphosphate supplementation did not affect liver weight compared with the control group. The liver plays a crucial role in humoral immunity and performs various metabolic and homeostatic functions. Changes in liver weight could directly or indirectly influence health and productivity [24]. Increases in liver weight might result from the accumulation of fat, protein, and water, or from hepatocyte hypertrophy [25]. Additionally, studies on broiler carcass quality have demonstrated that diets including LCPP resulted in reduced liver weights, which correlated directly with body weight [10,26].

This study suggests that polyphosphate supplementation had no adverse effects on organ weight and characteristics. Future studies should evaluate the effects of long-term administration of polyphosphate by optimal chain length on the physiological characteristics of laying hens.

Tibia characteristic

Table 6 demonstrates that polyphosphate supplementation affects the strength of the tibia. Although there was no significant difference in bone strength between the groups, P130 exhibited greater bone strength than the control group. The trends observed in the impact of different chain lengths of polyphosphate on tibia breaking strength are substantial and call for more detailed investigation in future studies [27]. The skeletal system of laying hens is a crucial component in the calcium supply chain, which is essential for eggshell formation. Studies have shown that dietary components can independently enhance egg production, eggshell quality, and bone characteristics in laying hens [28].

Therefore, these findings clarify how the structural diversity of polyphosphate influences the skeletal health of laying hens. The results also suggest that selecting the appropriate forms and feeding strategies of polyphosphate could significantly benefit skeletal health in laying hens [29].

Table 6. Effect of polyphosphate supplementation on tibia characteristics in laying hens

Item	Treatment ¹⁾				SEM	p-value
	Control	P3	P14	P130		
Bone Strength	19.92	21.46	25.74	26.61	1.63	0.16

¹⁾Control, basal diet; P3, basal diet + 0.1% short-chain polyphosphates; P14, basal diet + 0.1% medium-chain polyphosphates; P130, basal diet + 0.1% long-chain polyphosphates.

SEM, standard error of the mean.

Intestinal histomorphology

Table 7 displays the effects of polyphosphate on the microstructural characteristics of the digestive tract in laying hens, noting significant changes in the structural properties of the jejunum and ileum. Polyphosphate groups had no significant effect on the VH compared to control, however, VH increased significantly in the P130 group compared to the P14 group ($p < 0.05$) in the jejunum. An increase in VH indicates a higher number of intestinal cells and enhanced production of digestive enzymes [30]. Additionally, increased VH suggests a larger surface area for nutrient absorption, facilitating improved nutrient uptake [31]. These results reveal that the chain length of polyphosphate affects the nutrient absorption area in the jejunum, thus directly influencing nutrient absorption efficiency.

Significant differences were also observed in the ileum, particularly in CD, with significant reductions noted in the P3, P14, and P130 groups compared to the control ($p < 0.05$). The reduction in CD could indicate enhanced cellular regeneration and restructuring within the intestine, potentially benefiting the health and recovery capabilities of the digestive organs. Optimal CD are associated with intestinal cell regeneration, contributing to improved intestinal health and enhanced nutrient absorption efficiency [32]. Significant differences were noted in the ratio of villus height to crypt depth (VH/CD) in the ileum, with an elevated VH/CD ratio in the P130 group compared to the control ($p < 0.05$). A higher VH/CD ratio suggests an optimal structural morphology of the intestinal epithelium, indicative of a larger surface area for nutrient absorption [33]. [34] reported that biogenic polyphosphate nanoparticles (BPNPs) improved intestinal epithelial barrier function and prevented oxidative stress. Therefore, this study suggests that polyphosphate supplementation may improve barrier function, suppress intestinal inflammation, and promote epithelial cell regeneration, thereby optimizing the VH/CD ratio, improving intestinal mucosal health, and maintaining structural integrity.

Bacterial population and microbial diversity in the cecum

Table 8 illustrates variations in microbial populations within the cecum after polyphosphate supplementation. No significant differences were detected among treatments in the counts of *Lactobacilli*, Coliform bacteria, and *Shigella/Salmonella* ($p > 0.05$), with *Lactobacilli* showing the highest numbers, ranging from 8.35 to 8.50 Log CFU/g. This study suggested that polyphosphate supplementation did not influence changes in the bacterial population. Table 9 analyzed the microbial community in the cecal contents of laying hens using ASVs, the Chao1 index for

Table 7. Effect of polyphosphate supplementation on intestinal histomorphology in laying hens

Items	Treatment ¹⁾				SEM	p-value
	Control	P3	P14	P130		
Jejunum						
VH (um)	1250.85 ^{ab}	1227.00 ^{ab}	999.99 ^b	1377.35 ^a	78.53	0.04
CD (um)	172.99	154.45	143	154.37	6.21	0.42
VH/CD (um/um)	7.35	8.29	7.16	9.28	0.49	0.33
Ileum						
VH (um)	833.35	655.58	720.09	716.4	37.08	0.15
CD (um)	303.51 ^a	190.28 ^b	178.95 ^b	163.80 ^b	31.92	0.02
VH/CD (um/um)	2.94 ^b	3.51 ^{ab}	4.13 ^{ab}	4.48 ^a	0.34	0.01

¹⁾Control, basal diet; P3, basal diet + 0.1% short-chain polyphosphates; P14, basal diet + 0.1% medium-chain polyphosphates; P130, basal diet + 0.1% long-chain polyphosphates.

^{a,b)}Means within the same row with different letters differ significantly at $p < 0.05$. SEM, standard error of the mean; VH, villus height; CD, crypt depth.

Table 8. The effect of polyphosphate on the microbial population size in cecal contents

Items (Log CFU/g)	Treatment ¹⁾				SEM	p-value
	Control	P3	P14	P130		
Microbes on NA	6.21	6.88	6.04	5.93	0.14	0.07
Lactobacilli on MRS	8.50	8.35	8.39	8.39	0.09	0.95
Coliform Bacteria on MacConkey	5.37	5.06	4.76	4.52	0.39	0.9
<i>Shigella</i> and <i>Salmonella</i> on SS	4.52	4.15	4.00	3.88	0.40	0.95

¹⁾Control, basal diet; P3, basal diet + 0.1% short-chain polyphosphates; P14, basal diet + 0.1% medium-chain polyphosphates; P130, basal diet + 0.1% long-chain polyphosphates.

SEM, standard error of the mean.

Table 9. Alpha diversity indices of cecal contents in laying hens

Items	Treatment ¹⁾				SEM	p-value
	Control	P3	P14	P130		
ASVs	523.00 ^b	545.20 ^{ab}	566.00 ^{ab}	616.00 ^a	19.86	0.03
Chao1	527.56 ^b	551.46 ^{ab}	579.11 ^{ab}	631.05 ^a	22.24	0.03
Shannon	7.87	7.92	7.97	8.04	0.04	0.16

¹⁾Control, basal diet; P3, basal diet + 0.1% short-chain polyphosphates; P14, basal diet + 0.1% medium-chain polyphosphates; P130, basal diet + 0.1% long-chain polyphosphates.

^{a,b)}Means within the same row with different letters differ significantly at $p < 0.05$.

ASVs, amplicon sequence variants; SEM, standard error of the mean.

biological richness, and the Shannon index for diversity and evenness [35]. The data indicated that the P130 treatment resulted in higher values for ASVs and the Chao1 index compared to the control ($p < 0.05$), implying that longer chain lengths of polyphosphate contribute to increased microbial diversity and richness in the cecal contents. The Shannon index revealed no significant differences between the groups ($p > 0.05$). A diverse gut microbiome is believed to support a broad range of microbial strains, which contribute to overall health. While diversity is important, maintaining a balanced microbial community is more closely associated with optimal gut health [36]. Polyphosphate supplementation resulted in increased microbial diversity. To explore its potential impact on gut health, beta-diversity analysis and microbial community composition were assessed.

Beta diversity, a crucial concept in ecology and microbiology, reflects the variation in species composition across different environments or samples. Techniques such as PCoA and UPGMA tree are utilized to visualize and analyze beta diversity [37–39]. The effects of polyphosphate supplementation on the beta diversity of the microbial community in cecal contents are depicted in Fig. 1A and 1B. The beta diversity results showed that the PCoA plots and UPGMA tree revealed distinct bacterial communities between the control and P3 groups. Chang et al. [10] also reported differences in bacterial communities between the P130 and control groups in broilers. These results suggest that polyphosphate supplementation can significantly alter the composition of the gut microbiota, potentially resulting in distinct microbial communities.

Intestinal flora of the cecum

The relative abundance of microbial components is illustrated in Fig. 2. The cecal microbiota of laying hens at the phylum level predominantly comprises *Bacteroidetes* (28.95%) and *Bacillota* (55.46%). These two major species, *Bacteroidetes* and *Bacillota*, represent approximately 84% of the total microorganisms in relative abundance (Fig. 2A). Although there were no significant differences between groups at the phylum level, the P14 group tended to show the highest




Fig. 1. The similarity of the bacterial community in cecal contents. (A) Principal coordinate analysis (PCoA) plot based on the weighted UniFrac distance matrix, (B) unweighted pair-group mean average (UPGMA) tree based on the weighted UniFrac distance matrix. Control, basal diet; P3, basal diet + 0.1% short-chain polyphosphates; P14, basal diet + 0.1% medium-chain polyphosphates; P130, basal diet + 0.1% long-chain polyphosphates.

abundance of *Bacteroidota*. *Bacteroidota* are linked to increased fermentative activity and play roles in the degradation of complex carbohydrates and protein metabolism, enhancing gut health [40,41]. Some *Bacteroidota* species may become opportunistic pathogens, particularly in individuals with compromised immune systems or dysbiosis. The *Bacteroidota* phylum includes both beneficial and harmful bacterial species, making the relative proportions of these species critical [42].

Fig. 2B illustrates the classification components of gut flora at the family level. *Lachnospiraceae*,




Fig. 2. Microbiota relative abundance at the phylum level (A) and family level (B). Control, basal diet; P3, basal diet + 0.1% short-chain polyphosphates; P14, basal diet + 0.1% medium-chain polyphosphates; P130, basal diet + 0.1% long-chain polyphosphates.

Bacteroidaceae, and *Oscillospiraceae* constituted the largest percentages at this level. However, there were no significant differences between the groups at the family level. *Lachnospiraceae* and *Oscillospiraceae* belong to the phylum *Firmicutes* and possess beneficial functions, including butyrate

production, inflammation reduction, and fiber fermentation. Nevertheless, in the case of dysbiosis, the gut can lead to health issues such as inflammatory bowel disease and irritable bowel syndrome [43]. *Bacteroidaceae* falls within the phylum *Bacteroidetes* and contributes to nutrient absorption and the prevention of harmful bacteria overgrowth. Certain species within the *Bacteroidaceae* family can become opportunistic pathogens when the gut microbiome balance is disrupted, potentially leading to infections or exacerbating inflammatory conditions [44]. This study indicates that polyphosphate had no effect on cecal dysbiosis, as no health problems or mortalities occurred during the experimental period. In Table 10, polyphosphate supplementation showed no significant difference in the abundance of bacterial genera. The findings reveal that polyphosphate supplementation did not substantially alter the microbial composition in the ceca of laying hens.

Table 11 compares the results of this study with previous broiler research on polyphosphate supplementation, showing that performance, blood characteristics, organ characteristics, and intestinal morphology, etc. Specifically, changes in the broiler microbiome have been observed. Chang et al. [10] reported that P130 supplementation increased the abundance of *Phocaeicola* and *Faecalibacterium* at the genus level in the cecum of broilers. In addition, supplementation with P14 and P130 reduced the population size (Log CFU/g) of *Shigella* and *Salmonella* in the cecum of broilers. Moon et al [11] also found that P130 supplementation increased the population size of beneficial bacteria (*Streptococcus* spp.) in the jejunum of broilers. However, no significant effects on the gut bacterial composition of laying hens were observed in this study. This discrepancy could be attributed to factors such as bird breed, age, metabolic pathways and different nutritional requirements.

Considering these results, further research is necessary to evaluate the suitability of various polyphosphates. Specifically, exploring the impact of polyphosphate chain length and its effects during different stages of the laying period, such as the early, mid, and late stages, would be valuable. Such investigations could elucidate the potential advantages and optimal conditions for employing polyphosphates in poultry, enabling customization to enhance productivity and gut health.

CONCLUSION

In conclusion, polyphosphate supplementation enhanced the VH/CD ratio in the ileum and

Table 10. Abundance (%) of dominant bacterial genera in the cecal microbiota

Items	Treatment ¹⁾				SEM	p-value
	Control	P3	P14	P130		
<i>Bacteroides</i>	8.06	8.19	7.47	7.42	0.20	0.90
<i>Blautia</i>	3.30	3.02	3.72	3.29	0.15	0.42
<i>Faecalibacterium</i>	6.72	5.36	5.61	6.85	0.38	0.44
<i>Gemmiger</i>	0.74	0.82	0.74	0.92	0.04	0.88
<i>Lactobacillus</i>	2.05	2.92	3.36	3.08	0.28	0.77
<i>Limosilactobacillus</i>	1.67	1.48	1.70	1.07	0.15	0.75
<i>Mediterraneibacter</i>	10.18	10.30	9.94	10.17	0.07	0.99
<i>Olsenella</i>	1.67	1.28	1.10	1.57	0.13	0.35
<i>Parabacteroides</i>	2.02	1.77	2.20	2.10	0.09	0.36
<i>Paraphocaeicola</i>	2.90	2.24	2.00	2.46	0.19	0.34
<i>Phocaeicola</i>	8.22	10.28	10.58	6.68	0.92	0.12
<i>Prevotella</i>	2.34	1.95	2.91	2.88	0.23	0.31

¹⁾Control, basal diet; P3, basal diet + 0.1% short-chain polyphosphates; P14, basal diet + 0.1% medium-chain polyphosphates; P130, basal diet + 0.1% long-chain polyphosphates.

SEM, standard error of the mean.

Table 11. Comparison of the effects of polyphosphate supplementation in laying hens and broilers

	Laying hens	Broilers	Broilers
Tested	P3, P14, P130	P3, P14, P130	P130
Ref.	This study	Chang et al. [10]	Moon et al. [11]
Items			
Performance	- Egg production and egg mass increased in control and P3 compared to P130 - Feed intake in P3 increased	ND	- P130 improved body weight, average daily gain and decreased FCR during grower phase
Egg or meat quality	- No significant effect	- No significant effect	ND
Blood characteristic	- P3 and P130 decreased total protein levels	- P130 increased glucose and triglyceride levels - P3 decreased triglyceride levels	- P130 decreased blood urea nitrogen
Organ characteristics	- No significant effect	- Liver weight was lower in P130 than control - P3 increased jejunum length	- Duodenum and ileum length was shorter in the P130 than the control
Intestinal morphology	- CD in P3, P14, and P130 was lower than control - VH/CD ratio in P130 was higher than control	ND	ND
Population size of bacteria in the gut	- No significant effects	- P3 and P14 decreased coliform bacteria - P14 and P130 decreased <i>Shigella</i> and <i>Salmonella</i> in the cecum	- P130 increased coliforms and lactose negative enterobacteria in the jejunum - P130 increased <i>Streptococcus</i> spp. in the jejunum
Alpha diversity in the cecum	- P130 increased ASVs and Chao1	- P14 increased ASVs and Chao1	ND
Intestinal flora in the cecum	- No significant effects	- P3 and P14 decreased <i>Bacteroides</i> at the genus level - P130 increased <i>Phocaeicola</i> and <i>Faecilibacterium</i> - P3 increased <i>Barnesiella</i>	ND

ND, not detected; FCR, feed conversion ratio; CD, crypt depth; VH, villus height.

increased bacterial diversity in the cecum, without adversely affecting blood and organ characteristics in laying hens. Among the three polyphosphates with varying chain lengths, P130 was found to be more effective than P3 and P14, potentially improving nutrient absorption and the diversity of the gut microbiome in laying hens.

REFERENCES

- Adedokun SA, Olojede OC. Optimizing gastrointestinal integrity in poultry: the role of nutrients and feed additives. *Front Vet Sci.* 2019;5:348. <https://doi.org/10.3389/fvets.2018.00348>
- Han D, Ren T, Yang Y, Li Z, Du X, Zhang C, et al. Application and substitution of antibiotics in animal feeding. *Med Weter.* 2023;80:5-11. <https://doi.org/10.21521/mw.6830>
- Achbergerová L, Nahálka J. Polyphosphate - an ancient energy source and active metabolic regulator. *Microb Cell Fact.* 2011;10:63. <https://doi.org/10.1186/1475-2859-10-63>
- Kus F, Smolenski RT, Tomczyk M. Inorganic polyphosphate: regulator of cellular metabolism in homeostasis and disease. *Biomedicines.* 2022;10:913. <https://doi.org/10.3390/biomedicines10040913>
- Harada K, Itoh H, Kawazoe Y, Miyazaki S, Doi K, Kubo T, et al. Polyphosphate-mediated inhibition of tartrate-resistant acid phosphatase and suppression of bone resorption of osteoclasts. *PLOS ONE.* 2013;8:e78612. <https://doi.org/10.1371/journal.pone.0078612>

6. Morita K, Doi K, Kubo T, Takeshita R, Kato S, Shiba T, et al. Enhanced initial bone regeneration with inorganic polyphosphate-adsorbed hydroxyapatite. *Acta Biomater.* 2010;6:2808-15. <https://doi.org/10.1016/j.actbio.2009.12.055>
7. Pirttiniemi A, Adeshara K, Happonen N, Einarsdottir E, Katayama S, Salmenkari H, et al. Long-chain polyphosphates inhibit type I interferon signaling and augment LPS-induced cytokine secretion in human leukocytes. *J Leukoc Biol.* 2023;114:250-65. <https://doi.org/10.1093/jleuko/qiad058>
8. Wang Y, Li M, Li P, Teng H, Fan D, Du W, et al. Progress and applications of polyphosphate in bone and cartilage regeneration. *BioMed Res Int.* 2019;2019:5141204. <https://doi.org/10.1155/2019/5141204>
9. Shiba T. Inorganic polyphosphate and its chain-length dependency in tissue regeneration including bone remodeling and teeth whitening. In: Kulakovskaya T, Pavlov E, Dedkova E, editors. *Inorganic polyphosphates in eukaryotic cells.* Springer; 2016. p. 139-58.
10. Chang YQ, Moon SG, Wang YQ, Oh JS, Jeon SW, Lee AR, et al. Effects of polyphosphates with different chain lengths on digestive organ weight, carcass quality, and immune response, and intestinal microflora in broilers. *J Anim Sci Technol.* 2024. <https://doi.org/10.5187/jast.2024.e57>
11. Moon SG, Kothari D, Kim WL, Lee WD, Kim KI, Kim JI, et al. Feasibility of sodium long chain polyphosphate as a potential growth promoter in broilers. *J Anim Sci Technol.* 2021;63:1286-300. <https://doi.org/10.5187/jast.2021.e110>
12. Bonekamp RPRT, Lemme A, Wijtten PJA, Sparla JKWM. Effects of amino acids on egg number and egg mass of brown (heavy breed) and white (light breed) laying hens. *Poult Sci.* 2010;89:522-9. <https://doi.org/10.3382/ps.2009-00342>
13. Haugh H. The Haugh unit for measuring egg quality. *The U.S. Egg & Poultry Magazine*; 1937. p. 552-5, 572-3.
14. Aryal M, Poudel A, Satyal B, Gyawali P, Pokheral BR, Raut BK, et al. Evaluation of non-HDL-c and total cholesterol: HDL-c ratio as cumulative marker of cardiovascular risk in diabetes mellitus. *Kathmandu Univ Med J.* 2010;8:398-404. <https://doi.org/10.3126/kumj.v8i4.6239>
15. Ndazigaruye G, Kim DH, Kang CW, Kang KR, Joo YJ, Lee SR, et al. Effects of low-protein diets and exogenous protease on growth performance, carcass traits, intestinal morphology, cecal volatile fatty acids and serum parameters in broilers. *Animals.* 2019;9:226. <https://doi.org/10.3390/ani9050226>
16. Niu KM, Khosravi S, Kothari D, Lee WD, Lim JM, Lee BJ, et al. Effects of dietary multi-strain probiotics supplementation in a low fishmeal diet on growth performance, nutrient utilization, proximate composition, immune parameters, and gut microbiota of juvenile olive flounder (*Paralichthys olivaceus*). *Fish Shellfish Immunol.* 2019;93:258-68. <https://doi.org/10.1016/j.fsi.2019.07.056>
17. Jha R, Das R, Oak S, Mishra P. Probiotics (direct-fed microbials) in poultry nutrition and their effects on nutrient utilization, growth and laying performance, and gut health: a systematic review. *Animals.* 2020;10:1863. <https://doi.org/10.3390/ani10101863>
18. Tang SGH, Sieo CC, Ramasamy K, Saad WZ, Wong HK, Ho YW. Performance, biochemical and haematological responses, and relative organ weights of laying hens fed diets supplemented with prebiotic, probiotic and synbiotic. *BMC Vet Res.* 2017;13:248. <https://doi.org/10.1186/s12917-017-1160-y>
19. Kim CH, Kang HK. Effects of energy and protein levels on laying performance, egg quality, blood parameters, blood biochemistry, and apparent total tract digestibility on laying hens in an

- aviary system. *Animals*. 2022;12:3513. <https://doi.org/10.3390/ani12243513>
20. Sun M, Ma N, Liu H, Liu Y, Zhou Y, Zhao J, et al. The optimal dietary arginine level of laying hens fed with low-protein diets. *J Anim Sci Biotechnol*. 2022;13:63. <https://doi.org/10.1186/s40104-022-00719-X>
 21. Afsharmanesh M, Lotfi M, Mehdipour Z. Effects of wet feeding and early feed restriction on blood parameters and growth performance of broiler chickens. *Anim Nutr*. 2016;2:168-72. <https://doi.org/10.1016/j.aninu.2016.04.002>
 22. Elnesr SS, Alagawany M, Elwan HAM, Fathi MA, Farag MR. Effect of sodium butyrate on intestinal health of poultry – a review. *Ann Anim Sci*. 2020;20:29-41. <https://doi.org/10.2478/aoas-2019-0077>
 23. Marume U, Mokagane JM, Shole CO, Hugo A. *Citrullus lanatus* essential oils inclusion in diets elicit nutraceutical effects on egg production, egg quality, and physiological characteristics in layer hens. *Poult Sci*. 2020;99:3038-46. <https://doi.org/10.1016/j.psj.2020.01.029>
 24. Selim S, Abdel-Megeid NS, Abou-Elnaga MK, Mahmoud SF. Early nutrition with different diets composition versus fasting on immunity-related gene expression and histomorphology of digestive and lymphoid organs of layer-type chicks. *Animals*. 2021;11:1568. <https://doi.org/10.3390/ani11061568>
 25. Kim MJ, Lee JS, Ha OM, Jang JY, Cho SY. Effects of *Pueraria thunbergiana* bentham water extracts on hepatic alcohol metabolic enzyme system in rats. *J Korean Soc Food Sci Nutr*. 2002;31:92-7. <https://doi.org/10.3746/jkfn.2002.31.1.092>
 26. Qureshi AA, Burger WC, Prentice N, Bird HR, Sunde ML. Regulation of lipid metabolism in chicken liver by dietary cereals. *J Nutr*. 1980;110:388-93. <https://doi.org/10.1093/jn/110.3.388>
 27. Taylor CE, Henninger HB, Bachus KN. Cortical and medullary morphology of the tibia. *Anat Rec*. 2021;304:507-17. <https://doi.org/10.1002/ar.24479>
 28. Alfonso-Carrillo C, Benavides-Reyes C, de los Mozos J, Dominguez-Gasca N, Sanchez-Rodríguez E, Garcia-Ruiz AI, et al. Relationship between bone quality, egg production and eggshell quality in laying hens at the end of an extended production cycle (105 weeks). *Animals*. 2021;11:623. <https://doi.org/10.3390/ani11030623>
 29. Huang S, Kong A, Cao Q, Tong Z, Wang X. The role of blood vessels in broiler chickens with tibial dyschondroplasia. *Poult Sci*. 2019;98:6527-32. <https://doi.org/10.3382/ps/pez497>
 30. Mousavi A, Mahdavi AH, Riasi A, Soltani-Ghombavani M. Efficacy of essential oils combination on performance, ileal bacterial counts, intestinal histology and immunocompetence of laying hens fed alternative lipid sources. *J Anim Physiol Anim Nutr*. 2018;102:1245-56. <https://doi.org/10.1111/jpn.12942>
 31. Arslan C, Pirinç A, Eker N, Sur E, Ündağ İ, Kuşat T. Dietary encapsulated essential oil mixture influence on apparent nutrient digestibility, serum metabolic profile, lymphocyte histochemistry and intestinal morphology of laying hens. *Anim Biosci*. 2022;35:740-51. <https://doi.org/10.5713/ab.21.0275>
 32. Awad WA, Ghareeb K, Abdel-Raheem S, Böhm J. Effects of dietary inclusion of probiotic and synbiotic on growth performance, organ weights, and intestinal histomorphology of broiler chickens. *Poult Sci*. 2009;88:49-56. <https://doi.org/10.3382/ps.2008-00244>
 33. Ramirez SY, Peñuela-Sierra LM, Ospina MA. Effects of oregano (*Lippia origanoides*) essential oil supplementation on the performance, egg quality, and intestinal morphometry of Isa Brown laying hens. *Vet World*. 2021;14:595-602. <https://doi.org/10.14202/vetworld.2021.595-602>
 34. Feng G, Feng Y, Guo T, Yang Y, Guo W, Huang M, et al. Biogenic polyphosphate nanoparticles from *Synechococcus* sp. PCC 7002 exhibit intestinal protective potential in human intestinal epithelial cells in vitro and murine small intestine ex vivo. *J Agric Food Chem*. 2018;66:8026-35.

- <https://doi.org/10.1021/acs.jafc.8b03381>
35. He Y, Zhou BJ, Deng GH, Jiang XT, Zhang H, Zhou HW. Comparison of microbial diversity determined with the same variable tag sequence extracted from two different PCR amplicons. *BMC Microbiol.* 2013;13:208. <https://doi.org/10.1186/1471-2180-13-208>
 36. Barko PC, McMichael MA, Swanson KS, Williams DA. The gastrointestinal microbiome: a review. *J Vet Intern Med.* 2018;32:9-25. <https://doi.org/10.1111/jvim.14875>
 37. Lozupone CA, Hamady M, Kelley ST, Knight R. Quantitative and qualitative β diversity measures lead to different insights into factors that structure microbial communities. *Appl Environ Microbiol.* 2007;73:1576-85. <https://doi.org/10.1128/AEM.01996-06>
 38. Wang Y, Sun F, Lin W, Zhang S. AC-PCoA: adjustment for confounding factors using principal coordinate analysis. *PLoS Comput Biol.* 2022;18:e1010184. <https://doi.org/10.1371/journal.pcbi.1010184>
 39. Chang Q, Luan Y, Sun F. Variance adjusted weighted UniFrac: a powerful beta diversity measure for comparing communities based on phylogeny. *BMC Bioinform.* 2011;12:118. <https://doi.org/10.1186/1471-2105-12-118>
 40. Clausen U, Vital ST, Lambertus P, Gehler M, Scheve S, Wöhlbrand L, et al. Catabolic network of the fermentative gut bacterium *Phocaeicola vulgatus* (phylum Bacteroidota) from a physiologic-proteomic perspective. *Microb Physiol.* 2024;34:88-107. <https://doi.org/10.1159/000536327>
 41. Myhrstad MCW, Tunsjø H, Charnock C, Telle-Hansen VH. Dietary fiber, gut microbiota, and metabolic regulation—current status in human randomized trials. *Nutrients.* 2020;12:859. <https://doi.org/10.3390/nu12030859>
 42. Shin JH, Tillotson G, MacKenzie TN, Warren CA, Wexler HM, Goldstein EJC. Bacteroides and related species: the keystone taxa of the human gut microbiota. *Anaerobe.* 2024;85:102819. <https://doi.org/10.1016/j.anaerobe.2024.102819>
 43. Carding S, Verbeke K, Vipond DT, Corfe BM, Owen LJ. Dysbiosis of the gut microbiota in disease. *Microb Ecol Health Dis.* 2015;26:26191. <https://doi.org/10.3402/mehd.v26.26191>
 44. Wang C, Zhao J, Zhang H, Lee YK, Zhai Q, Chen W. Roles of intestinal bacteroides in human health and diseases. *Crit Rev Food Sci Nutr.* 2021;61:3518-36. <https://doi.org/10.1080/10408398.2020.1802695>