

# Modulation of microbial community and metabolism through *Lactiplantibacillus argentoratensis* AGMB00912 supplementation in weaning piglets

Ki-Nam Yoon<sup>1,2#</sup>, Han Gyu Lee<sup>3#</sup>, Seo-Joon Yeom<sup>1</sup>, Sang-Su Kim<sup>1</sup>, Jong-Heum Park<sup>1</sup>, Beom-Seok Song<sup>1</sup>, Seung-Won Yi<sup>3</sup>, Tai-Young Hur<sup>3</sup>, Jong-Bang Eun<sup>4</sup>, Seung-Hwan Park<sup>5</sup>, Ju Huck Lee<sup>5</sup>, Hyeun Bum Kim<sup>6</sup>, Ju-Hoon Lee<sup>7,8,9</sup>, Jae-Kyung Kim<sup>1\*</sup>

<sup>1</sup>Advanced Radiation Technology Institute, Korea Atomic Energy Research Institute, Jeongeup 56212, Korea

<sup>2</sup>Research Institute of Agricultural Science and Technology, Chonnam National University, Gwangju 61186, Korea

<sup>3</sup>Animal Diseases and Health Division, National Institute of Animal Science, Rural Development Administration, Wanju 55365, Korea

<sup>4</sup>Department of Food Science and Technology, Graduate School of Chonnam National University, Gwangju 61186, Korea


<sup>5</sup>Korean Collection for Type Cultures, Korea Research Institute of Bioscience and Biotechnology, Jeongeup 56212, Korea

<sup>6</sup>Department of Animal Biotechnology, Dankook University, Cheonan 31116, Korea

<sup>7</sup>Department of Agricultural Biotechnology, Seoul National University, Seoul 08826, Korea

<sup>8</sup>Department of Food and Animal Biotechnology, Seoul National University, Seoul 08826, Korea

<sup>9</sup>Center for Food and Bioconvergence, Seoul National University, Seoul 08826, Korea



Received: Sep 4, 2024  
Revised: Dec 10, 2024  
Accepted: Dec 15, 2024

#These authors contributed equally to this work.

## \*Corresponding author

Jae-Kyung Kim  
Advanced Radiation Technology Institute, Korea Atomic Energy Research Institute, Jeongeup 56212, Korea.  
Tel: +82-63-570-3219  
E-mail: [jkim@kaeri.re.kr](mailto:jkim@kaeri.re.kr)

Copyright © 2026 Korean Society of Animal Science and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

Dietary supplementation effects with *Lactiplantibacillus argentoratensis* strain AGMB00912 (LA) on gut microbiota and metabolic functions of weaned piglets were investigated. Eight 25-day-old weaned piglets were evenly divided into a control group and an LA-supplemented group, with the LA group receiving  $1.0 \times 10^8$  CFU/mL of LA daily for 10 days. Fecal samples taken on the 10<sup>th</sup> day were analyzed using 16S rRNA gene sequencing to assess microbial composition and metabolic function prediction. Supplementation with LA promoted a stable microbial environment by increasing the relative abundance of short-chain fatty acid-producing bacteria, including *Faecalitalea*, *Catenibacterium*, and *Butyrivibrio*, while reducing harmful genera like *Treponema* and *Campylobacter*. Administration of LA significantly influenced the metabolic activity of the microbial community, particularly by upregulating carbohydrate metabolism pathways, which enhanced the capacity for short-chain fatty acid production. This shift in microbial metabolism also extended to pathways involved in the biosynthesis of amino acids, lipids, cofactors, and vitamins, indicating an improved capacity for microbial-driven nutrient assimilation and utilization. Furthermore, LA supplementation promoted the biosynthesis of antimicrobial non-ribosomal peptides within the microbiome,

**ORCID**

Ki-Nam Yoon  
<https://orcid.org/0009-0001-1941-1200>  
 Han Gyu Lee  
<https://orcid.org/0000-0002-3531-1971>  
 Seo-Joon Yeom  
<https://orcid.org/0009-0003-9080-2436>  
 Sang-Su Kim  
<https://orcid.org/0009-0004-5810-3245>  
 Jong-Heum Park  
<https://orcid.org/0000-0002-9525-8543>  
 Beom-Seok Song  
<https://orcid.org/0000-0001-9531-6195>  
 Seung-Won Yi  
<https://orcid.org/0000-0001-5545-2969>  
 Tai-Young Hur  
<https://orcid.org/0000-0003-3129-2942>  
 Jong-Bang Eun  
<https://orcid.org/0000-0002-2917-4923>  
 Seung-Hwan Park  
<https://orcid.org/0000-0003-4613-4680>  
 Ju Huck Lee  
<https://orcid.org/0000-0001-6103-7593>  
 Hyeun Bum Kim  
<https://orcid.org/0000-0003-1366-6090>  
 Ju-Hoon Lee  
<https://orcid.org/0000-0003-0405-7621>  
 Jae-Kyung Kim  
<https://orcid.org/0000-0002-3746-7347>

**Competing interests**

No potential conflict of interest relevant to this article was reported.

**Funding sources**

This research was supported by the Korea Atomic Energy Research Institute (KAERI) Institutional Program (No. 523320 26), and by the project 'Development of gut microbiota for preventing intestinal diseases and its impact on host immunity in pigs' (Grant ID No. PJ01564402), funded by the Rural Development Administration of South Korea.

**Acknowledgements**

Not applicable.

**Availability of data and material**

The high-throughput sequencing data generated in this study have been deposited in the SRA database under accession number PRJNA1045807.

**Authors' contributions**

Conceptualization: Yoon KN, Lee HG, Park JH, Song BS, Yi SW, Hur TY, Park SH, Ju Huck Lee, Kim HB, Lee Ju-Hoon, Kim JK.  
 Data curation: Yoon KN, Yeom SJ, Kim SS.  
 Formal analysis: Yoon KN, Yeom SJ, Kim SS.  
 Methodology: Yoon KN, Lee HG, Hur TY, Kim JK.  
 Validation: Yoon KN, Lee HG, Park JH, Song BS, Yi SW, Hur TY, Eun JB, Park SH, Ju Huck Lee, Kim HB, Lee Ju-Hoon, Kim JK.  
 Writing - original draft: Yoon KN, Lee HG, Kim JK.  
 Writing - review & editing: Yoon KN, Lee HG, Yeom SJ, Kim SS, Park JH, Song BS, Yi SW, Hur TY, Eun JB, Park SH, Lee Ju Huck, Kim HB, Lee Ju-Hoon, Kim JK.

crucial for inhibiting the growth of pathogenic microorganisms and maintaining microbial balance. The modulation of microbial metabolism is also predicted to reduce glycan degradation and increase peptidoglycan biosynthesis, contributing to enhanced gut barrier function and a more regulated immune response. These metabolic changes within the microbial community are predicted to stabilize the gut microbiota, providing enhanced disease resistance and supporting the overall health and growth of weaned piglets.

**Keywords:** Weaning transition, Probiotics, *Lactiplantibacillus argenterotensis* AGMB00912, Gut microbiota, Metabolic function prediction

**INTRODUCTION**

The gut microbiota plays a vital role in supporting the health and development of piglets, particularly during the weaning transition [1]. In the suckling phase, the gut microbial community is altered and influenced by oligosaccharides present in the sow's milk, which promote the proliferation of beneficial bacteria, including *Lactobacillus* [2]. These oligosaccharides also support the proliferation of genera such as *Escherichia* and *Streptococcus*, which contribute to the development of an anaerobic intestinal environment [3], conducive to colonization by various genera including *Bacteroides*, *Bifidobacterium* and *Clostridium*, thus enriching the diversity of the intestinal microbial community [4]. However, a solid-type weaning diet, which refers to the solid food given during the weaning period, dramatically alters the bacterial communities due to its high proportion of grain and crude protein content [5]. Cao et al. [5] reported that soybean and pectin-rich diets could potentially reduce the proportion of *Lactobacillus* and increase the relative abundance of *Prevotella* in the large intestine. Moreover, the abrupt proliferation of *Escherichia* and *Shigella* is attributed to the high protein levels in the diet [6]. This disruption of the intestinal microbiota creates an environment susceptible to infection by Proteobacteria, such as *Escherichia* and *Salmonella*, which are typical post-weaning diarrhea pathogens [7,8]. Infection-induced inflammation of the intestine also creates a favorable environment for the growth of Proteobacteria [9]. Nitric oxide, generated during the intestinal inflammatory response, is converted into nitrate, which supports the growth of *Escherichia coli* strains carrying the nitrate reductase gene [10]. Additionally, increased blood flow to the inflamed intestine raises oxygen levels, thus resulting in an increased proportion of facultative anaerobes such as Proteobacteria [11]. This shift disrupts anaerobic conditions and initiates a cycle of adverse conditions which ultimately lead to a loss of bacterial diversity [2]. Therefore, maintaining gut homeostasis during the weaning transition by regulating microbial communities is a crucial challenge in the swine industry.

Probiotic microorganisms have been evaluated as non-antibiotic approaches to restore intestinal microbial balance and inhibit pathogenic microbial infections by producing health-promoting bioactive compounds such as short-chain fatty acids (SCFA), bacteriocins, enzymes, and vitamins [12,13]. *Lactiplantibacillus plantarum*, a member of the beneficial probiotic group, includes the subspecies *Lactiplantibacillus argenterotensis*, previously known as *Lactobacillus argenterotensis*. This bacterium is a gram-positive, facultative anaerobe capable of both homo- and heterofermentation [14]. This bacterium produces SCFA and other metabolites like lactate and acetate, which contribute to gut health by supporting beneficial microbial functions and reducing pathogenic populations through the production of bioactive compounds like hydrogen peroxide. Additionally, the ability of *L. argenterotensis* to ferment carbohydrates through the Embden-Meyerhoff-Parnas and phosphoketolase pathways enhances its metabolic versatility [15]. In our previous study, we isolated *L. argenterotensis* AGMB00912 (LA) from the stool of healthy swine and demonstrated its *in vitro* antimicrobial activity against pathogenic microorganisms, which was primarily mediated by

**Ethics approval and consent to participate**

All animal experiments were approved by the Animal Ethics Committee of the National Institute of Animal Science, Republic of Korea (approval No. NIAS 2021-503).

the production of SCFA and improvements in the intestinal microbiota [16]. Building upon these findings, the current pilot-scale study focuses on comparing the gut microbiome of weaned piglets with and without dietary supplementation with LA, aiming to assess its potential impact on the intestinal microbial community structure during the weaning period.

## MATERIALS AND METHODS

### Bacterial culture and preparation

The LA used for supplementation was isolated in our previous study [16] and inoculated into De Man-Rogosa-Sharpe broth (BD Difco™, Becton Dickinson). The culture was incubated at 37°C for 24 h, followed by centrifugation at 3,264×g for 20 min at 4°C. After centrifugation, the cell pellet was suspended in phosphate-buffered saline 1X (Gibco) and diluted to  $1.0 \times 10^8$  CFU/mL. The LA was then administered to the piglets in the LA group via oral gavage each day.

### Animal experiment and sample collection

Eight 25-day-old castrated male piglets (Landrace × Yorkshire,  $5.97 \pm 0.43$  kg) from the same herd were purchased from a commercial farm. After three days, the piglets were randomly assigned to one of two groups: piglets administered a normal diet for only 10 days (control,  $n = 4$ ), and piglets administered a normal diet daily supplemented with  $1.0 \times 10^8$  colony forming units of LA ( $n = 4$ ). The diet was prepared following the nutritional guidelines outlined in the 'Korean Feeding Standard for Pigs' (Table 1). All animal experiments were approved by the Animal Ethics Committee of the National Institute of Animal Science, Korea (approval No. NIAS 2021-503). The experiment was conducted over a total of 10 days, and on the final day (day 10), stool samples (100 g) were collected from each piglet through gentle rectal stimulation. The samples were immediately stored at  $-80^\circ\text{C}$ .

**Table 1. Ingredients and chemical composition of the experimental diet**

Items	
Ingredients	
Corn	73.74
Soybean meal 44%	22.20
Soybean oil	0.86
L-Lysine-HCl	0.17
Dicalcium phosphate	1.15
Limestone	0.88
Vitamin-mineral premix1	0.50
NaCl	0.50
Calculated composition	
Metabolizable energy (kcal/kg)	3,300
Crude protein	16.0
Lysine	0.95
Methionine	0.26
Calcium	0.66
Phosphorus	0.56

Values supplied per kilogram premix feed concentrations: Vitamin A 5,000,000 IU; vitamin E, 1,000 mg; Vitamin B<sub>1</sub>, 150 mg; Vitamin B<sub>2</sub>, 300 mg; Vitamin B<sub>12</sub>, 1,500 mg; niacin amide, 1,500 mg; DL-calcium pantothenate, 1,000 mg; folic acid, 200 mg; Vitamin H, 10 mg; choline chloride, 2,000 mg; min 3,800 mg; zinc, 1,500 mg; iron, 4,000 mg; Cu, 500 mg; I, 250 mg; Co, 100 mg; Mg, 200 mg.

until analysis.

### DNA extraction and 16S rRNA gene sequencing

The total DNA was extracted using a DNeasy PowerSoil Pro Kit (Qiagen) from 200 mg of feces collected per sample following the protocol provided by the manufacturer. DNA concentrations were measured using a Victor Nivo (PerkinElmer). A universal primer set targeting the V3–V4 regions (341F–805R) was used to prepare the 16S rRNA gene amplicons [17] using the following polymerase chain reaction (PCR) conditions for the first PCR: 3 min at 95 °C for heat activation, followed by 25 cycles of 30 s at 95 °C, 30 s at 55 °C, and 30 s at 72 °C, with a final extension of 5 min at 72 °C. The PCR product was purified with AMPure beads (Agencourt Biosciences). After purification, 10 µL of the PCR product was amplified for library construction using NexteraXT Indexed Primer. The second PCR had similar conditions, with 10 cycles. The purified product was quantified by qPCR (KAPA Library Quantification kits for Illumina Sequencing) and qualified using TapeStation D1000 ScreenTape (Agilent Technologies), then sequenced on the MiSeq™ platform (Illumina).

### 16S rRNA gene analysis

Each amplicon sequence variant (ASV) was analyzed via BLAST+ (v2.9.0) using the NCBI 16S rRNA gene database, assigning taxonomy based on the highest similarity. Hits with query coverage or identity below 85% were discarded. Multiple sequence alignments were performed with MAFFT (v7.475), and a phylogenetic tree was built using FastTreeMP (v2.1.10). Microbial community analyses were carried out through QIIME2 (v1.9) [18] using ASV abundance and taxonomy data. Species diversity and evenness in the microbial communities were calculated using the Shannon and Inverse Simpson indices. Alpha diversity was assessed via rarefaction curves and Chao1 values. Beta diversity, based on weighted and unweighted UniFrac distances, was analyzed to identify variations between comparative groups.

### Biomarker analysis

Linear discriminant analysis effect size (LEfSe) analysis was conducted to identify biomarkers with significant differential abundance across groups. The analysis was performed using the microbiomeMarker package (v1.2.1) in R (v4.0.1). Initially, the ASV table, taxonomic classifications, and sample metadata were integrated into a phyloseq object. Statistical significance for the Wilcoxon rank-sum test was set at < 0.05. Normalization was performed using the Counts Per Million method. A Kruskal-Wallis test cut-off of 0.05 was applied to detect features with significant differential abundances.

### Metabolic function prediction

To infer the metagenome's functional composition from 16S rRNA gene sequences, the analysis pipeline utilized PICRUSt2. Metadata was loaded from a tab separated values file using the readr package. The ggpicrust2 package was employed to convert the predicted metagenome abundance data into Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway abundance using the ko2kegg\_abundance function. Differential abundance analysis across treatment groups was performed using the ALDEx2 method within the ggpicrust2 package's pathway\_daa function. This method provided statistically significant pathways, adjusting for multiple testing using the Benjamini-Hochberg procedure to control the false discovery rate. The top 104 features with the lowest adjusted *p*-values were selected for further annotation. To elucidate the biological interpretation of the data, KEGG Orthology annotations for these features were obtained. The

subset of KEGG pathway abundances corresponding to the top 104 features was extracted for downstream analysis.

### Statistical analyses

Gut microbiota diversity was analyzed using QIIME2 (v1.9). Alpha diversity was assessed with Chao1, Shannon, and Simpson indices, while group differences were evaluated using the Kruskal-Wallis test. Beta diversity significance was assessed via permutational multivariate analysis of variance (PERMANOVA) with the vegan package (v2.6.4) in R, using both unweighted and weighted UniFrac distances. Statistical significance was set at  $p < 0.05$ . The LEfSe analysis employed an LDA score cut-off of  $\geq 3$  to determine the effect size, indicating biologically relevant features. The Kruskal-Wallis and Wilcoxon rank-sum tests were applied with an alpha level of 0.05. Core microbiota analysis was performed with a 20% sample prevalence and 0.2% relative abundance. Predicted metagenomic function differences were analyzed using ALDEx2. Adjusted  $p$ -values were calculated using the Benjamini-Hochberg method, with a significance threshold set at  $p < 0.05$ .

## RESULTS

### DNA sequencing data

Sequencing of the 16S rRNA genes from the fecal samples produced high-quality reads across both groups (Table 2, 3). The LA group generated 147,406–161,312 reads per sample, while the control

**Table 2.** 16S rRNA gene analysis preprocessing and quality control of ASV data

Group	Name	Total bases (bp)	Total reads	GC (%)	AT (%)	Q20 (%)	Q30 (%)
Control	Con_1	61,611,088	204,688	54.1	45.9	87.5	77.3
	Con_2	50,813,014	168,814	54.2	45.8	87.0	76.6
	Con_3	48,181,070	160,070	53.6	46.4	87.8	77.8
	Con_4	49,518,714	164,514	53.7	46.3	87.0	76.3
LA	LA_1	44,369,206	147,406	53.5	46.5	87.2	76.7
	LA_2	42,261,002	140,402	53.7	46.3	86.3	75.1
	LA_3	46,568,312	154,712	53.3	46.7	87.8	77.3
	LA_4	48,554,912	161,312	53.5	46.5	88.3	78.3

ASV, amplicon sequence variant; LA, *Lactiplantibacillus argentoratensis* strain AGMB00912; GC, guanine-cytosine; AT, adenine-thymine.

**Table 3.** Sequence processing and ASV inference summary

Group	Name	Raw data	Adapter & Primer Trimming	Pre-processing Length Trimming	Quality Filter	QC Remain (%)	denoised For	denoised Rev	merged Pair	non-chimeric	ASV Length Filter	ASV Remain (%)
Control	Con_1	102,344	101,133	101,133	81,908	80.03	77,240	79,607	61,708	43,034	43,034	42.05
	Con_2	84,407	83,292	83,292	66,928	79.29	62,539	64,756	48,555	32,749	32,749	38.80
	Con_3	80,035	79,080	79,080	64,870	81.05	60,544	62,857	47,056	30,447	30,447	38.04
	Con_4	82,257	81,257	81,257	65,302	79.39	61,111	63,156	47,936	35,872	35,872	43.61
LA	LA_1	73,703	72,530	72,530	59,330	80.50	56,612	58,043	48,824	27,261	27,259	36.98
	LA_2	70,201	69,330	69,330	54,436	77.54	52,023	53,208	43,651	29,027	29,027	41.35
	LA_3	77,356	76,436	76,436	63,280	81.80	60,930	62,150	52,756	32,045	32,045	41.43
	LA_4	80,656	79,653	79,653	67,308	83.45	64,718	66,005	56,721	34,034	34,032	42.19

ASV, amplicon sequence variant; LA, *Lactiplantibacillus argentoratensis* strain AGMB00912; QC, quality control.

group ranged from 160,070–204,688 reads. The average guanine-cytosine (GC) and adenine-thymine (AT) contents remained consistent across all samples at approximately 53.5% and 46.5%, respectively. After stringent quality control and filtering steps, the final dataset retained 523,362 high-quality, informative reads, with an average of 64,554 reads per sample.


### Alpha and beta diversity

Fig. 1 shows alpha rarefaction curves from 16S rRNA gene sequencing, illustrating species richness in fecal samples from the control and LA groups. Each curve represents a stool sample, with the X-axis showing the number of sequences and the Y-axis showing observed species richness. Stabilization of the curves suggests that the sequencing depth effectively captured microbial diversity, validating the reliability of the subsequent ASV-based analyses. The alpha diversity of bacterial communities in the fecal samples from the weaned piglets was assessed using observed features and Chao1 (indicators of species richness) along with Shannon and Simpson indices (indicating species evenness) (Fig. 2A, 2B, 2C, and 2D). The results showed that the species richness and diversity indices of the control group (observed features:  $455.50 \pm 16.09$ , Chao1:  $458.91 \pm 18.20$ , Shannon:  $6.27 \pm 0.33$ , Simpson:  $0.94 \pm 0.03$ ) were significantly higher than those of the LA treatment group (observed features:  $274.25 \pm 92.79$ , Chao1:  $275.73 \pm 13.00$ , Shannon:  $3.94 \pm 0.31$ , Simpson:  $0.72 \pm 0.04$ ) ( $p < 0.001$ ). The principal coordinates analysis (PCoA) plots, derived from both unweighted (Fig. 2E) and weighted (Fig. 2F) UniFrac distances, demonstrated significant distinctions in the microbial community separation between the control and LA groups in weaned piglets ( $p = 0.021$  and  $0.028$ , respectively). Therefore, dietary supplementation with LA significantly modulates the gut microbiota in weaning pigs.


### Probiotic supplementation modulates the microbial communities of weaned piglets

The relative abundance of the microbial populations in LA-treated weaned piglets was analyzed. At the phylum level, the LA group predominantly exhibited Bacillota (94.15%–95.86%) and Bacteroidota (0.56%–3.05%) (Fig. 3A). The control group showed a distribution with Bacillota ranging from 38.68%–54.66% and Bacteroidota from 15.83%–20.85%. Additionally, the relative abundances of Proteobacteria and Spirochaetes in the control group ranged from 2.06%–36.87% and 2.88%–34.91%, respectively. In the LA group, the distributions of these phyla were 0.11%–0.45% and 0.07%–0.28%, respectively (Fig. 3A).

*Streptococcus* and *Clostridium* exhibited a relatively high proportion in the LA group, accounting



**Fig. 1. Comparative rarefaction analysis of microbial diversity in the control and *Lactiplantibacillus argenteratensis* AGMB00912 (LA) groups.** The number of sequences was normalized to the minimum number of sequences between the control and LA groups to account for differences in sampling depth. Each curve represents an individual stool sample. The X-axis represents the number of sequences, and the Y-axis represents species.




**Fig. 2.** Alpha diversity indices in the gut microbiomes of control and *Lactiplantibacillus argenterotensis* AGMB00912 (LA) treated weaning piglets. Species richness was measured using (A) observed features and (B) Chao1 diversity indices. Species evenness was measured using (C) Shannon and (D) Simpson diversity indices. Principal coordinate analysis plots for different groups of weaning piglets. The control group (red oval) and LA group (blue oval) were significantly clustered based on unweighted (E) and weighted (F) UniFrac distance metrics.


for 45.89%–57.48% and 4.87%–11.65%, while the control group exhibited a microbial community structure characterized by a higher relative abundance of other genera and a lower proportion of *Streptococcus* and *Clostridium* (Fig. 3B). The relative abundance of *Streptococcus* in the control group ranged from 0.12%–2.57%, differing from the distribution observed in the LA group. The relative abundances of the following genera were distributed differently between the control and LA groups: *Prevotella* (4.14%–11.49% in the control group vs 0.24%–1.59% in the LA group), *Treponema* (2.03%–34.59% vs 0.07%–0.28%), *Campylobacter* (0.09%–6.51% vs 0.00%–0.20%), *Escherichia* (0.49%–34.80% vs 0.00%–0.11%), and *Intestinimonas* (0.98%–24.70% vs 0.43%–1.49%).

### Biomarker analysis of probiotic-treated weaned piglets

Fig. 4 demonstrates the significantly different taxa in the intestinal microbiota between the control and LA groups. Fig. 4A displays the LEfSe analysis cladogram, spanning from the phylum to genus levels. Fig. 4B presents a histogram displaying the species-level differences in abundance, as indicated by LDA scores > 3. These results revealed the predominant presence of Proteobacteria, Spirochaetes, and Bacteroidota in the control group at the phylum level (Fig. 3). At the genus level, the relative abundance of *Treponema*, *Campylobacter*, *Bacteroides*, and *Bullifex* was significantly higher in the control group than that in the LA group ( $p < 0.05$ ). Conversely, LA supplementation significantly increased the proportion of *Porcinola*, *Ligilactobacillus*, *Faecalitalea*, *Catenibacterium*, *Methanosphaera*, *Bifidobacterium*, *Butyrivibrio*, *Abyssovibrio*, and *Collinsella* ( $p < 0.05$ ). To visualize the varying abundance of bacterial genera in the control and LA groups, a hierarchical clustering heat map was generated (Fig. 5). The heat map revealed contrasting relative abundances between the two



**Fig. 3.** Gut microbiota composition at the phylum and genus levels in the control and *Lactiplantibacillus argenteratensis* AGMB00912 (LA) treatment groups. The relative abundance of each taxa at the phylum (A) and genus (B) levels.



**Fig. 4.** Differential abundance of bacteria between the control and *Lactiplantibacillus argenteratensis* AGMB00912 (LA) groups, as determined using the linear discriminant analysis (LDA) effect size algorithm. The cladogram shows differential abundance at the phylum, class, order, family, and genus levels (A). The histogram depicts differential abundance at the species level (B). A *p*-value of <0.05 was deemed significant in both the Kruskal–Wallis and Wilcoxon tests. A discriminative feature had a Log<sub>10</sub> LDA score of 3. The length of each histogram represents the LDA score, indicating the degree of influence of each species.




*Campylobacter*, and *Treponema* were found only in the control group, with none in the LA group (Fig. 6A and 6B). Therefore, LA supplementation may selectively inhibit pathogenic bacterial genera such as *Escherichia* and *Campylobacter*, potentially contributing to a more balanced and beneficial gut microbiota composition in weaned piglets.

### Probiotic treatment regulates the gut-related metabolic function of weaned piglets


To predict the metabolic functions of gut microbiota in weaned piglets treated with LA, PICRUSt2 was used to assess the abundance of KEGG pathways. The heat map depicted the top 22 categories of pathways impacted by the gut microbiota in each group, including cell growth and death, transport and catabolism, membrane transport, signal transduction, folding, sorting and degradation, replication and repair, translation, immune disease, infectious disease, immune system, digestive system, endocrine system, energy metabolism, glycan biosynthesis, terpenoids and polyketides, cofactors and vitamins, carbohydrate metabolism, amino acid metabolism, lipid metabolism, and xenobiotic biodegradation (Fig. 7).

### Impact of probiotic supplementation on fundamental biological processes and systemic health in weaned piglets

Fig. 8 illustrates the 24 subcategories of metabolic differences. Metabolic activities associated with cell growth and death (specifically the p53 signaling pathway and apoptosis), in addition to transport and catabolism (including lysosome and peroxisome), decreased significantly in the LA group ( $p < 0.05$ ), compared to those of the control group. While RNA degradation was activated in the control group ( $p = 0.044$ ), LA supplementation notably alleviated this effect and concurrently activated pathways related to DNA damage repair, including DNA replication, base excision repair, nucleotide excision repair, and mismatch repair. Additionally, the LA group displayed the upregulation of ABC transporter metabolism and the phosphotransferase system (membrane transport), aminoacyl-tRNA biosynthesis, and ribosome (translation), as well as the phosphatidylinositol signaling system. The LA group also exhibited downregulation of the peroxisome proliferator-activated receptors (PPAR) signaling pathway metabolism, while the



**Fig. 6. Core microbiome in different groups of weaning piglets.** Heatmap depicting the core operational taxonomic units and their prevalence at different detection thresholds in the control (A) and *Lactiplantibacillus argenteratensis* strain AGMB00912 (LA) groups (B).



**Fig. 7.** Effect of *Lactiplantibacillus argenteratensis* strain AGMB00912 (LA) on gut microbiota that regulate functional pathways in weaning piglets. Based on the analysis of 16S rRNA V3-V4 gene expression in feces, PICRUST2 software was used for functional pathway prediction. Heat map plot of varying functional pathways between the control and LA groups.


insulin signaling pathway was upregulated.

### Modulation of immune function and disease resistance

In the control group, metabolism related to bacterial infections was activated, while pathways for bacterial invasion of epithelial cells ( $p = 0.043$ ), *Escherichia coli* infection ( $p = 0.038$ ), and Shigellosis ( $p = 0.044$ ) were suppressed in the LA group (Fig. 9). Immune system pathways, including antigen processing and nucleotide-binding and oligomerization domain (NOD)-like receptor signaling, were significantly suppressed, while primary immunodeficiency pathways were activated in the LA group.


### Enhancement of nutritional metabolism

Compared to the control group, LA supplementation upregulated ( $p < 0.05$ ) carbohydrate metabolism (ascorbate and aldarate, glycolysis, pentose phosphate, fructose, mannose, galactose, starch, sucrose, amino sugar, nucleotide sugar, inositol phosphate, pyruvate, propanoate, butanoate, and C5-branched dibasic acid), as well as amino acid metabolism (glycine, serine, threonine,



**Fig. 8.** The key differences among the control and *Lactiplantibacillus argentoratensis* strain AGMB00912 (LA) groups include cell growth and death, transport and catabolism, membrane transport, signal transduction, folding, sorting and degradation, replication and repair, and translation and endocrine system pathways. ABC transporters, ATP-binding cassette transporters; RNA, ribonucleic acid; tRNA, transfer ribonucleic acid; DNA, deoxyribo nucleic acid; PPAR, peroxisome proliferator-activated receptor.

cysteine, methionine, valine, leucine, isoleucine, lysine, histidine, tyrosine, phenylalanine, and



**Fig. 9.** The key differences among the control and *Lactiplantibacillus argenteratensis* strain AGMB00912 (LA) groups include carbohydrate and amino acid metabolism pathways. NOD, nucleotide-binding and oligomerization domain.


tryptophan) (Fig. 10). In addition, lipid metabolism (fatty acid, primary and secondary bile acid, glycerolipid, linoleic acid, and sphingolipid) and cofactor and vitamin metabolism (sepiapterin reductase, thiamine, vitamin B6, nicotinate, nicotinamide, pantothenate, CoA, and folate) were also increased in LA-treated weaned piglets (Fig. 11). In energy metabolism, oxidative phosphorylation and nitrogen and sulfur metabolisms were suppressed, while photosynthesis, methane production, and carbon fixation in photosynthetic organisms were upregulated in the LA group ( $p < 0.05$ ).

### Modulation of antibiotic biosynthesis and glycan metabolic pathways

Analysis of the biosynthesis of terpenoids, polyketides, and other secondary metabolites showed that LA supplementation can significantly increase the biosynthesis of the penicillin, cephalosporin, novobiocin, streptomycin, neomycin, kanamycin, gentamicin, tetracycline, polyketide sugar, ansamycins, and vancomycin groups of antibiotics ( $p < 0.05$ ). In contrast, LA supplementation downregulated phenylpropanoid, flavone, flavonol, isoquinoline alkaloid, tropane, piperidine, and pyridine alkaloid biosynthesis (Fig. 12). Moreover, LA supplementation mitigated the degradation of other glycans ( $p = 0.044$ ), enhanced peptidoglycan biosynthesis ( $p = 0.005$ ), and significantly reduced the synthesis of glycosaminoglycans and lipopolysaccharides ( $p < 0.05$ ).

### Detoxification and enhancement of xenobiotic biodegradation

Fig. 13 predicts results for xenobiotic biodegradation and metabolism, indicating a significant reduction in the activation of degradation pathways for benzoate, bisphenol, dioxin, xylene, chloroalkane, chloroalkene, and naphthalene in the LA group ( $p < 0.05$ ). In addition, the degradation pathways for fluoroacetate and caprolactam were also significantly downregulated in




**Fig. 10.** The key differences among the control and *Lactiplantibacillus argentoratensis* strain AGMB00912 (LA) groups include cofactors and vitamins, lipid metabolism, and energy metabolism pathways.

the LA-treated piglets ( $p = 0.021$  and  $p = 0.042$ , respectively).


## DISCUSSION

During the weaning phase, piglets undergo significant changes in their intestinal microbiome due




**Fig. 11.** The key differences among the control and *Lactiplantibacillus argentoratensis* strain AGMB00912 (LA) groups include biosynthesis of other secondary metabolites, terpenoids and polyketides, and glycan biosynthesis metabolism pathways.

to separation from the mother sow and sudden dietary changes [2,19]. This study investigated the development of the gut microbiota in weaned piglets, with a focus on the effects of LA, a probiotic



**Fig. 12.** The key differences among the control and *Lactiplantibacillus argentoratensis* strain AGMB00912 (LA) groups include cofactors and vitamins, lipid metabolism, and energy metabolism pathways.

strain, on piglet intestinal microbial structure. Our previous pan-genome analysis indicated that LA supplementation produces SCFA, such as lactate, formate, and acetate—key characteristics of LA. Safety evaluations using the virulence factor database and comprehensive antibiotic resistance database confirmed the absence of virulence factors and antibiotic resistance genes, supporting LA as a safe probiotic for commercial use [16]. This study showed that dietary supplementation with LA for 10 days altered the microbial communities in weaned piglets. Alpha diversity, which measures species diversity and richness, is known to fluctuate during the post-weaning period. While some studies report an increase in alpha diversity during weaning [20], others, such as those by Hu et al. [21] and Gresse et al. [2], observed a decrease due to weaning-induced gut dysbiosis. In our study, the control LA group exhibited significantly higher alpha diversity indices (Observed, Chao1,



**Fig. 13.** The key differences among the control and *Lactiplantibacillus argentoratensis* strain AGMB00912 (LA) groups include xenobiotic biodegradation and metabolism pathways.

Shannon, and Simpson) compared to the LA-treated group. A weaning diet typically reduces the proportion of LA while promoting the proliferation of genera such as *Prevotella*, *Bacteroides*, *Bifidobacterium*, *Clostridium*, *Escherichia* and *Shigella*, leading to increased microbial diversity and imbalance [3,4]. These findings suggest that LA supplementation may help stabilize microbial diversity, preventing abrupt changes in the gut microbiome. Furthermore, beta diversity analysis revealed significant differences between the control and LA groups, with each group forming distinct microbial clusters. This pattern indicates that LA supplementation induced a unique microbial structure in the LA group, consistent with previous findings that probiotic strains like *Bifidobacterium* and *Lactobacillus* can modulate the gut microbiota in weaned piglets [22,23].

Taxonomic analysis was conducted to further explore the observed differences. The results showed a disruption in microbial balance at both the phylum and genus levels in the control group, while the LA-supplemented group maintained a more consistent microbial structure. Bacillota and Bacteroidota are the two dominant phyla in piglet gut microbiota [21], with Bacillota genera such as *Streptococcus*, *Clostridium*, and *Lactobacillus* playing key roles in producing beneficial SCFA through starch and fiber degradation [24]. The Bacillota to Bacteroidota ratio is an important marker of intestinal community balance and is linked to host health [25]. Generally, an increase in Bacteroidota and a decrease in Bacillota are associated with poor health, as these shifts can affect energy harvest and trigger inflammatory responses [26]. Our findings indicated that LA supplementation enriched Bacillota and preserved the Bacillota to Bacteroidota ratio. Similarly, Guevarra et al. [27] found that supplementation with the probiotic *Pediococcus acidilactici* modulated this ratio, supporting our results.

Additionally, The LA group had a lower relative abundance of Proteobacteria and Spirochaetes compared to the control group. Proteobacteria, which includes pathogens such as *Campylobacter*, *Escherichia*, *Helicobacter* and *Salmonella* [21], are often enriched in the intestinal microbiota of piglets suffering from post-weaning diarrhea [28,29]. An increase in Proteobacteria is a common marker of intestinal disorders. Spirochaetes, particularly the genus *Treponema*, are also known to induce colitis in infected hosts [30]. These findings suggest that LA supplementation may help mitigate microbial imbalances and support gut homeostasis during the weaning transition. Hierarchical clustering analysis further confirmed distinct microbial distributions between the control and LA groups.

To verify the statistical significance of these microbial shifts, LEfSe analysis was performed. Consistent with the taxonomy analysis results, LA treatment significantly reduced the proportion of Proteobacteria and Spirochaetes at the phylum level, and *Treponema* and *Campylobacter* at the genus level. *Campylobacter* is commonly transmitted through the fecal-oral route from sow to piglets, often causing enteritis, especially in piglets deprived of colostrum [31]. Additionally, probiotic treatments have been shown to inhibit the growth of *Treponema* in weaned piglets, as demonstrated by Zhang et al. [32], supporting the results of this study. Supplementation with LA also significantly increased the relative abundance of beneficial bacteria, including *Porcincola*, *Ligilactobacillus*, *Faecalitalea*, *Catenibacterium*, *Methanosphaera*, *Bifidobacterium*, *Butyrivibrio*, *Abysivirga*, and *Collinsella*. *Porcincola*, a gram-positive genus, contains a biosynthetic gene cluster for sactipeptide-like peptides [33], which exhibit antibacterial properties [34]. *Ligilactobacillus*, formerly part of the *Lactobacillus salivarius* group, is commonly found in fermented foods and used as a probiotic [35]. It possesses digestive enzymes, produces bacteriocins, and exhibits antioxidant activity [36]. *Faecalitalea*, a Bacillota member, produces SCFA and has positive effects on insulin secretion and responsiveness [37]. *Catenibacterium*, a gram-positive anaerobe, synthesizes acetate, lactate, butyrate, and isobutyrate from glucose [38]. *Methanosphaera*, belonging to the Archaea domain, may improve feed efficiency and reduce methane emissions [39]. Within the phylum Actinobacteria, *Bifidobacterium* predominates in healthy mammalian intestines and enhances gut health, immunity, and antioxidant activity in weaned piglets [40,41]. Pang et al. [22] also showed that *Bifidobacterium* promotes growth performance by maintaining gut homeostasis and modulating the intestinal microbiota. *Butyrivibrio*, an anaerobic butyrate-producing bacterium, was isolated from animal and human intestines [42]. *Collinsella*, capable of producing SCFAs such as acetate, formate, and lactate, also modulates bile acid and plasma cholesterol levels [43]. *Collinsella* contains genes for butyrate kinase and phosphate butyryltransferase, suggesting a specialized role in butyrate production [44]. *Abysivirga* ferments carbohydrates to enhance nutrient availability and digestibility [45]. In summary, LA supplementation played a crucial role in modulating the gut microbiota of weaned piglets by reducing the abundance of harmful microorganisms such as *Campylobacter* and *Treponema*, while enriching SCFA-producing bacteria.

The metabolic functions of microbial communities can be categorized using shotgun metagenomic sequencing. However, this method poses challenges, particularly in the presence of contamination, and is more expensive than 16S rRNA gene sequencing [46,47]. As an alternative, methods like PICRUSt2 have been developed to predict functional profiles based on taxonomic composition. PICRUSt2 predicts metabolic functions from 16S rRNA marker sequences and supports several gene family databases, including KEGG orthologs [46]. In this study, PICRUSt2 was used to predict differentially abundant metabolic functions between the control and LA groups based on 16S rRNA gene sequences. The results revealed distinct expression profiles of the top 22 pathway categories between the two groups (Fig. 7).

One notable pathway, the p53 signaling pathway, is activated in response to intestinal epithelial damage to maintain intestinal integrity. This pathway can lead to cell cycle arrest for DNA repair or trigger apoptosis [48]. Compared to the control, LA supplementation suppressed the p53 signaling pathway, reducing apoptosis and RNA degradation while upregulating DNA repair-related metabolic functions. Additionally, the PPAR- $\gamma$  signaling pathway, which regulates inflammation in the colon induced by pathogenic bacteria [49], was predicted to be significantly downregulated in the LA group. Lysosome and peroxisome metabolism were also enhanced by LA supplementation, preserving cellular integrity against pathogens [50], maintaining the balance of reactive oxygen species, and protecting cells from oxidative stress [51]. Furthermore, the antigen processing pathway, which prepares antigens for presentation to immune cells, and the NOD-like

receptor signaling pathway, which recognizes pathogenic ligands and activates immune responses [52], were modulated by LA treatment. Similar to these findings, Sun et al. [53] reported that *Lactobacillus gasseri* strain JM1 exerts immunomodulatory effects via the NOD2-mediated signaling pathway. Moreover, the bacterial invasion of epithelial cells, pathogenic *Escherichia coli* infection, and Shigellosis were suppressed in the LA group. These findings suggest that LA supplementation can improve fundamental biological processes and enhance systemic health by modulating immune function and disease resistance in weaned piglets.

The weaning transition, marked by the shift from sow milk to solid feed, significantly affects the digestive system and nutrient absorption in piglets [1,4]. In this study, dietary supplementation with LA appeared to upregulate carbohydrate metabolism across 15 distinct pathways compared to the control. Carbohydrates are essential for cellular structure and serve as the primary energy source for living organisms [54]. Carbohydrate metabolism, mediated by the gut microbiota, produces SCFA through processes such as digestion, absorption, and fermentation [55]. The increased proportion of Bacillota genera, including *Streptococcus*, *Clostridium*, and *Lactobacillus*, following LA supplementation, likely enhanced the production of beneficial SCFA via carbohydrate metabolism. The relative abundance of pathways related to amino acid, lipid, cofactor, and vitamin metabolism was significantly increased in the LA group compared to the control. Amino acids like valine, leucine, isoleucine, phenylalanine, and tyrosine are crucial SCFA precursors and fuel the tricarboxylic acid cycle [56]. Additionally, the metabolism of phenylalanine, tyrosine, and tryptophan has been linked to an increase in beneficial bacteria, which helps prevent gut inflammation [57]. Bile acids regulate glucose, lipid, and energy metabolism, while sphingolipids play a role in cell signaling and apoptosis [58]. Linoleic acid, an essential fatty acid, supports cell membrane composition, modulates inflammation, and serves as an energy source for gut epithelial cells [59]. Vitamins B6, pantothenate, and CoA are essential for cellular metabolism, with vitamin B6 playing a key role in fatty acid biosynthesis [60,61].

Intestinal bacteria produce antimicrobial peptides, either ribosomal or non-ribosomal, depending on their biosynthesis pathway [62]. In this study, LA treatment increased the biosynthesis of non-ribosomal peptides, including penicillin, cephalosporin, novobiocin, streptomycin, neomycin, kanamycin, gentamicin, tetracycline, ansamycins, and vancomycin in weaned piglets. These peptides inhibit the growth of pathogenic microorganisms. Additionally, the LA group showed a significant rise in peptidoglycan biosynthesis and a reduction in glycan degradation, indicating a thriving bacterial population that may enhance barrier protection and modulate immune responses, potentially altering the microbiota composition [63,64]. Lipopolysaccharides, from the outer membrane of gram-negative bacteria, are pathogen-associated molecular patterns made of lipid and polysaccharide components [58,65]. The findings of this study suggest that LA supplementation reduced lipopolysaccharide activation in weaned piglets. Furthermore, the activation of degradation pathways for toxic compounds—such as benzoate, bisphenol, dioxin, xylene, chloroalkane, chloroalkene, and naphthalene—was predicted in the LA group, suggesting detoxification of harmful substances in the gut [66,67]. These results indicate that LA supplementation could modulate the gut microbiome of weaned piglets by enhancing antimicrobial peptide production, strengthening gut barrier function, and reducing the risk of endotoxin and toxin exposure.

## CONCLUSION

In weaned piglets, dietary supplementation with LA modulated the gut microbiota, suggesting its potential as a valuable additive in the swine industry. Supplementation with LA promoted microbial homeostasis and enhanced metabolic functions of microbial communities, including the

biosynthesis of beneficial compounds like antibiotics, glycan structures, and SCFA. These findings suggest that LA may contribute to enhancing immune function and promoting the overall health and growth of weaned piglets. However, this was a preliminary study with a small sample size and only one method of LA administration. Therefore, further large-scale studies are needed to validate these findings and provide a more comprehensive understanding of the gut microbiome's response to LA supplementation.

## REFERENCES

- Zheng L, Duarte ME, Sevarolli Loftus A, Kim SW. Intestinal health of pigs upon weaning: challenges and nutritional intervention. *Front Vet Sci.* 2021;8:628258. <https://doi.org/10.3389/fvets.2021.628258>
- Gresse R, Chaucheyras-Durand F, Fleury MA, van de Wiele T, Forano E, Blanquet-Diot S. Gut microbiota dysbiosis in postweaning piglets: understanding the keys to health. *Trends Microbiol.* 2017;25:851-73. <https://doi.org/10.1016/j.tim.2017.05.004>
- Petri D, Hill JE, van Kessel AG. Microbial succession in the gastrointestinal tract (GIT) of the preweaned pig. *Livest Sci.* 2010;133:107-9. <https://doi.org/10.1016/j.livsci.2010.06.037>
- Bian G, Ma S, Zhu Z, Su Y, Zoetendal EG, Mackie R, et al. Age, introduction of solid feed and weaning are more important determinants of gut bacterial succession in piglets than breed and nursing mother as revealed by a reciprocal cross-fostering model. *Environ Microbiol.* 2016;18:1566-77. <https://doi.org/10.1111/1462-2920.13272>
- Cao KF, Zhang HH, Han HH, Song Y, Bai XL, Sun H. Effect of dietary protein sources on the small intestine microbiome of weaned piglets based on high-throughput sequencing. *Lett Appl Microbiol.* 2016;62:392-8. <https://doi.org/10.1111/lam.12559>
- Tian L, Bruggeman G, van den Berg M, Borewicz K, Scheurink AJW, Bruininx E, et al. Effects of pectin on fermentation characteristics, carbohydrate utilization, and microbial community composition in the gastrointestinal tract of weaning pigs. *Mol Nutr Food Res.* 2017;61:1600186. <https://doi.org/10.1002/mnfr.201600186>
- Schokker D, Zhang J, Zhang LL, Vastenhouw SA, Heilig HGHJ, Smidt H, et al. Early-life environmental variation affects intestinal microbiota and immune development in new-born piglets. *PLOS ONE.* 2014;9:e100040. <https://doi.org/10.1371/journal.pone.0100040>
- Fouhse JM, Zijlstra RT, Willing BP. The role of gut microbiota in the health and disease of pigs. *Anim Front.* 2016;6:30-6. <https://doi.org/10.2527/af.2016-0031>
- Spees AM, Wangdi T, Lopez CA, Kingsbury DD, Xavier MN, Winter SE, et al. Streptomycin-induced inflammation enhances *Escherichia coli* gut colonization through nitrate respiration. *mBio.* 2013;4:e00430-13. <https://doi.org/10.1128/mBio.00430-13>
- Bäumler AJ, Sperandio V. Interactions between the microbiota and pathogenic bacteria in the gut. *Nature.* 2016;535:85-93. <https://doi.org/10.1038/nature18849>
- Zeng MY, Inohara N, Nuñez G. Mechanisms of inflammation-driven bacterial dysbiosis in the gut. *Mucosal Immunol.* 2017;10:18-26. <https://doi.org/10.1038/mi.2016.75>
- Chávarri M, Díez-Gutiérrez L, Marañón I, Barrón LJR. Secondary metabolites from probiotic metabolism. In: Dhanasekaran D, Sankaranarayanan A, editors. *Advances in probiotics: microorganisms in food and health.* Academic Press; 2021. p. 259-76.
- Zhai L, Wu J, Lam YY, Kwan HY, Bian ZX, Wong HLX. Gut-microbial metabolites, probiotics and their roles in type 2 diabetes. *Int J Mol Sci.* 2021;22:12846. <https://doi.org/10.3390/ijms222312846>
- Zheng J, Wittouck S, Salvetti E, Franz CMAP, Harris HMB, Mattarelli P, et al. A taxonomic

- note on the genus *Lactobacillus*: description of 23 novel genera, emended description of the genus *Lactobacillus* Beijerinck 1901, and union of *Lactobacillaceae* and *Leuconostocaceae*. *Int J Syst Evol Microbiol*. 2020;70:2782-858. <https://doi.org/10.1099/ijsem.0.004107>
15. Quatravaux S, Remize F, Bryckaert E, Colavizza D, Guzzo J. Examination of *Lactobacillus plantarum* lactate metabolism side effects in relation to the modulation of aeration parameters. *J Appl Microbiol*. 2006;101:903-12. <https://doi.org/10.1111/j.1365-2672.2006.02955.x>
  16. Yoon KN, Yang J, Yeom SJ, Kim SS, Park JH, Song BS, et al. *Lactiplantibacillus argentoratensis* AGMB00912 protects weaning mice from ETEC infection and enhances gut health. *Front Microbiol*. 2024;15:1440134. <https://doi.org/10.3389/fmicb.2024.1440134>
  17. Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, et al. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res*. 2013;41:e1. <https://doi.org/10.1093/nar/gks808>
  18. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol*. 2019;37:852-7. <https://doi.org/10.1038/s41587-019-0209-9>
  19. Pereira WA, Franco SM, Reis IL, Mendonça CMN, Piazzentin ACM, Azevedo POS, et al. Beneficial effects of probiotics on the pig production cycle: an overview of clinical impacts and performance. *Vet Microbiol*. 2022;269:109431. <https://doi.org/10.1016/j.vetmic.2022.109431>
  20. Holman DB, Gzyl KE, Mou KT, Allen HK. Weaning age and its effect on the development of the swine gut microbiome and resistome. *mSystems*. 2021;6:e00682-21. <https://doi.org/10.1128/mSystems.00682-21>
  21. Hu J, Nie Y, Chen J, Zhang Y, Wang Z, Fan Q, et al. Gradual changes of gut microbiota in weaned miniature piglets. *Front Microbiol*. 2016;7:1727. <https://doi.org/10.3389/fmicb.2016.01727>
  22. Pang J, Liu Y, Kang L, Ye H, Zang J, Wang J, et al. *Bifidobacterium animalis* promotes the growth of weaning piglets by improving intestinal development, enhancing antioxidant capacity, and modulating gut microbiota. *Appl Environ Microbiol*. 2022;88:e01296-22. <https://doi.org/10.1128/aem.01296-22>
  23. Chen L, Xu Y, Chen X, Fang C, Zhao L, Chen F. The maturing development of gut microbiota in commercial piglets during the weaning transition. *Front Microbiol*. 2017;8:1688. <https://doi.org/10.3389/fmicb.2017.01688>
  24. Hu C, Xing W, Liu X, Zhang X, Li K, Liu J, et al. Effects of dietary supplementation of probiotic *Enterococcus faecium* on growth performance and gut microbiota in weaned piglets. *AMB Express*. 2019;9:33. <https://doi.org/10.1186/s13568-019-0755-z>
  25. Miragoli F, Patrone V, Prandini A, Sigolo S, Dell'Anno M, Rossi L, et al. Implications of tributyrin on gut microbiota shifts related to performances of weaning piglets. *Microorganisms*. 2021;9:584. <https://doi.org/10.3390/microorganisms9030584>
  26. Elokil AA, Magdy M, Melak S, Ishfaq H, Bhuiyan A, Cui L, et al. Faecal microbiome sequences in relation to the egg-laying performance of hens using amplicon-based metagenomic association analysis. *Animal*. 2020;14:706-15. <https://doi.org/10.1017/S1751731119002428>
  27. Guevarra RB, Kim ES, Cho JH, Song M, Cho JH, Lee JH, et al. Gut microbial shifts by synbiotic combination of *Pediococcus acidilactici* and lactulose in weaned piglets challenged with Shiga toxin-producing *Escherichia coli*. *Front Vet Sci*. 2022;9:1101869. <https://doi.org/10.3389/fvets.2022.1101869>
  28. Karasova D, Crhanova M, Babak V, Jerabek M, Brzobohaty L, Matesova Z, et al. Development of piglet gut microbiota at the time of weaning influences development of postweaning diarrhea: a field study. *Res Vet Sci*. 2021;135:59-65. <https://doi.org/10.1016/j.rvsc.2020.12.022>

29. Tran THT, Everaert N, Bindelle J. Review on the effects of potential prebiotics on controlling intestinal enteropathogens *Salmonella* and *Escherichia coli* in pig production. *J Anim Physiol Anim Nutr.* 2018;102:17-32. <https://doi.org/10.1111/jpn.12666>
30. Mølbak L, Klitgaard K, Jensen TK, Fossi M, Boye M. Identification of a novel, invasive, not-yet-cultivated *Treponema* sp. in the large intestine of pigs by PCR amplification of the 16S rRNA gene. *J Clin Microbiol.* 2006;44:4537-40. <https://doi.org/10.1128/JCM.01537-06>
31. Babakhani FK, Bradley GA, Joens LA. Newborn piglet model for campylobacteriosis. *Infect Immun.* 1993;61:3466-75. <https://doi.org/10.1128/iai.61.8.3466-3475.1993>
32. Zhang D, Ji H, Liu H, Wang S, Wang J, Wang Y. Changes in the diversity and composition of gut microbiota of weaned piglets after oral administration of *Lactobacillus* or an antibiotic. *Appl Microbiol Biotechnol.* 2016;100:10081-93. <https://doi.org/10.1007/s00253-016-7845-5>
33. Wylensek D, Hitch TCA, Riedel T, Afrizal A, Kumar N, Wortmann E, et al. A collection of bacterial isolates from the pig intestine reveals functional and taxonomic diversity. *Nat Commun.* 2020;11:6389. <https://doi.org/10.1038/s41467-020-19929-w>
34. Lee H, Churey JJ, Worobo RW. Biosynthesis and transcriptional analysis of thurincin H, a tandem repeated bacteriocin genetic locus, produced by *Bacillus thuringiensis* SF361. *FEMS Microbiol Lett.* 2009;299:205-13. <https://doi.org/10.1111/j.1574-6968.2009.01749.x>
35. Tohno M, Tanizawa Y, Sawada H, Sakamoto M, Ohkuma M, Kobayashi H. A novel species of lactic acid bacteria, *Ligilactobacillus pabuli* sp. nov., isolated from alfalfa silage. *Int J Syst Evol Microbiol.* 2022;72. <https://doi.org/10.1099/ijsem.0.005587>
36. Yang J, Shang P, Zhang B, Wang J, Du Z, Wang S, et al. Genomic and metabonomic methods reveal the probiotic functions of swine-derived *Ligilactobacillus salivarius*. *BMC Microbiol.* 2023;23:242. <https://doi.org/10.1186/s12866-023-02993-9>
37. Sanna S, van Zuydam NR, Mahajan A, Kurilshikov A, Vich Vila A, Vösa U, et al. Causal relationships among the gut microbiome, short-chain fatty acids and metabolic diseases. *Nat Genet.* 2019;51:600-5. <https://doi.org/10.1038/s41588-019-0350-x>
38. Kageyama A, Benno Y. *Catenibacterium mitsuokai* gen. nov., sp. nov., a Gram-positive anaerobic bacterium isolated from human faeces. *Int J Syst Evol Microbiol.* 2000;50:1595-9. <https://doi.org/10.1099/00207713-50-4-1595>
39. Yang J, Chen R, Peng Y, Chai J, Li Y, Deng F. The role of gut archaea in the pig gut microbiome: a mini-review. *Front Microbiol.* 2023;14:1284603. <https://doi.org/10.3389/fmicb.2023.1284603>
40. Ruiz L, Delgado S, Ruas-Madiedo P, Sánchez B, Margolles A. Bifidobacteria and their molecular communication with the immune system. *Front Microbiol.* 2017;8:2345. <https://doi.org/10.3389/fmicb.2017.02345>
41. Splichalova A, Donovan SM, Tlaskalova-Hogenova H, Stranak Z, Splichalova Z, Splichal I. Monoassociation of preterm germ-free piglets with *Bifidobacterium animalis* subsp. *lactis* BB-12 and its impact on infection with *Salmonella typhimurium*. *Biomedicines.* 2021;9:183. <https://doi.org/10.3390/biomedicines9020183>
42. Palevich N, Kelly WJ, Leahy SC, Altermann E, Rakonjac J, Attwood GT. The complete genome sequence of the rumen bacterium *Butyrivibrio hungatei* MB2003. *Stand Genomic Sci.* 2017;12:72. <https://doi.org/10.1186/s40793-017-0285-8>
43. Bag S, Ghosh TS, Das B. Complete genome sequence of *Collinsella aerofaciens* isolated from the gut of a healthy Indian subject. *Genome Announc.* 2017;5:e01361-17. <https://doi.org/10.1128/genomea.01361-17>
44. Qin P, Zou Y, Dai Y, Luo G, Zhang X, Xiao L. Characterization of a novel butyric acid-producing bacterium *Collinsella aerofaciens* subsp. *Shenzhenensis* subsp. nov. *Microorganisms.* 2019;7:78. <https://doi.org/10.3390/microorganisms7030078>

45. Schouw A, Leiknes Eide T, Stokke R, Pedersen RB, Steen IH, Bødtker G. *Abyssivirga alkaniphila* gen. nov., sp. nov., an alkane-degrading, anaerobic bacterium from a deep-sea hydrothermal vent system, and emended descriptions of *Natranaerovirga pectinivora* and *Natranaerovirga hydrolytica*. *Int J Syst Evol Microbiol*. 2016;66:1724-34. <https://doi.org/10.1099/ijsem.0.000934>
46. Douglas GM, Maffei VJ, Zaneveld JR, Yurgel SN, Brown JR, Taylor CM, et al. PICRUSt2 for prediction of metagenome functions. *Nat Biotechnol*. 2020;38:685-8. <https://doi.org/10.1038/s41587-020-0548-6>
47. Sun S, Jones RB, Fodor AA. Inference-based accuracy of metagenome prediction tools varies across sample types and functional categories. *Microbiome*. 2020;8:46. <https://doi.org/10.1186/s40168-020-00815-y>
48. Harris SL, Levine AJ. The p53 pathway: positive and negative feedback loops. *Oncogene*. 2005;24:2899-908. <https://doi.org/10.1038/sj.onc.1208615>
49. Dubuquoy L, Rousseaux C, Thuru X, Peyrin-Biroulet L, Romano O, Chavatte P, et al. PPAR $\gamma$  as a new therapeutic target in inflammatory bowel diseases. *Gut*. 2006;55:1341-9. <https://doi.org/10.1136/gut.2006.093484>
50. Barisch C, Holthuis JCM, Cosentino K. Membrane damage and repair: a thin line between life and death. *Biol Chem*. 2023;404:467-90. <https://doi.org/10.1515/hsz-2022-0321>
51. Bellec K, Cordero JB. The peroxisome: a new player in intestinal epithelial repair. *Dev Cell*. 2020;53:131-2. <https://doi.org/10.1016/j.devcel.2020.03.023>
52. Motta V, Soares F, Sun T, Philpott DJ. NOD-like receptors: versatile cytosolic sentinels. *Physiol Rev*. 2015;95:149-78. <https://doi.org/10.1152/physrev.00009.2014>
53. Sun L, Tian W, Guo X, Zhang Y, Liu X, Li X, et al. *Lactobacillus gasseri* JM1 with potential probiotic characteristics alleviates inflammatory response by activating the PI3K/Akt signaling pathway in vitro. *J Dairy Sci*. 2020;103:7851-64. <https://doi.org/10.3168/jds.2020-18187>
54. Goh YJ, Klaenhammer TR. Insights into glycogen metabolism in *Lactobacillus acidophilus*: impact on carbohydrate metabolism, stress tolerance and gut retention. *Microb Cell Fact*. 2014;13:94. <https://doi.org/10.1186/s12934-014-0094-3>
55. Portincasa P, Bonfrate L, Vacca M, De Angelis M, Farella I, Lanza E, et al. Gut microbiota and short chain fatty acids: implications in glucose homeostasis. *Int J Mol Sci*. 2022;23:1105. <https://doi.org/10.3390/ijms23031105>
56. Neis EPJG, Dejong CHC, Rensen SS. The role of microbial amino acid metabolism in host metabolism. *Nutrients*. 2015;7:2930-46. <https://doi.org/10.3390/nu7042930>
57. Liu H, Tan B, Huang B, Li J, Wang J, Liao P, et al. Involvement of calcium-sensing receptor activation in the alleviation of intestinal inflammation in a piglet model by dietary aromatic amino acid supplementation. *Br J Nutr*. 2018;120:1321-31. <https://doi.org/10.1017/S0007114518002891>
58. Lin TL, Shu CC, Chen YM, Lu JJ, Wu TS, Lai WF, et al. Like cures like: pharmacological activity of anti-inflammatory lipopolysaccharides from gut microbiome. *Front Pharmacol*. 2020; 11:554. <https://doi.org/10.3389/fphar.2020.00554>
59. Nishida A, Ando Y, Kimura I, Miyamoto J. Involvement of gut microbial metabolites derived from diet on host energy homeostasis. *Int J Mol Sci*. 2022;23:5562. <https://doi.org/10.3390/ijms23105562>
60. Theodoulou FL, Sibon OCM, Jackowski S, Gout I. Coenzyme A and its derivatives: renaissance of a textbook classic. *Biochem Soc Trans*. 2014;42:1025-32. <https://doi.org/10.1042/BST20140176>
61. Parra M, Stahl S, Hellmann H. Vitamin B $_6$  and its role in cell metabolism and physiology. *Cells*. 2018;7:84. <https://doi.org/10.3390/cells7070084>

62. Garcia-Gutierrez E, Mayer MJ, Cotter PD, Narbad A. Gut microbiota as a source of novel antimicrobials. *Gut Microbes*. 2019;10:1-21. <https://doi.org/10.1080/19490976.2018.1455790>
63. Vollmer W, Bertsche U. Murein (peptidoglycan) structure, architecture and biosynthesis in *Escherichia coli*. *Biochim Biophys Acta*. 2008;1778:1714-34. <https://doi.org/10.1016/j.bbamem.2007.06.007>
64. Irazoki O, Hernandez SB, Cava F. Peptidoglycan muropeptides: release, perception, and functions as signaling molecules. *Front Microbiol*. 2019;10:500. <https://doi.org/10.3389/fmicb.2019.00500>
65. Nativel B, Couret D, Giraud P, Meilhac O, d'Hellencourt CL, Viranaïcken W, et al. *Porphyromonas gingivalis* lipopolysaccharides act exclusively through TLR4 with a resilience between mouse and human. *Sci Rep*. 2017;7:15789. <https://doi.org/10.1038/s41598-017-16190-y>
66. Walczak-Nowicka ŁJ, Herbet M. Sodium benzoate—harmfulness and potential use in therapies for disorders related to the nervous system: a review. *Nutrients*. 2022;14:1497. <https://doi.org/10.3390/nu14071497>
67. Liu R, Cai D, Li X, Liu B, Chen J, Jiang X, et al. Effects of bisphenol A on reproductive toxicity and gut microbiota dysbiosis in male rats. *Ecotoxicol Environ Saf*. 2022;239:113623. <https://doi.org/10.1016/j.ecoenv.2022.113623>