

Comparative effects of deoxynivalenol and deepoxy-deoxynivalenol on porcine oocyte maturation and embryonic development, and the partial protective role of resveratrol


Hoyong Choi¹, Jaehyeok Yoon¹, Junghui Jo¹, Xiang Zhang¹, Kukbin Ji¹, Inchul Choi², Jeongwoong Lee^{3*}, Min Kyu Kim^{1,4*}

¹Department of Animal Science and Biotechnology, College of Agriculture and Life Sciences, Chungnam National University, Daejeon, Korea

²Department of Animal Biosystems and Dairy Science, College of Agriculture and Life Sciences, Chungnam National University, Daejeon, Korea

³Biotherapeutics Translational Research Center, Korea Research Institute of Bioscience and Biotechnology, Daejeon, Korea

⁴MK Biotech Co., LTD., Daejeon, Korea



Received: Feb 11, 2026

Revised: Mar 26, 2026

Accepted: Mar 27, 2026

*Corresponding author

Jeongwoong Lee

E-mail: jwlee@kribb.re.kr

Min Kyu Kim

E-mail: kminkyu@cnu.ac.kr

Copyright © 2026 Korean Society of Animal Science and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Hoyong Choi

<https://orcid.org/0000-0001-9527-7544>

Jaehyeok Yoon

<https://orcid.org/0009-0008-2789-7088>

Junghui Jo

<https://orcid.org/0009-0005-3814-9913>

Xiang Zhang

<https://orcid.org/0009-0006-0562-7242>

Kukbin Ji

<https://orcid.org/0000-0002-3144-9746>

Abstract

Mycotoxin contamination of animal feed is a major threat to livestock reproduction. Deoxynivalenol (DON), a trichothecene mycotoxin, disrupts cellular function by binding to ribosomes and inducing ribotoxic stress, oxidative imbalance, and apoptosis. Its primary microbial metabolite, deepoxy-deoxynivalenol (DOM-1), is thought to be less toxic. However, its reproductive effects remain unclear. DON contamination is widely reported in cereal grains used in animal feed, and pigs exposed to contaminated diets have been shown to exhibit detectable DON and DOM-1 levels in blood, urine, and fetal tissues (approximately 8–9 ng/g), highlighting the biological relevance of DON exposure during reproduction. Here, we evaluated the effects of DON (250, 500, and 1,000 ng/mL) and DOM-1 (250, 500, and 1,000 ng/mL) on porcine oocyte maturation and embryonic development *in vitro*, and examined whether resveratrol (Res, 2 μ M) could mitigate DON-induced toxicity. Exposure to DON inhibited cumulus cell expansion, maturation rates and impaired developmental competence, at a concentration of 1,000 ng/mL, developmental arrest was complete. DON-treated oocytes showed increased ROS, decreased GSH, upregulated ER stress and apoptosis-related gene levels (ATF4, XBP1, CHOP, BAX), with downregulated anti-apoptosis gene levels (BCL2). In contrast, DOM-1 had no significant effects compared with those of the controls, except for a modest reduction in the blastocyst rate at the highest concentration. Although Res did not show restorative effects on cumulus cell expansion and nuclear maturation, Res supplementation reduced DON-induced ER stress markers, increased BCL2 levels, and induced blastocyst development in DON-exposed embryos. These findings demonstrate that DON exerts dose-dependent toxicity in porcine oocytes via oxidative stress and ER stress-mediated apoptosis, whereas DOM-1 is largely nontoxic. Res partially protected against DON-induced oocyte damage and confirms its potential as a supplement to combat mycotoxins.

Keywords: Reproduction, Oocyte, Mycotoxin, Deoxynivalenol, Deepoxy-deoxynivalenol, Endoplasmic reticulum stress

Inchul Choi
<https://orcid.org/0000-0001-5011-2658>
 Jeongwoong Lee
<https://orcid.org/0000-0002-2649-9677>
 Min Kyu Kim
<https://orcid.org/0000-0001-7099-9735>

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This research was supported by a grant of Korean ARPA-H Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare, Republic of Korea (grant number: RS-2025-25454860).

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Choi H, Ji K, Kim MK.
 Data curation: Yoon J, Choi I, Lee J.
 Formal analysis: Choi H, Jo J.
 Methodology: Choi H, Zhang X, Ji K.
 Software: Choi H, Yoon J.
 Validation: Zhang X, Lee J, Kim MK.
 Investigation: Choi H, Ji K, Kim MK.
 Writing - original draft: Choi H, Choi I, Kim MK.
 Writing - review & editing: Choi H, Yoon J, Jo J, Zhang X, Ji K, Choi I, Lee J, Kim MK.

Ethics approval and consent to participate

Ethical research standards related to the use of ovaries and semen were approved by the Institutional Animal Care and Use Committee (IACUC) of Chungnam National University (Approval no: 202103A-CNU-002).

Declaration of generative AI

No AI tools were used in this article.

INTRODUCTION

Mycotoxin contamination of animal feed is a major problem affecting global livestock production. Mycotoxins are toxic secondary metabolites produced by fungal species such as *Fusarium*, *Aspergillus*, and *Penicillium*, and they remain in the feed even after processing and storage methods. Pigs have a poor ability to detoxify toxins in their bodies, which makes them particularly vulnerable to mycotoxins [1,2]. Deoxynivalenol (DON) and zearalenone (ZEN) are toxins derived from *Fusarium*, that impair feed intake, growth performance, and reproductive efficiency. Even low concentrations of DON and ZEN, these mycotoxins reduce productivity and cause reproductive disorders, resulting in economic losses [1–3]. In particular, DON is one of the most prevalent *Fusarium*-derived trichothecene mycotoxins found in cereals, such as wheat, barley, and maize. Upon ingestion, DON exerts various toxic effects, including vomiting, diarrhea, growth retardation, immunodeficiency, and intestinal damage [4–6]. At the cellular level, DON exerts its toxicity primarily by binding to the ribosomal peptidyl transferase center, thereby inhibiting protein synthesis and activating mitogen-activated protein kinase (MAPK) pathways, which lead to ribotoxic stress, oxidative damage, and immune dysfunction [7,8]. These mechanisms cause not only gastrointestinal and immune dysfunctions, but also substantial reproductive disorders in pigs, the species most sensitive to DON exposure [4]. Previous studies have shown that DON impairs ovarian function, damages follicular development, and compromises oocyte quality, thereby reducing fertilization and embryo viability in humans and pigs [9–11]. Deepoxy-deoxynivalenol (DOM-1) is the primary microbial metabolite of DON, generated through the de-epoxidation of the epoxide group at the C12–C13 position by the intestinal microbiota. Since the interaction between the epoxide group and the ribosome causes inhibition of protein synthesis, these structural changes reduce the toxicity of DON [12,13]. Consequently, DOM-1 is considered to be much less toxic than the original compound [12,14,15]. Ruminants are highly protected from DON toxicity because of their diverse gastrointestinal microbiomes that efficiently convert DON to DOM-1 in the rumen [16]. By contrast, pigs have a limited capacity to transform DON into DOM-1 and are more sensitive to DON [17]. DOM-1 has been detected in biological fluids, including plasma, urine, and follicular fluid, indicating its systemic circulation and potential interaction with reproductive tissues [18,19]. Although DOM-1 is considered a lower-toxicity metabolic, its effects on oocyte quality and embryonic development have not been precisely determined [12,14,15].

Conflicting results have been reported regarding the effects of these mycotoxins on reproductive capacity. In bovine studies, DOM-1 induced endoplasmic reticulum (ER) stress in follicular membrane cells, leading to apoptosis [18,20]. By contrast, other studies have reported that DOM-1 is less toxic than DON [12,14,15], and in some cases, no significant differences were observed between DOM-1 treated samples and controls [21]. These conflicting results may be because of differences in experimental models, concentrations, exposure duration, and assessment parameters, suggesting uncertainty regarding the effects of DOM-1. Therefore, understanding the role of DOM-1 in the reproductive context is essential to assess the true risk of DON contamination in swine production and its broader impact on animal and human health.

Given the adverse effects of DON and its metabolite DOM-1 on oocyte quality and embryonic development, it is important to develop effective protective strategies. Resveratrol (Res) is a natural polyphenolic compound found in berries is known to have antioxidant and anti-inflammatory properties. Res has been reported to mitigate DON-induced toxicity, including intestinal inflammation and oxidative stress [22–24]. In addition to its role in gut health, Res has received attention in the field of reproductive physiology. Supplementation with Res improves the quantity and quality of mitochondria, enhances oocyte competence and embryo development rates, reduces

ROS levels, and suppresses the expression of apoptosis related genes in bovine and porcine oocytes [24,25–27]. Based on these findings, the present study was designed to evaluate the impact of DON and DOM-1 on porcine oocyte maturation and embryonic development, and to determine whether Res supplementation can counteract their negative effects, thereby improving oocyte quality and developmental potential.

MATERIALS AND METHODS

This study adhered to ethical research protocols for handling porcine ovaries and semen, with all methods officially approved by the Institutional Animal Care and Use Committee (IACUC) of Chungnam National University (Approval No.: 202103A-CNU-002). All chemicals and reagents were purchased from Sigma Chemical, unless otherwise stated. DON and DOM-1 were purchased from Romer Labs.

In vitro maturation

Porcine ovaries were collected from a local slaughterhouse within 3 h of slaughter. These samples were preserved in 0.9% sterile saline at 30 °C and transported to the laboratory in a saline solution supplemented with 75 mg/mL penicillin G and 50 mg/mL streptomycin sulfate. Cumulus-oocyte complexes (COCs) were harvested from 3 to 6 mm antral follicles using a 10-mL syringe. Following the selection of COCs with at least three layers of cumulus cells, incubation was carried out at 38.5 °C in a humidified, 5% CO₂ incubator. The culture medium consisted of TCM-199 (500 µL), supplemented with 2.5 mM fructose, 0.4 mM L-cysteine, 1 mM sodium pyruvate, 0.13 mM kanamycin, 10% (v/v) pFF, 10 ng/mL EGF, 10 IU/mL PMSG, and 10 IU/mL hCG (MSD Animal Health). After an initial 22 h maturation period, the COCs were rinsed three times in hormone-free *in vitro* maturation (IVM) medium and further cultured for 20–22 h. Oocyte maturation was confirmed by identifying the extrusion of the first polar body in the perivitelline space after denuding.

Electrical parthenogenesis

After 44 h of IVM, cumulus cells were removed from COCs using 1 mg/mL hyaluronidase in Tyrode's lactate HEPES buffered medium (114 mM NaCl, 3.2 mM KCl, 2 mM NaHCO₃, 0.4 mM NaH₂PO₄, 10 mM Na-Lactate, 0.5 mM MgCl₂·6H₂O, 10 mM HEPES, 2 mM CaCl₂·2H₂O, 0.01% PVA, 12 mM sorbitol, and 0.25 mM Na-pyruvate, with pH 7.2–7.4 and osmotic pressure at 295–310 mOsm; NCSU-W). Maturity was determined by presence of the 1st polar body in the perivitelline space, and only mature oocytes (Stage metaphase II [MII]) were used in the experiments. The selected oocytes were equilibrated in an electrical activation medium containing 0.3 M mannitol, 0.05% BSA, 0.1 mM MgSO₄, and 0.1 mM CaCl₂. The oocytes were then placed in a Fusion Chamber (BTX, 45-0104) loaded with electrical activation medium; and a direct-current (DC) pulse of 1.8 kV/cm for 30 µs was applied as a single pulse, using a BTX Electro-Cell Manipulator (LF101). After the electrical pulse, the oocytes were placed in porcine zygote medium 5 (PZM-5) with 1.9 mM N-6-dimethylaminopurine (6-DMAP) for 3 h, and cultured in PZM-5 for 6 d at 38.5 °C in 5% CO₂. The cleavage rate was assessed on day 2, and embryo development was assessed on day 6.

Evaluation of cumulus cell expansion

After 44 h of maturation, the degree of cumulus cell expansion was evaluated using the method specified in previous studies [27,28]. Degrees of 0, 1, and 2, indicated the degree of expansion of

cumulus cells, respectively. All experiments were independently repeated at least three times. For each replicate, approximately 15–20 COCs were randomly allocated to each experimental group.

In vitro fertilization

After 44 h of IVM, cumulus cells were removed from COCs using 1 mg/mL hyaluronidase in NCSU-W. MII-stage oocytes denuded of cumulus cells were placed in 60- μ L droplets of modified Tris-buffered medium (mTBM) containing 113.1 mM NaCl, 3.0 mM KCl, 7.5 mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 20.0 mM Tris base, 11.0 mM glucose, 5.0 mM Na-pyruvate, 10 μ g/mL gentamicin, 1.0 mM caffeine, and 0.2% BSA, with a pH of 7.4 and an osmotic pressure of 290 to 300 mOsm. The mTBM drops were then covered with mineral oil. For insemination, sperm were separated from the diluent and buffer via centrifugation. Centrifugation was performed twice: the first at 500 $\times$ g for 5 min, and then at 300 $\times$ g for 2 min. Subsequently, sperm quality was assessed using the CASA system (MEDISCIENCE PLANNING). To achieve the optimal sperm density, 1 mL of mTBM was added to resuspend the sperm pellets. Thereafter, 20 μ L of 2×10^6 sperm/mL was added to the oocytes in 60 μ L mTBM droplets and incubated for 6 h in 5% CO_2 in air at 38.5°C. Fertilized oocytes were rinsed three times and cultured in 25 μ L of PZM-5 at 38.5°C under a 5% CO_2 atmosphere for 6 days.

Embryo development and total blastocyst counts

Cleavage and blastocyst rates were evaluated on day 2 and day 6 after *in vitro* fertilization (IVF). Cleavage and the blastocyst rates were calculated as the percentage of embryos that underwent cleavage and formed blastocysts, respectively, relative to the total number of MII oocytes following fertilization. Depending on the experiment, total blastocysts were counted, and blastocysts were visualized through immunofluorescence staining. The total blastocyst yield was determined on day 6. To evaluate blastocyst quality through immunofluorescence, blastocysts were initially washed in NCSU-W (3 times) and then subjected to Hoechst-33342 staining (5 μ g/mL) for 15 min. After the staining, each blastocyst was mounted on a slide with glycerol and a coverslip. Nuclear morphology and total cell counts were analyzed using a fluorescence microscope (Nikon).

Detection of intracellular glutathione and ROS levels

Following IVM, cumulus cells were mechanically removed to obtain denuded oocytes for assessment of intracellular ROS and GSH levels. To quantify these levels, oocytes were stained using Invitrogen CellTracker™ Blue (CMF2HC) and the Image-iT™ LIVE Green ROS Detection Kit (Invitrogen), respectively. After being rinsed three times in NCSU-W, oocytes were initially incubated with 25 μ M CellTracker™ Blue CMF2HC dye for 15 min at 38.5°C to evaluate GSH. Subsequently, after three times washes in 5% FBS-PBS (v/v), samples were further incubated for 30 min at 38.5°C with 25 μ M 5-(and 6)-carboxy-2',7'-dichlorodihydrofluorescein diacetate for ROS measurement. All oocytes were washed three times after staining, and images were captured using a fluorescence microscope equipped with appropriate filter (GSH: 370 nm; ROS: 460 nm). The average fluorescence intensity was measured by analyzing digital images of the oocytes using ImageJ software. (normalized to the average background intensity).

TUNEL staining

To assess apoptosis within blastocysts on day 6 of culture, a TUNEL assay was performed using a commercial kit (In Situ Cell Death Detection Kit, Roche Diagnostics). Initially, the embryos underwent a 40 min fixation in 4% paraformaldehyde, followed by three consecutive washes with PBS containing 1% PVA. Subsequently, blastocysts were permeabilized for 40 min with

1.0% Triton X-100. After additional washing steps, samples were treated with the TUNEL reaction mixture (enzyme label solution = 1:9) for 40 min at 38°C under dark conditions. For nuclear visualization, the blastocysts were stained with Hoechst 33342 for 15 min at 38°C, after being washed with 0.05% PBS. Finally, apoptotic signals were visualized and captured using a fluorescence microscope (Nikon Intensilight C-HGFI).

Quantitative real-time PCR

30–50 COCs were collected per group and stored at –80°C until use. Total mRNA was extracted using RNeasy Micro Kit (Qiagen) according to the manufacturer's instructions. The total RNA concentration was measured using a Biospec-Nano Spectrophotometer (Shimadzu). Complementary DNA (cDNA) was synthesized from mRNA using Maxime™ RT-PCR PreMix (iNtRON). The cDNA was amplified by quantitative real-time PCR (qRT-PCR) using a CFX96 Touch Real-Time PCR System (Bio-Rad Laboratories) and SYBR® Green Master Mix (SmartGENE™). The amplification cycles were as follows: 95°C for 5 min, followed by 39 cycles of 95°C for 10 s and 60°C for 30 s, concluding with a final melting curve from 65°C to 95°C, increasing by 0.5°C every 5 s. Primers were designed from gene sequences from NCBI. The primers used targeted ER stress; Activating Transcription Factor 4 (ATF4), X-box-binding protein-1 (XBP1), and C/EBP Homologous Protein (CHOP) and related to apoptosis; Bcl-2-associated X protein (BAX) and B-cell lymphoma 2 (BCL2), which are listed in Table 1. Relative gene expression levels of each gene were normalized using the expression of Glyceraldehyde-3-phosphate dehydrogenase (GAPDH), using the equation: $R = 2^{-[\Delta Ct_{\text{sample}} - \Delta Ct_{\text{control}}]}$.

Statistical analysis

All data were obtained from at least three independent replicates and are presented as mean ± SEM values. Statistical analyses were conducted using GraphPad Prism 10 software, version 10.4.2. Data were analyzed using one-way ANOVA with Tukey tests. *p*-values < 0.05 were considered to be statistically significant.

Experiment design

Three experiments were conducted to evaluate the effects of DON and its metabolite DOM-1 on porcine oocyte maturation and embryonic development, and to examine the potential protective

Table 1. Specific primers used for real-time PCR experiments

Primer name	Accession No.	Sequence (5' → 3')
ATF4-F	NM_001123078.1	CCCCTTGTTGGTCTGTCCG
ATF4-R		TGGCTGCTGTCTTGTCTGCT
BAX-F	XM_003127290.5	GGTCGCGCTTTTCTACTTTG
BAX-R		CGATCTCGAAGGAAGTCCAG
BCL2-F	NM_214285.1	AAACAATGCAGCAGCTGAGA
BCL2-R		AACCACCCAGCTAGAGTCA
CHOP-F	NM_001144845.1	TCTGGCTTGGCTGACTGAGGAG
CHOP-R		TTTCCGTTTCTGGGTCTTCTTTGG
XBP1-F	NM_001271738.1	GGAGTTAAGACAGCGCTTGG
XBP1-R		GAGATGTTCTGGAGGGGTGA
GAPDH-F	NM_001206359.1	GTCGGTTGTGGATCTGACCT
GAPDH-R		TTGACGAAGTGGTCGTTGAG

role of Res.

Experiment 1: Dose-dependent effects of DON and DOM-1

Preliminary experiments were performed to assess the effects on oocytes, and establish appropriate concentrations of the tested substances. Oocytes were subjected to IVM under the following treatments:

- (1) Control (CON); (2) DON 250 ng/mL; (3) DON 500 ng/mL; (4) DON 1,000 ng/mL;
- (5) DOM-1 250 ng/mL; (6) DOM-1 500 ng/mL; and (7) DOM-1 1,000 ng/mL.

After 44 h of IVM, maturation rates were assessed and developmental competence was evaluated following parthenogenetic activation by cleavage (day 2) and blastocyst formation (day 6).

Experiment 2: Mechanistic assessment of DON and DOM-1 toxicity

To investigate the mechanisms underlying mycotoxin effects, oocytes were matured *in vitro* under the following treatments (CON, DON 500–1,000 ng/mL, DOM-1 500–1,000 ng/mL). After 44 h of IVM, the following endpoints were assessed. Cumulus expansion was assessed by morphological scoring. ROS and GSH levels were measured in MII oocytes. In addition, the expression of ER stress- and apoptosis-related genes in cumulus–oocyte complexes (COCs) was analyzed using RT-qPCR. Developmental competence following IVF was evaluated by assessing cleavage rates on day 2, blastocyst formation, total cell number, and apoptosis on day 6 using the TUNEL assay.

Experiment 3: Protective effects of Res against DON-induced toxicity

To test whether Res could mitigate DON-induced damage, oocytes were matured under the following treatments:

- (1) CON; (2) Res 2 μ M; (3) DON 500 ng/mL; and (4) DON 500 ng/mL + Res 2 μ M.

The concentration of Res was selected based on previous studies [24,29]. After 44 h of IVM, maturation rates and cumulus expansion were recorded. ROS/GSH levels and ER stress- and apoptosis-related gene expression in COCs were examined. Developmental competence was then evaluated by IVF (cleavage on day 2, blastocyst formation on day 6). The concentration of resveratrol was chosen according to effective doses reported in previous studies.

RESULTS

Effects of DON and DOM-1 on porcine oocyte maturation and embryonic development *in vitro*

To evaluate the effects of DON and its metabolite DOM-1 on porcine oocyte maturation and subsequent embryonic development, oocytes were subjected to IVM in the presence of increasing concentrations of each mycotoxin, followed by parthenogenetic activation via electrostimulation (Fig. 1). DON exposure resulted in a pronounced, dose-dependent impairment of oocyte maturation and subsequent embryonic development (Table 2). Even at a moderate concentration, DON significantly reduced the oocyte maturation rate (35.00%, $p < 0.05$) compared with that of control group (91.00%, $p < 0.05$). In addition, DON markedly compromised blastocyst formation (10.00%, $p < 0.05$) compared to the control group (39.93%, $p < 0.05$). At the highest concentration tested, oocyte maturation was almost completely inhibited, and no embryonic cleavage or blastocyst




Fig. 1. Results of parthenogenesis experiments according to concentration of mycotoxin treatment. The experiment was replicated at least five times. Within each category, groups marked with different letters are significantly different ($p < 0.05$).

Table 2. Results of parthenogenesis experiment according to concentration of mycotoxin treatment

	Concentration (ng)	Number of oocyte examined	Percentage of oocyte maturation (n)	Number of embryo examined	Percentage of embryo cleavage (n)	Percentage of blastocyst (n)
CON	-	120	91.00 ± 1.5 (109) ^a	109	94.67 ± 4.0 (106) ^a	39.93 ± 5.8 (42) ^a
DON	250	103	77.33 ± 1.3 (80) ^a	80	99.03 ± 1.6 (76) ^a	36.83 ± 1.9 (29) ^a
	500	99	35.00 ± 4.0 (36) ^b	34	83.90 ± 8.6 (30) ^a	10.00 ± 9.23 (4) ^b
	1,000	114	4.67 ± 2.9 (6) ^c	6	6.67 ± 6.7 (3) ^b	0 ^c
DOM-1	250	141	85.33 ± 3.2 (120) ^a	120	99.43 ± 2.9 (109) ^a	42.53 ± 7.2 (50) ^a
	500	122	86.00 ± 1.0 (105) ^a	102	93.07 ± 3.7 (95) ^a	42.73 ± 9.3 (43) ^a
	1,000	111	80.33 ± 5.9 (89) ^a	87	91.83 ± 6.3 (80) ^a	23.74 ± 5.6 (20) ^a

Within each category, groups marked with different letters are significantly different ($p < 0.05$).

CON, control; DON, deoxynivalenol; DOM-1, deepoxy-deoxynivalenol.

development was observed, indicating severe toxicity of DON toward both nuclear maturation and developmental competence. By contrast, as shown in Table 2, DOM-1 exposure did not significantly affect oocyte maturation or early embryonic cleavage across the tested concentration ranges, and both parameters remained comparable to those of the control group ($p < 0.05$). However, blastocyst development was adversely affected at the highest concentration of DOM-1 (23.00%), whereas lower concentrations supported blastocyst formation at levels similar to those of the control (39.93%, $p < 0.05$). These findings suggest that DOM-1 exerts substantially weaker reproductive toxicity than DON, with detrimental effects on embryonic development observed only at elevated concentrations. Based on these results, because a significant difference emerged at concentrations ≥ 500 ng/mL, the treatment range for subsequent experiments was defined as 500–1,000 ng/mL.

Effect of DON and DOM-1 on cumulus expansion and oxidative stress in porcine COCs

To evaluate the effects of DON and DOM-1 on cumulus expansion, porcine COCs were cultured in a maturation medium supplemented with the respective treatments. After 44 h of IVM, cumulus expansion was observed under a microscope and scored according to established criteria (Fig. 2). The results showed that COCs in both DON-treated groups had significantly reduced cumulus expansion compared with that of the control, with the 1,000 ng/mL DON group displaying a more pronounced reduction than the 500 ng/mL group. By contrast, DOM-1 treatment did not significantly affect cumulus expansion, and the COCs displayed morphology similar to that of the control group. These findings suggest that, unlike DON, DOM-1 has minimal detrimental effects




Fig. 2. Effect of DON and DOM-1 on cumulus expansion of porcine COCs during *in vitro* maturation. (A) The degree of cumulus cell expansion was classified into three grades: Degree 0 (no cumulus expansion), Degree 1 (moderate expansion), and Degree 2 (full expansion). (B) Average the degree of cumulus cell expansion. Within each category, groups marked with different letters are significantly different ($p < 0.05$). Scale bar = 100 μm. COCs, cumulus-oocyte complexes; DON, deoxynivalenol; DOM-1, deepoxy-deoxynivalenol.

on cumulus cell function during oocyte maturation. To further determine whether these differences in expansion were associated with changes in oxidative stress, intracellular ROS and GSH levels were examined in MII-stage oocytes (Fig. 3). ROS levels were higher in both DON-treated groups than in the control and DOM-1 groups, with the highest levels observed in the 1,000 ng/mL DON group. By contrast, GSH levels were markedly lower in the DON groups than in the control and DOM-1 groups. The DOM-1 groups had slightly lower GSH than the control, but the difference was small. These findings show that DON causes oxidative stress in porcine oocytes by increasing ROS levels and reducing GSH levels, whereas DOM-1 has only a mild effect.

Effect of DON and DOM-1 on pro- and anti-apoptotic gene regulation in porcine COCs

To confirm whether DON and DOM-1 induce apoptosis in porcine COCs during IVM, gene expression analysis was performed using RT-qPCR (Fig. 4). For ER stress-related genes, the expression of activating transcription factor 4 (ATF4) and X-box binding protein 1 (XBP1) was significantly higher in the DON-treated groups, compared with the control, and both genes showed a dose-dependent increase with rising DON concentrations ($p < 0.05$). The expression of C/EBP homologous protein (CHOP/DDIT3) was significantly elevated only in the 1,000 ng/mL DON group compared with all other groups ($p < 0.05$). For apoptosis-related genes, the expression of BCL2-associated X protein (BAX) was significantly increased in the DON groups compared with the control, with the 1,000 ng/mL DON group showing the highest expression and significantly exceeding the DOM-1 group ($p < 0.05$). By contrast, the anti-apoptotic gene B-cell lymphoma 2 (BCL2) was markedly reduced in the DON groups. The 1,000 ng/mL DON group showed the lowest BCL2 expression compared with all other groups ($p < 0.05$), whereas the 500 ng/mL DON




Fig. 3. The effects of DON and DOM-1 on intracellular ROS and GSH levels in porcine oocytes. (A) Oocytes were stained with CellTracker Blue (a–e) and 2', 7'-dichlorodihydrofluorescein diacetate (H₂DCFDA) (f–j) to detect intracellular levels of GSH and ROS, respectively. (B) Quantification of the intracellular GSH and ROS levels in MII oocytes. Data are presented as mean ± SEM. For each treatment group, a total of 30–40 oocytes were utilized. Different letters above bars indicate significant differences ($p < 0.05$). Scale bar = 100 μm. DON, deoxynivalenol; DOM-1, deepoxy-deoxynivalenol; GSH, glutathione.




Fig. 4. The expression levels of genes related to (A–C) ER stress and (D–E) apoptosis in COCs cultured under different concentrations of mycotoxins. COCs were cultured for 2 days. The experiment was replicated at least three times. Within each category, groups marked with different letters are significantly different ($p < 0.05$). ER, endoplasmic reticulum; COCs, cumulus-oocyte complexes.

group also showed significantly lower expression than that of both the control and the 500 ng/mL DOM-1 group ($p < 0.05$).

Effect of DON and DOM-1 on porcine preimplantation development

Because parthenogenetic activation evaluates only the intrinsic developmental competence of oocytes, an additional IVF experiment was conducted to assess fertilization capacity and subsequent embryonic development under conditions closer to physiological reproduction.

Cleavage rates were evaluated 2 d after IVF (Fig. 5A). The cleavage rate in the 500 ng/mL DON group was not significantly different from those in the control and DOM-1 groups ($p > 0.05$). However, no cleaved embryos were observed in the 1,000 ng/mL DON group. Both DOM-1 groups showed cleavage rates comparable to the control ($p > 0.05$).

On day 6, blastocyst formation was observed (Fig. 5B). The blastocyst rate was significantly reduced in the DON groups compared with that in control group, and in the 1,000 ng/mL DON group, no embryos developed to the blastocyst stage. By contrast, both DOM-1 groups exhibited blastocyst formation rates similar to those of the control group ($p > 0.05$). Detailed results are presented in Table 3.

Blastocyst quality was assessed by Hoechst staining (Fig. 6A). Total cell numbers were significantly lower in the 500 ng/mL DON group than in the control and DOM-1 500 ng/mL groups ($p < 0.05$). No expanded blastocysts were available for staining in the 1,000 ng/mL DON group.

TUNEL staining was performed to determined apoptosis levels (Fig. 6B). Blastocysts derived




Fig. 5. The effects of DON and DOM-1 on development. (A) The cleavage rate was observed on 2 days of culture and (B) the blastocyst rate on 6 days after IVF. The experiment was replicated at least five times. Within each category, groups marked with different letters are significantly different ($p < 0.05$). DON, deoxynivalenol; DOM-1, deepoxy-deoxynivalenol; IVF, *in vitro* fertilization.

Table 3. Effects of DON and DOM-1 on development

	Concentration (ng)	Number of embryo examined	Percentage of embryo cleavage (n)	Percentage of blastocyst (n)
CON	-	228	94.92 ± 1.7 (216) ^a	18.92 ± 1.6 (43) ^a
DON	500	111	91.20 ± 3.5 (108) ^a	6.4 ± 2.6 (7) ^b
	1,000	4	0 ^b	0 ^c
DOM-1	500	175	88.06 ± 2.8 (157) ^a	18.44 ± 2.7 (35) ^a
	1,000	179	91.18 ± 1.3 (179) ^a	16.78 ± 1.8 (33) ^a

Within each category, groups marked with different letters are significantly different ($p < 0.05$).

CON, control; DON, deoxynivalenol; DOM-1, deepoxy-deoxynivalenol.

from the 500 ng/mL DON group contained significantly more TUNEL-positive cells than those from the control and DOM-1 groups ($p < 0.05$). No significant differences were observed between the control and DOM-1 groups. No expanded blastocysts were obtained from the 1,000 ng/mL DON group for TUNEL analysis.

Protective effects of Res on porcine oocytes exposed to DON

To investigate whether Res could mitigate the detrimental effects of DON on porcine oocytes, we examined its effects on oocyte maturation, apoptosis-related gene expression, and subsequent embryonic development.

When oocytes were cultured with 2 μ M Res, 500 ng/mL DON, or 500 ng/mL DON plus 2 μ M Res (DON + Res), no significant differences in maturation rate or cumulus expansion were observed between the control and Res groups. By contrast, both the DON and DON + Res groups showed significantly impaired oocyte maturation and reduced cumulus expansion compared with the control and Res-only groups. Moreover, Res supplementation did not restore the maturation or




Fig. 6. The Effects of DON and DOM-1 on embryo development. Embryos were cultured for 6 days after IVF. (A) Total cell number of porcine blastocysts. (B) Hoechst staining and TUNEL assay in porcine blastocysts. The experiment was replicated at least three times. For each treatment group, a total of 30–40 oocytes were utilized. Within each category, groups marked with different letters are significantly different ($p < 0.05$). Scale bar = 100 µm. DON, deoxynivalenol; DOM-1, deepoxy-deoxynivalenol; IVF, *in vitro* fertilization.

cumulus expansion of DON-treated COCs (Fig. 7). To further assess the protective role of Res, the expression of ER stress- and apoptosis-related genes was examined using RT-qPCR (Fig. 8). As expected, the DON group exhibited significantly elevated expression of ATF4, XBP1, and CHOP compared with the Control and Res groups ($p < 0.05$). Importantly, Res co-treatment (DON + Res) significantly reduced ATF4 and XBP1 expression relative to DON alone, although CHOP expression remained elevated. For apoptosis-




Fig. 7. The effect of resveratrol on cumulus cell expansion and oocytes maturation in DON-exposed oocyte. (A) Oocyte maturation on the 2 days of culture. (B) Average the degree of cumulus cell expansion. (C) The degree of cumulus cell expansion was classified into three grades: Degree 0 (no cumulus expansion), Degree 1 (moderate expansion), and Degree 2 (full expansion). Within each category, groups marked with different letters are significantly different ($p < 0.05$). Scale bar = 100 µm. DON, deoxynivalenol.




Fig. 8. The expression levels of genes related to (A–C) ER stress and (D–E) apoptosis in COCs cultured with Res and DON. COCs were cultured for 2 days. The experiment was replicated at least three times. Within each category, groups marked with different letters are significantly different ($p < 0.05$). ER, endoplasmic reticulum; COCs, Cumulus-oocyte complexes; DON, deoxynivalenol.

related markers, BAX expression was significantly higher in the DON group than in the control and Res groups, whereas expression in the DON + Res group did not differ significantly from that in the DON group. By contrast, BCL2 expression was significantly reduced in the DON group, whereas its levels in the Res and DON + Res groups were comparable to those in the control group ($p > 0.05$). These data indicate that Res alleviates DON-induced ER stress but only partially mitigates pro-apoptotic signaling.

Finally, the effects of Res on the developmental potential of DON-treated oocytes were evaluated using IVF (Fig. 9). On day 2, cleavage rates did not differ significantly among the groups ($p > 0.05$; Fig. 8A). However, by day 6, the DON group failed to produce blastocysts, whereas the DON + Res group produced embryos that developed to the blastocyst stage. Although the blastocyst rate in the DON + Res group was lower than that in the control and Res groups ($p < 0.05$), the presence of blastocysts indicates a partial protective effect of Res against DON-induced developmental arrest (Fig. 9B). Detailed results are presented in Table 4.

DISCUSSION

This study demonstrated that DON exerts profound toxicity on porcine oocytes, whereas its metabolite DOM-1 has minimal detrimental effects. DON exposure impairs cumulus expansion, reduces maturation rates, induced oxidative and ER stress, triggers apoptosis, and ultimately compromises developmental competence, by contrast DOM-1 exhibits minimal or no detrimental effects under the




Fig. 9. The effects of resveratrol on oocytes development in DON-exposed oocytes. (A) The cleavage rate was observed on 2 days of culture and (B) the blastocyst rate on 6 days after IVF. The experiment was replicated at least five times. Within each category, groups marked with different letters are significantly different ($p < 0.05$). DON, deoxynivalenol; IVF, *in vitro* fertilization.

Table 4. Effects of resveratrol on oocytes development in DON-exposed oocytes

	Concentration	Number of oocyte examined	Percentage of oocyte maturation (n)	Number of embryo examined	Percentage of embryo cleavage (n)	Percentage of blastocyst (n)
CON	-	172	85.50 ± 2.3 (147) ^a	142	91.23 ± 1.6 (130) ^a	14.30 ± 1.8 (20) ^a
Resveratrol	2 µM	209	87.00 ± 2.1 (183) ^a	169	92.24 ± 3.0 (154) ^a	13.82 ± 1.7 (19) ^a
DON	500 ng	166	45.40 ± 5.0 (75) ^b	70	80.24 ± 5.6 (57) ^b	0 ^b
DON + Resveratrol	500 ng + 2 µM	157	50.40 ± 3.1 (76) ^b	74	79.20 ± 4.6 (59) ^a	5.17 ± 3.3 (4) ^b

Within each category, groups marked with different letters are significantly different ($p < 0.05$).

CON, control; DON, deoxynivalenol; DOM-1, deepoxy-deoxynivalenol.

same conditions. Furthermore, Res supplementation partially alleviated DON-induced stress responses and increased the proportion of cleaved embryos that developed into blastocysts, highlighting its potential as a protective agent against mycotoxin-induced reproductive toxicity.

Interestingly, our findings suggest that DON-induced developmental arrest may involve two distinct windows of sensitivity in porcine oocytes and embryos. The first window occurs during oocyte maturation, when DON disrupts cumulus expansion, increases oxidative and ER stress, and shifts apoptotic gene expression profiles, thereby reducing oocyte competence prior to fertilization. The second window occurs during zygotic genome activation (ZGA), which occurs after the 4-cell stage [30]. At this stage, embryonic development becomes highly dependent on *de novo* transcription and protein synthesis. Given that DON exerts its toxicity by binding to the ribosomal peptidyl transferase center and inhibiting protein synthesis [31,32], it is plausible that DON-exposed zygotes, even if they reach the 4-cell stage, fail to progress further because of impaired transcriptional activation and translational capacity. This dual impact—first at the maturation stage and later during ZGA—provides a mechanistic explanation for the severe reduction in blastocyst formation observed in our study. By contrast, DOM-1, which lacks the epoxide moiety required for ribosomal binding [33,34], did not impair ZGA-dependent development, which is consistent with its minimal cytotoxicity in oocytes. In this study, DON exposure was limited to the oocyte

maturation period, therefore, the observed effects were directly attributed to alterations in oocyte quality. Although impaired oocyte maturation may subsequently influence early embryonic development, including stages around the maternal–zygotic transition, this process was not directly examined in the current study. Thus, any potential involvement of DON in the maternal–zygotic transition should be interpreted with caution and warrants further investigation.

During IVF, oocytes rely predominantly on stored maternal transcripts and post-transcriptional control, with minimal *de novo* transcription; whereas, cumulus cells remain transcriptionally active and exchange metabolites and redox equivalents with the oocyte via gap junctions in transzonal projections. Thus, ROS and GSH depletion detected in MII oocytes likely reflect disrupted cumulus–oocyte metabolic coupling rather than primary transcriptional changes within the oocyte itself [35]. Cumulus cells are key suppliers of cysteine, pyruvate, and NADPH, and help sustain intra-oocyte GSH, which is a principal antioxidant for meiotic competence. Perturbation of cumulus function is therefore expected to reduce oocyte GSH levels and elevate ROS levels, ultimately compromising maturation quality. Our results showing higher ROS levels and lower GSH levels in the DON groups are consistent with this biology and with the literature linking cumulus redox support to oocyte competence [36–38]. To localize the transcriptional response, we assayed COCs and found that DON activated ER-stress-related markers (ATF4, XBP1, and CHOP) and shifted the apoptotic balance toward cell death, with significantly increased BAX expression and markedly decreased BCL2 expression. These changes align with the known actions of DON, which trigger ribotoxic and ER stress pathways, and apoptosis in porcine reproductive cells and embryos [39]. Conversely, DOM-1, a deep-epoxidized metabolite, showed minimal changes in ER stress or apoptosis markers in COCs and had a negligible impact on ROS levels in oocytes, consistent with its markedly reduced cytotoxicity relative to DON. Together, these findings support a model in which DON first impairs cumulus-mediated redox support during maturation (resulting in increased ROS and decreased GSH in the oocyte without changes in oocyte gene-expression), and secondarily drives ER stress-linked apoptosis at the cumulus–oocyte complex level, culminating in poor cleavage and blastocyst outcomes.

The present study examined the potential of Res to counteract the toxic effects of DON on porcine oocytes. Res was selected because of its well-documented antioxidant and anti-apoptotic properties, as previous studies have shown that it reduces intracellular ROS, enhances GSH levels, and improves oocyte competence in mammalian reproduction [25,40]. Previous studies have demonstrated that dietary Res supplementation during gestation and lactation improves the antioxidant status of both sows and piglets, accompanied by modulation of antioxidant-related gene expression and activation of the Kelch-like ECH-associated protein 1 (KEAP1)–NRF2 signaling pathway in the placenta [41]. Considering that measurable levels of DON have been detected in the body fluids of sows and their fetuses that consumed DON-contaminated feed in previous studies [42], we hypothesized that supplementation of the oocyte culture medium with Res would exert antioxidant effects, thereby alleviating DON-induced toxicity and improving cumulus cell function and oocyte maturation. Although Res supplementation did not restore cumulus expansion or nuclear maturation impaired by DON, it exerted beneficial effects at the molecular level within COCs. Specifically, Res can attenuate the DON-induced upregulation of ER stress markers such as ATF4 and XBP1 and stabilize the expression of the anti-apoptotic gene BCL2 [29]. These results suggest that Res partially relieves ER stress in cumulus cells, thereby maintaining a more favorable microenvironment for the enclosed oocytes.

At the developmental level, the protective effects of Res were more apparent. While cleavage rates in the DON + Res group showed only a modest, non-significant trend toward improvement compared with the DON group, blastocyst development was partially restored. Notably, embryos

derived from DON + Res oocytes progressed beyond the cleavage stage and reached the blastocyst stage, which was completely blocked in the DON-only group. This finding suggests that although Res cannot fully rescue the maturation process oocytes that succeed in maturation under DON + Res conditions may retain high developmental competence, possibly supported by improved cumulus-mediated signaling or antioxidant defense during maturation.

Collectively, these findings highlight Res as a potential mitigator of DON-induced damage in porcine oocytes, particularly through modulation of ER stress and partial restoration of developmental potential. Nevertheless, the inability of Res to fully restore cumulus expansion, nuclear maturation, and blastocyst rates to control levels underscores the need for additional strategies, possibly involving combined antioxidant or signaling-targeted interventions, to comprehensively overcome the reproductive toxicity of DON.

REFERENCES

1. Awuchi CG, Ondari EN, Nwozo S, Odongo GA, Eseoghene IJ, Twinomuhwezi H, et al. Mycotoxins' toxicological mechanisms involving humans, livestock and their associated health concerns: a review. *Toxins*. 2022;14:167. <https://doi.org/10.3390/toxins14030167>
2. Kępińska-Pacelik J, Biel W. Alimentary risk of mycotoxins for humans and animals. *Toxins*. 2021;13:822. <https://doi.org/10.3390/toxins13110822>
3. Pierron A, Alassane-Kpembi I, Oswald IP. Impact of mycotoxin on immune response and consequences for pig health. *Anim Nutr*. 2016;2:63-8. <https://doi.org/10.1016/j.aninu.2016.03.001>
4. McGraw MS, Daigneault BW. Environment to embryo: intersections of contaminant exposure and preimplantation embryo development in agricultural animals. *Biol Reprod*. 2022;107:869-80. <https://doi.org/10.1093/biolre/ioac121>
5. Alizadeh A, Braber S, Akbari P, Kraneveld A, Garssen J, Fink-Gremmels J. Deoxynivalenol and its modified forms: are there major differences? *Toxins*. 2016;8:334. <https://doi.org/10.3390/toxins8110334>
6. Holanda DM, Kim SW. Mycotoxin occurrence, toxicity, and detoxifying agents in pig production with an emphasis on deoxynivalenol. *Toxins*. 2021;13:171. <https://doi.org/10.3390/toxins13020171>
7. Pierron A, Alassane-Kpembi I, Oswald IP. Impact of two mycotoxins deoxynivalenol and fumonisin on pig intestinal health. *Porcine Health Manag*. 2016;2:21. <https://doi.org/10.1186/s40813-016-0041-2>
8. Sweeney T. Is exposure to endocrine disrupting compounds during fetal/post-natal development affecting the reproductive potential of farm animals? *Domest Anim Endocrinol*. 2002;23:203-9. [https://doi.org/10.1016/S0739-7240\(02\)00157-1](https://doi.org/10.1016/S0739-7240(02)00157-1)
9. Gerez JR, Desto SS, Bracarense APFRL. Deoxynivalenol induces toxic effects in the ovaries of pigs: an ex vivo approach. *Theriogenology*. 2017;90:94-100. <https://doi.org/10.1016/j.theriogenology.2016.10.023>
10. Malekinejad H, Schoevers EJ, Daemen IJMM, Zijlstra C, Colenbrander B, Fink-Gremmels J, et al. Exposure of oocytes to the Fusarium toxins zearalenone and deoxynivalenol causes aneuploidy and abnormal embryo development in pigs. *Biol Reprod*. 2007;77:840-7. <https://doi.org/10.1095/biolreprod.107.062711>
11. Han J, Wang QC, Zhu CC, Liu J, Zhang Y, Cui XS, et al. Deoxynivalenol exposure induces autophagy/apoptosis and epigenetic modification changes during porcine oocyte maturation. *Toxicol Appl Pharmacol*. 2016;300:70-6. <https://doi.org/10.1016/j.taap.2016.03.006>
12. Pierron A, Mimoun S, Murate LS, Loiseau N, Lippi Y, Bracarense APFL, et al. Microbial biotransformation of DON: molecular basis for reduced toxicity. *Sci Rep*. 2016;6:29105.

- <https://doi.org/10.1038/srep29105>
13. Hooft JM, Bureau DP. Deoxynivalenol: mechanisms of action and its effects on various terrestrial and aquatic species. *Food Chem Toxicol.* 2021;157:112616. <https://doi.org/10.1016/j.fct.2021.112616>
 14. Bracarense APFL, Pierron A, Pinton P, Gerez JR, Schatzmayr G, Moll WD, et al. Reduced toxicity of 3-epi-deoxynivalenol and de-epoxy-deoxynivalenol through deoxynivalenol bacterial biotransformation: in vivo analysis in piglets. *Food Chem Toxicol.* 2020;140:111241. <https://doi.org/10.1016/j.fct.2020.111241>
 15. Tassis PD, Reisinger N, Nagl V, Tzika E, Schatzmayr D, Mittas N, et al. Comparative effects of deoxynivalenol, zearalenone and its modified forms de-epoxy-deoxynivalenol and hydrolyzed zearalenone on boar semen in vitro. *Toxins.* 2022;14:497. <https://doi.org/10.3390/toxins14070497>
 16. Gallo A, Mosconi M, Trevisi E, Santos RR. Adverse effects of Fusarium toxins in ruminants: a review of in vivo and in vitro studies. *Dairy.* 2022;3:474-99. <https://doi.org/10.3390/dairy3030035>
 17. Pan T, Guo R, Wang W, Liu X, Xia B, Jiang L, et al. Mechanism of mitigating on Deoxynivalenol-induced intestinal toxicity in swine and its dietary regulation strategy. *J Integr Agric.* 2025;24:2449-64. <https://doi.org/10.1016/j.jia.2024.07.037>
 18. Guerrero-Netro HM, Estienne A, Chorfi Y, Price CA. The mycotoxin metabolite deepoxy-deoxynivalenol increases apoptosis and decreases steroidogenesis in bovine ovarian theca cells. *Biol Reprod.* 2017;97:746-57. <https://doi.org/10.1093/biolre/iox127>
 19. Guerrero-Netro HM, Barreta MH, Costa E, Goetten A, Dupras R, Mills L, et al. Effects of the mycotoxin metabolite de-epoxy-deoxynivalenol (DOM-1) on embryo development and sperm motility in cattle. *J Appl Toxicol.* 2021;41:1180-7. <https://doi.org/10.1002/jat.4102>
 20. Reyes-Perea AD, Guerrero-Netro HM, Meza-Serrano E, Estienne A, Price CA. The mycotoxin de-epoxy-deoxynivalenol (DOM-1) increases endoplasmic reticulum stress in ovarian theca cells. *Toxins.* 2023;15:228. <https://doi.org/10.3390/toxins15030228>
 21. Springler A, Hessenberger S, Reisinger N, Kern C, Nagl V, Schatzmayr G, et al. Deoxynivalenol and its metabolite deepoxy-deoxynivalenol: multi-parameter analysis for the evaluation of cytotoxicity and cellular effects. *Mycotoxin Res.* 2017;33:25-37. <https://doi.org/10.1007/s12550-016-0260-z>
 22. Lee K, Wang C, Chaille JM, Machaty Z. Effect of resveratrol on the development of porcine embryos produced in vitro. *J Reprod Dev.* 2010;56:330-5. <https://doi.org/10.1262/jrd.09-174K>
 23. Wang F, Tian X, Zhang L, He C, Ji P, Li Y, et al. Beneficial effect of resveratrol on bovine oocyte maturation and subsequent embryonic development after in vitro fertilization. *Fertil Steril.* 2014;101:577-86. <https://doi.org/10.1016/j.fertnstert.2013.10.041>
 24. Qiu Y, Nie X, Yang J, Wang L, Zhu C, Yang X, et al. Effect of resveratrol supplementation on intestinal oxidative stress, immunity and gut microbiota in weaned piglets challenged with deoxynivalenol. *Antioxidants.* 2022;11:1775. <https://doi.org/10.3390/antiox11091775>
 25. Kwak SS, Cheong SA, Jeon Y, Lee E, Choi KC, Jeung EB, et al. The effects of resveratrol on porcine oocyte in vitro maturation and subsequent embryonic development after parthenogenetic activation and in vitro fertilization. *Theriogenology.* 2012;78:86-101. <https://doi.org/10.1016/j.theriogenology.2012.01.024>
 27. Lee S, Jin JX, Taweechaipaisankul A, Kim GA, Lee BC. Synergistic effects of resveratrol and melatonin on in vitro maturation of porcine oocytes and subsequent embryo development. *Theriogenology.* 2018;114:191-8. <https://doi.org/10.1016/j.theriogenology.2018.03.040>
 28. Giarretta E, Spinaci M, Bucci D, Tamanini C, Galeati G. Effects of resveratrol on vitrified porcine oocytes. *Oxid Med Cell Longev.* 2013;2013:920257. <https://doi.org/10.1155/2013/920257>

29. Oestrup O, Hall V, Petkov SG, Wolf XA, Hyldig S, Hyttel P. From zygote to implantation: morphological and molecular dynamics during embryo development in the pig. *Reprod Domest Anim.* 2009;44 Suppl 3:39-49. <https://doi.org/10.1111/j.1439-0531.2009.01482.x>
30. Wang FM, Galson DL, Roodman GD, Ouyang H. Resveratrol triggers the pro-apoptotic endoplasmic reticulum stress response and represses pro-survival XBP1 signaling in human multiple myeloma cells. *Exp Hematol.* 2011;39:999-1006. <https://doi.org/10.1016/j.exphem.2011.06.007>
31. Jukam D, Shariati SAM, Skotheim JM. Zygotic genome activation in vertebrates. *Dev Cell.* 2017;42:316-32. <https://doi.org/10.1016/j.devcel.2017.07.026>
32. Pestka JJ, Smolinski AT. Deoxynivalenol: toxicology and potential effects on humans. *J Toxicol Environ Health B Crit Rev.* 2005;8:39-69. <https://doi.org/10.1080/10937400590889458>
33. Bae HK, Pestka JJ. Deoxynivalenol induces p38 interaction with the ribosome in monocytes and macrophages. *Toxicol Sci.* 2008;105:59-66. <https://doi.org/10.1093/toxsci/kfn102>
34. Vanhoutte I, De Mets L, De Boevre M, Uka V, Di Mavungu JD, De Saeger S, et al. Microbial detoxification of deoxynivalenol (DON), assessed via a *Lemna minor* L. Bioassay, through biotransformation to 3-epi-DON and 3-epi-DOM-1. *Toxins.* 2017;9:63. <https://doi.org/10.3390/toxins9020063>
35. Yao Y, Long M. The biological detoxification of deoxynivalenol: a review. *Food Chem Toxicol.* 2020;145:111649. <https://doi.org/10.1016/j.fct.2020.111649>
36. Hu Y, Zhang R, Zhang S, Ji Y, Zhou Q, Leng L, et al. Transcriptomic profiles reveal the characteristics of oocytes and cumulus cells at GV, MI, and MII in follicles before ovulation. *J Ovarian Res.* 2023;16:225. <https://doi.org/10.1186/s13048-023-01291-2>
37. Sutton-McDowall ML, Gilchrist RB, Thompson JG. The pivotal role of glucose metabolism in determining oocyte developmental competence. *Reproduction.* 2010;139:685-95. <https://doi.org/10.1530/REP-09-0345>
38. von Mengden L, Klamt F, Smitz J. Redox biology of human cumulus cells: basic concepts, impact on oocyte quality, and potential clinical use. *Antioxid Redox Signal.* 2020;32:522-35. <https://doi.org/10.1089/ars.2019.7984>
39. Liu Y, Xiao X, Wang L, Fu Y, Yao S, Liu X, et al. The dose-dependent dual effects of alpha-ketoglutarate (AKG) on cumulus oocyte complexes during in vitro maturation. *Cell Commun Signal.* 2024;22:472. <https://doi.org/10.1186/s12964-024-01827-z>
40. Chen Q, Cui Y, Zhao J, Zeng W, Jin N, Yang L, et al. Cellular apoptosis induced by deoxynivalenol. *Indian J Microbiol.* 2022;62:61-9. <https://doi.org/10.1007/s12088-021-00965-z>
41. González-Garzón AC, Ramón-Ugalde JP, Ambríz-García DA, Vazquez-Avendaño JR, Hernández-Pichardo JE, Rodríguez-Suastegui JL, et al. Resveratrol reduces ROS by increasing GSH in vitrified sheep embryos. *Animals.* 2023;13:3602. <https://doi.org/10.3390/ani13233602>
42. Meng Q, Li J, Wang C, Shan A. Biological function of resveratrol and its application in animal production: a review. *J Anim Sci Biotechnol.* 2023;14:25. <https://doi.org/10.1186/s40104-022-00822-z>
43. Dänicke S, Brüssow KP, Goyarts T, Valenta H, Ueberschär KH, Tiemann U. On the transfer of the *Fusarium* toxins deoxynivalenol (DON) and zearalenone (ZON) from the sow to the full-term piglet during the last third of gestation. *Food Chem Toxicol.* 2007;45:1565-74. <https://doi.org/10.1016/j.fct.2007.02.016>