

Dietary supplementation with coated zinc oxide enhanced growth performance by modulating systemic immunity and microbial composition in weaned pigs

Jinuk Nam^{1#}, Kyeong Il Park^{2#}, Jinho Cho^{3#}, Hyeun Bum Kim^{4#}, Jinmu Ahn², Youngmin Park⁵, Sheena Kim⁴, Sangwoo Park⁶, Yanhong Liu⁵, Hyunjin Kyoung^{2,5*}, Minho Song^{2*}

¹Department of Nutrition, University of California, Davis, CA, USA


²Department of Animal Science and Biotechnology, Chungnam National University, Daejeon, Korea

³Department of Animal Science, Chungbuk National University, Cheongju, Korea

⁴Department of Animal Biotechnology, Dankook University, Cheonan, Korea

⁵Department of Animal Science, University of California, Davis, CA, USA

⁶Division of Animal Bioscience and Integrated Biotechnology, Gyeongsang National University, Jinju, Korea



Received: Mar 18, 2025

Revised: Apr 7, 2025

Accepted: Apr 7, 2025

#These authors contributed equally to this work.

*Corresponding author

Hyunjin Kyoung

E-mail: hjkyoung@cnu.ac.kr

Minho Song

E-mail: mhsong@cnu.ac.kr

Copyright © 2026 Korean Society of Animal Science and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Jinuk Nam

<https://orcid.org/0009-0003-8696-5697>

Kyeong Il Park

<https://orcid.org/0000-0002-3590-3993>

Jinho Cho

<https://orcid.org/0000-0001-7151-0778>

Abstract

Dietary zinc oxide (ZnO) is widely adopted in the swine industry to alleviate post-weaning diarrhea and promote the growth performance of weaned pigs. However, its low rate of absorption in the gastrointestinal tract leads to excessive excretion, which causes severe environmental pollution. To overcome this limitation, dietary coated ZnO (CZO) has been studied to enhance zinc absorption and mitigate its environmental influence. In the present study, we investigated the effects of dietary CZO on growth performance, frequency of diarrhea, nutrient digestibility, hematological and immune responses, and fecal microbiota of weaned pigs. A total of 72 weaned pigs (7.30 ± 0.01 kg of average initial body weight; 28 days old) were randomly assigned to three dietary treatments (4 pigs/pen; 6 replicates/treatment): basal weaner diet (CON), CON with 2,500 ppm of ZnO (HZO), and CON with 200 mg/kg of CZO. The experiment was conducted for 6 weeks. The CZO group increased ($p < 0.05$) average daily gain and gain to feed ratio and showed tendency to reduce ($p = 0.094$) the frequency of diarrhea compared with the CON group during the first two weeks after weaning. The CZO group showed a tendency for the lowest levels of serum tumor necrosis factor- α ($p = 0.072$) on day 7 and hematocrit ($p = 0.083$) on day 14 and the highest ($p = 0.072$) levels of the immunoglobulin M on day 14 among the treatment groups. The HZO group tended to have the lowest ($p = 0.078$) serum cortisol levels on day 7 among all treatments. At the genus level of gut microbiota analysis, the dietary CZO group showed a higher ($p < 0.05$) relative abundance of *Prevotella*, *Eubacterium*, and *Lactobacillus* than pigs treated with the CON diet. The HZO group showed a higher ($p < 0.05$) relative abundance of *Eubacterium* and *Lactobacillus* than the CON group. Our results demonstrated that dietary CZO supplementation enhanced growth

Hyeun Bum Kim
<https://orcid.org/0000-0003-1366-6090>
 Jinmu Ahn
<https://orcid.org/0009-0005-1490-2974>
 Youngmin Park
<https://orcid.org/0009-0003-7522-1211>
 Sheena Kim
<https://orcid.org/0000-0002-5410-1347>
 Sangwoo Park
<https://orcid.org/0000-0003-2288-1374>
 Yanhong Liu
<https://orcid.org/0000-0001-7727-4796>
 Hyunjin Kyoung
<https://orcid.org/0000-0001-5742-5374>
 Minho Song
<https://orcid.org/0000-0002-4515-5212>

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This research was supported by Chungnam National University.

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Nam J, Park KI, Cho J, Kim HB, Kyoung H, Song M.
 Data curation: Nam J, Park KI, Cho J, Kim HB, Kyoung H, Song M.
 Formal analysis: Nam J, Ahn J, Park Y, Kim S, Park S, Liu Y.
 Methodology: Nam J, Park KI, Cho J, Kim HB.
 Software: Nam J, Park KI, Cho J, Kim HB.
 Validation: Park KI, Cho J, Kim HB, Kyoung H, Song M.
 Investigation: Nam J, Park KI, Cho J, Kim HB.
 Writing - original draft: Nam J, Park KI, Cho J, Kim HB.
 Writing - review & editing: Nam J, Park KI, Cho J, Kim HB, Ahn J, Park Y, Kim S, Park S, Liu Y, Kyoung H, Song M.

Ethics approval and consent to participate

The experimental protocol used in this experiment was reviewed and approved by the Institutional Animal Care and Use Committee of Chungnam National University, Daejeon, Republic of Korea (approval# 202103A-CNU-080).

Declaration of generative AI

No AI tools were used in this article.

performance and reduced the incidence of diarrhea by modulating systemic immune responses and shifting fecal microbial compositions in weaned pigs, suggesting its potential as an alternative to the high dose ZnO diets.

Keywords: Diarrhea, Fecal microbiota, Growth performance, Systemic immunity, Zinc

INTRODUCTION

Weaning is considered a critical and challenging stage for piglets because of their immature intestinal system and underdeveloped immune functions [1,2]. During this stage, piglets are exposed to an increased risk of post-weaning diarrhea (PWD), which is a fatal condition that hampers the development of piglets and impairing intestinal barrier function by disrupting intestinal mucosa and tight junction proteins, thereby reducing feed intake and causing severe growth retardation [3,4]. Dietary zinc is considered an essential trace element for animals primarily to support immunity, growth, and overall health [5,6]. In swine production, zinc oxide (ZnO) is commonly used at high doses (2,000 to 3,000 ppm) and many previous studies have shown that dietary ZnO has been supplemented in nursery diets to alleviate PWD by supporting intestinal development while reducing gut permeability [7]. However, long-term supplementation with high levels of ZnO raised some serious concerns. It increased the resistance of antibacterial activity by selecting specific strains such as *Escherichia coli* that are more metal tolerant, accumulates toxicity in the kidney and liver, and leads to substantial environmental pollution when excreted through swine manure, owing to their low absorption rates [8–10]. To address these issues, the European Union has prohibited the use of high levels of ZnO in the feed and other countries have begun to phase out the use of ZnO to address environmental concerns [11]. Therefore, various studies are being conducted to control dietary ZnO doses and enhance the bioavailability of zinc, aiming to reduce the reliance on high levels of ZnO in weaner diets [12–14].

ZnO is highly soluble in acidic environments, causing it to be broken down into zinc ions by gastric acid, resulting in a relatively small amount of ZnO reaching the small intestine [15,16]. In addition, it can bind to substances such as phytate to form insoluble complexes, further reducing its bioavailability [17]. Enteric coating technology acts as a protective barrier and has been developed to enhance the delivery of ZnO into the small intestine. With the application of the coating, the inner core of ZnO reduced the dissociation by gastric acid, allowing it to reach the small intestine where it can effectively perform its function and be released by pancreatic lipase [13,18]. In an *in vitro* study, Shen et al. [15] reported that the dissociation percentage under acidic conditions was higher for uncoated ZnO than for coated ZnO. Moreover, previous studies showed that dietary inclusion of coated ZnO improved growth rate, nutrient digestibility, and alleviated PWD when compared with a basal nursery diet and achieved similar effects to those observed with pharmacological levels of ZnO [13,18,19]. However, the effects of lipid-coated ZnO still remain controversial, and relatively few studies have examined its effect on the changes of fecal microbiota in the weaning piglets. Therefore, our study aimed to investigate the effects of dietary supplementation with coated ZnO as an alternative to high doses of conventional ZnO on growth performance, nutrient digestibility, hematological parameters, systemic immune responses, and fecal microbiota of weaned pigs.

MATERIALS AND METHODS

Origins of the tested product

The lipid encapsulated ZnO used in the present study was supplied by (Animal Care Company).

According to the protocol from manufacturer, this product is a double-coated form of zinc derived from ZnO in the form of lipid matrix composed of fatty acids and hydrogenated palm oil. The ZnO content of the product was 51%.

Experimental design, animals, and diets

A total of 72 weaned piglets ([Landrace × Yorkshire] × Duroc) with average initial body weight (BW) of 7.30 ± 0.01 kg around 28 days age were conducted for 6 weeks. Pigs were randomly assigned to three dietary treatments (4 pigs/pen; 6 replicates/treatment; block = initial BW) with pen as the experimental unit. The dietary treatments were 1) a basal weaner diet based on corn-soybean meal (CON), 2) CON supplemented with 2,500 ppm high levels of conventional ZnO (HZO), and 3) CON supplemented with 200 mg/kg dietary coated ZnO (CZO) for 6 weeks. The basal diet was designed to meet or exceed the nutritional requirements of the National Research Council [20] for weaned pigs (Table 1). The environments of the pig room were controlled, with the temperature set between 28°C to 32°C and allowing *ad libitum* access to feed and water in the pens of the same sized (232 × 175 cm; width × length) during the trial.

Table 1. Composition of basal diet for weaned pigs (as-fed basis)

Item	Basal diet
Ingredient (%)	
Corn	49.86
Soybean meal (44%)	25.00
Whey powder	12.50
Soy protein concentrate	6.25
Soybean oil	3.00
Limestone	1.14
Monocalcium phosphate	1.05
Vitamin premix ¹⁾	0.20
Mineral premix ²⁾	0.20
L-Lysine-HCl	0.45
DL-Methionine	0.16
L-Threonine	0.13
L-Valine	0.06
Total	100.00
Calculated energy and nutrient	
Metabolizable energy (kcal/kg)	3,465
Crude protein (%)	21.26
Calcium (%)	0.81
Phosphorous (%)	0.65
Lysine (%)	1.53
Methionine (%)	0.47
Threonine (%)	0.95
Tryptophan (%)	0.25

¹⁾Vitamin premix provided the following quantities of vitamin per kilogram of complete diet: vitamin A, 12,000 IU; vitamin D₃, 2,500 IU; vitamin E, 30 IU; vitamin K₃, 3 mg; D-pantothenic acid, 15 mg; nicotinic acid, 40 mg; choline, 400 mg; and vitamin B₁₂, 12 µg.

²⁾Mineral premix provided the following quantities of mineral per kilogram of complete diet: Fe, 90 mg from iron sulfate; Cu, 8.8 mg from copper sulfate; Zn, 100 mg from zinc oxide; Mn, 54 mg from manganese oxide; I, 0.35 mg from potassium iodide; Se, 0.30 mg from sodium selenite.

Data and sample collection

For measuring growth performance, individual BW and feed residual amount were weighed and noted on days 1, 14, and 42 and the average daily gain (ADG), average daily feed intake (ADFI), and gain to feed ratio (G:F) were calculated. Fecal scores in each pen were visually monitored by two independent observers during the first two weeks. The fecal scores ranged from 1 to 5, with 1 indicating hard and dry feces, 2 indicating soft feces, 3 indicating moist feces, 4 indicating mild diarrhea, and 5 indicating watery diarrhea. Diarrhea percentages were determined by recording the number of days each pens received a score of 4 or higher, and these days were expressed as a percentage. In the last week of the trial, 0.2% of chromium oxide (Daejung Chemicals & Metals) was added to the diets for the each pen as an indigestible marker and feces were collected during the last 3 days of the experiment via rectal palpation after 4 days of adaptation and stored at -80°C for apparent total tract digestibility (ATTD) analysis. In fecal microbiota analysis, feces were obtained from three randomly selected pigs per treatment on the first and final day of the experiment and stored at -80°C until analysis. Blood samples (10 mL) were collected from the jugular vein of one randomly chosen pig per pen using with or without ethylenediaminetetraacetic acid (EDTA) as the anticoagulant vacutainer tubes (BD Vacutainer Systems) on days 1, 7, and 14. Serum samples were collected from the blood samples in non-EDTA tubes which were allowed to clot at room temperature and centrifuged for 15 min at $3,000\times g$ at 4°C and stored at -80°C for immune response analysis.

Nutrient digestibility analysis

Feed and frozen fecal samples were dried in a forced-air dry oven at 65°C for 72 hours. After drying, these samples were ground using a electric grinder (Hamilton Beach) for chemical analysis. All grounded samples were analyzed for dry matter (DM), and energy using a bomb calorimeter (Parr 1281 Bomb Calorimeter, Parr Instrument), and crude protein (CP) using the Kjeldahl method according to the procedures outlined by the Association of Official Analytical Chemists [21,22]. Chromium concentrations in the diets and fecal samples were estimated using an absorption spectrophotometer (Hitachi Z-5000 Absorption Spectrophotometer, Hitachi High-Tech). The ATTD values for DM, CP, and energy in each treatment were determined based on a previous study [23].

Blood samples analysis

Hematological parameters were measured from blood samples collected in EDTA tubes using the Scil Vet abc hematology analyzer (Scil Animal Care Company) adjusted for porcine blood. Measurement included total white blood cell (WBC) count, red blood cell (RBC) count, and hematocrit (HCT) level. Systemic immune responses were analyzed by measuring serum concentrations of tumor necrosis factor- α (TNF- α) and cortisol using ELISA kits (R&D Systems), as well as immunoglobulin A (IgA), immunoglobulin G (IgG), and immunoglobulin M (IgM) using ELISA kits (Bethyl Laboratories). Absorbance was measured at 450 nm using a microplate reader and the concentration of each sample was determined based on their standard curve.

Fecal microbiota analysis

The total DNA was extracted from the use of 200 mg of feces from each sample using the QIAamp Fast DNA Stool Mini Kit (QIAGEN) according to the manufacturer's guidelines. DNA concentrations were then measured with the Colibri Microvolume Spectrometer (Titertek Berthold). Samples with the ratio of OD₂₆₀/OD₂₈₀ ranging from 1.80 to 2.15 were processed additionally. To amplify the V5 to V6 regions of the 16S rRNA gene, the sets of polymerase chain reaction (PCR) primers 799F-mod6 and 1114R were used [24]. After amplification, the products were purified using a Wizard® SV Gel

and PCR Clean Up System purification kit (Promega). Purified amplicons of the 16S rRNA gene were sequenced using an Illumina MiSeq platform (Macrogen). Raw sequence data produced using the Illumina MiSeq platform were subjected to quality assessment using FastQC. Sequencing errors that were inconsistent with the PCR primers contained ambiguous bases, or were less than 200 bp in length were eliminated using Mothur software [25]. The QIIME2 and Microbial Helper pipelines performed de novo operational taxonomic unit (OTU) selection using the Deblur algorithm with a 97% identity cutoff. Alpha (observed OTUs, Chao1, Shannon, and Simpson) and Beta diversity based on principal coordinate analysis (PCoA) using Bray-Curtis indices were measured to assess microbial richness and evenness and comparing microbial communities within each dietary treatment as well as comparing microbial communities across different dietary treatments, respectively.

Statistical analyses

All data except the frequency of diarrhea and fecal microbiota were analyzed using PROC GLM of SAS (ver. 9.4, SAS Institute) in a randomized complete block design (block = initial BW). The experiment unit was the pen. The statistical models for growth performance, digestibility, hematological parameters, and immune responses of pigs included diet as the main effect and initial BW as a covariate. The Chi-square test was used to analyze the frequency of diarrhea. The MicrobiomeAnalyst (<https://www.microbiomeanalyst.ca/>) webtool was used to analyze the alpha and beta diversity statistics using the Kruskal-Wallis test and PERMANOVA, respectively. Taxonomic classification was analyzed by STAMP software ver. 2.1.3 [26] using a two-sided Welch's *t*-test. Results were presented as means $\pm$ SEM, except alpha diversity presented as means $\pm$ SD. Statistically differences and tendency were considered at $p < 0.05$ and $0.05 \leq p < 0.10$, respectively.

RESULTS

Growth performance and nutrient digestibility

During the first two weeks of the study, the pigs fed CZO exhibited a significant increase ($p < 0.05$) in ADG and G:F compared with those fed CON (Table 2). However, there were no differences in the growth performance from day 15 to 42 or in the overall period among the dietary treatments. The frequency of diarrhea was measured at 13.87% in CON, 11.05% in HZO, and 10.16% in the CZO group. The pigs in the CZO group tended to have a lower ($p = 0.094$) incidence of diarrhea frequency than those in the CON group. No differences were observed in the ATTD of DM, CP, and energy among the dietary treatments (Table 3).

Hematological parameters and immune responses

No significant differences were observed in WBC and RBC counts on days 1, 7, and 14 among the dietary treatments (Table 4). However, pigs fed with CZO tended to have the lowest HCT level on day 14 ($p = 0.083$). Regarding systemic immunity, pigs in the CZO group showed a tendency toward the lowest serum TNF- α level on day 7 ($p = 0.072$) and the highest IgM level on day 14 ($p = 0.072$) among the dietary treatments. In addition, pigs in the HZO group tended to have the lowest serum cortisol concentration on day 7 ($p = 0.078$) compared to other treatments. No significant differences were observed in IgG and IgA levels among the dietary treatments.

Fecal microbiota diversities

The alpha diversity indices are presented in Table 5. There were no differences in alpha diversity indices among the dietary treatments on day 1. However, pigs fed dietary CZO showed an increase ($p < 0.05$)

Table 2. Effects of dietary coated ZnO on growth performance of weaned pigs¹⁾

Item ²⁾	CON	HZO	CZO	SEM	p-value
Day 1 to 14					
Initial BW (kg)	7.29	7.30	7.29	0.32	0.995
Final BW (kg)	10.65	11.12	11.75	0.41	0.673
ADG (g/d)	240 ^b	273 ^{ab}	319 ^a	20.02	0.045
ADFI (g/d)	435	428	424	25.26	0.963
G:F (g/g)	0.552 ^b	0.638 ^{ab}	0.752 ^a	0.042	0.021
Day 15 to 42					
Initial BW (kg)	10.65	11.12	11.75	0.41	0.673
Final BW (kg)	21.75	22.15	22.45	0.34	0.576
ADG (g/d)	396	394	382	12.75	0.953
ADFI (g/d)	958	954	957	30.45	0.988
G:F (g/g)	0.413	0.413	0.399	0.025	0.981
Day 1 to 42					
Initial BW (kg)	7.29	7.30	7.29	0.32	0.995
Final BW (kg)	21.75	22.15	22.45	0.34	0.576
ADG (g/d)	344	354	361	7.12	0.482
ADFI (g/d)	784	779	779	23.21	0.962
G:F (g/g)	0.439	0.454	0.463	0.017	0.738
Frequency of diarrhea ³⁾ (%)	13.87	11.05	10.16		0.094

¹⁾Each value is the mean of 6 replicates (4 pigs/pen).

²⁾CON, basal weaner diet based on corn and soybean meal; HZO, CON + 2,500 ppm ZnO; CZO, CON + 200 mg/kg dietary coated ZnO; BW, body weight; ADG, average daily gain; ADFI, average daily feed intake; G:F, gain to feed ratio.

³⁾Frequency of diarrhea for the first 2 weeks after weaning (%) = (number of diarrhea score of 4 or higher / number of pen days) × 100. Data was analyzed using the Chi-square test.

^{a,b)}Means in the same row with different superscripts are different ($p < 0.05$).

Table 3. Effects of dietary coated ZnO on nutrient and energy apparent total tract digestibility of weaned pigs¹⁾

Item ²⁾	CON	HZO	CZO	SEM	p-value
DM (%)	72.45	75.05	77.01	5.45	0.413
Energy (%)	74.26	76.63	78.03	5.19	0.569
CP (%)	71.48	73.58	73.55	6.28	0.638

¹⁾Each value is the mean of 6 replicates (1 pig/pen).

²⁾CON, basal weaner diet based on corn and soybean meal; HZO, CON + 2,500 ppm ZnO; CZO, CON + 200 mg/kg dietary coated ZnO; DM, dry matter; CP, crude protein.

in the observed OTUs and Chao1 indices on day 42 compared with those fed HZO. Microbial beta diversity among the dietary treatments was visualized using PCoA plots as shown in 2D and 3D (Fig. 1). No differences were found ($r^2 = 0.24$, $p > 0.05$) in the Bray–Curtis distance on day 1 among the dietary treatments. However, differences in the clustering of microbial communities were detected ($r^2 = 0.43$, $p < 0.05$; Figs. 1C and 1D) on day 42 among the dietary treatments.

Fecal bacterial taxonomic relative abundances

The relative abundances of bacterial taxa at the phylum and genus levels among the dietary treatments are presented in Figs. 2 and 3, respectively. In terms of fecal microbial composition, Firmicutes (CON, 57.42%; HZO, 50.48%; CZO, 58.26%) was the most dominant phylum among the dietary treatments on day 1, followed by Bacteroidetes (CON, 25.46%; HZO, 23.99%; CZO,

Table 4. Effects of dietary coated ZnO on blood profile and immune responses of weaned pigs¹⁾

Item ²⁾	CON	HZO	CZO	SEM	p-value
WBC ($\times 10^3/\mu\text{L}$)					
Day 1	13.54	13.18	14.60	2.59	0.585
Day 7	23.15	19.81	20.08	1.31	0.429
Day 14	24.65	23.27	18.80	2.48	0.511
RBC ($\times 10^6/\mu\text{L}$)					
Day 1	4.65	4.89	4.68	0.23	0.405
Day 7	6.05	6.28	6.29	0.41	0.894
Day 14	6.75	6.45	6.40	0.32	0.475
HCT (%)					
Day 1	26.08	25.02	25.42	0.82	0.835
Day 7	31.55	28.63	29.54	1.85	0.775
Day 14	36.67 ^a	31.05 ^{ab}	30.25 ^b	1.91	0.083
TNF- α (pg/mL)					
Day 1	140.10	132.64	133.35	25.52	0.515
Day 7	109.01 ^a	80.29 ^b	78.57 ^b	9.51	0.072
Day 14	105.44	84.55	85.91	38.91	0.507
Cortisol (ng/mL)					
Day 1	115.05	118.74	116.72	18.64	0.691
Day 7	123.43 ^a	81.18 ^b	86.54 ^{ab}	12.52	0.078
Day 14	105.61	98.53	97.53	17.45	0.303
IgG (mg/mL)					
Day 1	5.26	4.95	5.01	0.76	0.776
Day 7	3.67	3.79	3.83	1.12	0.675
Day 14	3.75	4.52	4.26	0.31	0.268
IgM (mg/mL)					
Day 1	1.47	1.68	1.23	0.32	0.514
Day 7	1.46	1.48	1.47	0.22	0.638
Day 14	1.22 ^b	1.61 ^{ab}	1.65 ^a	0.13	0.072
IgA (mg/mL)					
Day 1	0.28	0.31	0.21	0.07	0.496
Day 7	0.33	0.31	0.30	0.10	0.728
Day 14	0.39	0.54	0.44	0.13	0.464

¹⁾Each value is the mean of 6 replicates (1 pig/pen).²⁾CON, basal weaner diet based on corn and soybean meal; HZO, CON + 2,500 ppm ZnO; CZO, CON + 200 mg/kg dietary coated ZnO; WBC, white blood cell; RBC, red blood cell; HCT, hematocrit; TNF- α , tumor necrosis factor- α ; IgG, immunoglobulin G; IgM, immunoglobulin M; IgA, immunoglobulin A.^{a,b}Means in the same row with different superscripts are different ($p < 0.10$).


23.08%). On day 42, Firmicutes remained the most dominant phylum among the dietary treatments (CON, 69.37%; HZO, 66.14%; CZO, 70.81%). In contrast, Bacteroidetes abundance was decreased (CON, 6.18%; HZO, 18.25%; CZO, 11.11%) and Actinobacteria was increased (CON, 1.52%; HZO, 14.33%; CZO, 2.97%). *Prevotella* was the most dominant fecal genus among the dietary treatments (CON, 24.87%; HZO, 32.67%; CZO, 27.36%) on day 1. On day 42, the dietary CZO group showed a higher ($p < 0.05$) proportions of *Prevotella* (18.62% vs. 7.94%), *Eubacterium* (11.97% vs. 1.32%) and *Lactobacillus* (9.58% vs. 5.13%) than the CON group (Fig. 4). In addition, pigs fed CZO had a higher ($p < 0.005$) abundance of *Eubacterium* (11.97% vs. 1.42%) than those fed HZO.

Table 5. Effects of dietary coated ZnO on bacterial alpha diversity of weaned pigs¹⁾

Item ²⁾	CON	HZO	CZO	p-value
Day 1				
Observed OTUs	492.33 ± 10.30	504.33 ± 122.10	423.00 ± 68.74	0.697
Chao1	494.25 ± 104.20	506.62 ± 123.02	424.81 ± 68.50	0.699
Shannon	4.46 ± 0.35	4.40 ± 0.79	4.06 ± 0.54	0.769
Simpson	0.972 ± 0.010	0.940 ± 0.063	0.943 ± 0.041	0.732
Day 42				
Observed OTUs	417.67 ^{ab} ± 75.21	300.67 ^b ± 20.07	462.33 ^a ± 9.88	0.033
Chao1	420.06 ^{ab} ± 76.18	302.16 ^b ± 19.62	466.43 ^a ± 12.41	0.032
Shannon	3.77 ± 0.35	3.43 ± 0.46	4.09 ± 0.18	0.397
Simpson	0.946 ± 0.016	0.950 ± 0.017	0.952 ± 0.006	0.750

¹⁾Each value is the mean of 3 replicates and presented as mean ± SD.

²⁾CON, basal weaner diet based on corn and soybean meal; HZO, CON + 2,500 ppm ZnO; CZO, CON + 200 mg/kg dietary coated ZnO; OTUs, operational taxonomic units.

^{a,b)}Means in the same row with different superscripts are different ($p < 0.05$).

Fig. 1. Principal coordinates analysis (PCoA) based on Bray-Curtis distance of bacterial communities of weaned pigs ($n = 3$). Permutational multivariate analysis of variance (PERMANOVA) was used for statistical differences of clustering distances. Beta diversity analysis were represented for fecal bacteria (A), (C) at day 1 and day 42 for ordination 2D and (B), (D) at day 1 and day 42 for PCoA 3D. CON, basal weaner diet based on corn and soybean meal; HZO, CON + 2,500 ppm ZnO; CZO, CON + 200 mg/kg dietary coated ZnO.

The dietary HZO group had higher ($p < 0.05$) *Lactobacillus* abundance (19.67% vs. 5.13% and 9.58%, respectively) than the CON and CZO groups. Additionally, pigs fed HZO had increased ($p < 0.05$) the proportions of *Prevotella* (14.46% vs. 7.94%) compared with those fed CON.




Fig. 2. The relative abundance of the fecal microbiota at the phylum level among dietary treatments on day 1 and day 42. The proportions of top five bacteria are presented at the phylum level in each time period. CON, basal weaner diet based on corn and soybean meal; HZO, CON + 2,500 ppm ZnO; CZO, CON + 200 mg/kg dietary coated ZnO.




Fig. 3. The relative abundance of the fecal microbiota at the genus level among dietary treatments on day 1 and day 42. CON, basal weaner diet based on corn and soybean meal; HZO, CON + 2,500 ppm ZnO; CZO, CON + 200 mg/kg dietary coated ZnO.

DISCUSSION

Due to weaning stress, piglets often suffer from physiological disorders including digestive and immune dysfunction, which reduce feed efficiency, impair nutrient absorption, and delay growth rate [27,28]. To mitigate these challenges and prevent susceptibility to infections, pharmacological levels of ZnO are commonly supplemented in the weaner diets [29–31]. This supplementation not only promotes growth performance but also ameliorates the intestinal barrier environment, enhances systemic immune function, and modulates gut microbiota composition, particularly




Fig. 4. The bar plots verifying relative abundance differences of fecal microbiota of weaned pigs at the genus level among dietary treatments. *Prevotella* (A), *Eubacterium* (B), and *Lactobacillus* (C) at day 42. CON, basal weaner diet based on corn and soybean meal; HZO, CON + 2,500 ppm ZnO; CZO, CON + 200 mg/kg dietary coated ZnO. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$.

during the first and second critical weeks of post-weaning [32]. However, inorganic zinc has relatively low bioavailability compared with its organic forms, with most of it being discharged from the feces which could be the main source of soil pollution [33,34]. Therefore, we hypothesized that low doses of coated ZnO could enhance the zinc absorption rate in the gastrointestinal (GI) tract while maintaining an efficacy similar to that of high levels of ZnO. In the present study, the dietary CZO group showed improved ADG and feed efficiency during the first two weeks post-weaning compared with the CON group, which is consistent with previous studies [16,18,35]. These results suggest that low doses of coated ZnO promote growth performance through the slow release of Zn by pancreatic lipase and more effective delivery to the small intestine, achieving growth promoting effects similar to those of high levels of conventional ZnO forms. In swine production, ZnO is often added to a weaner diet to alleviate PWD caused by nutritional and physiological stress and *E. coli* infection [36]. In the present study, the CZO group tended to have a lower frequency of diarrhea compared with the CON group during the first two weeks of our study. Additionally, HCT levels in hematological parameters were lower in pigs fed CZO than in those fed CON, which is an indicator of dehydration status typically accompanied by an increase of frequency of diarrhea. Therefore, our results suggest the supplementation with coated ZnO enhanced the intestinal health of pigs, which in turn positively affects their growth performance during the critical period when piglets are most vulnerable to digestive disturbances.

Higher ATTD values of DM, CP, and energy were observed in previous studies with the administration of chelated or coated ZnO in the weaner diet [12,16] which can be attributed to the enhanced activations of digestive enzymes [37] and improved morphology of the small intestine, [38] which positively affects nutrient absorption. However, in contrast to previous reports, our results showed no differences in the ATTD of DM, CP, and energy among the dietary treatments. Biswas et al. [39] reported that the ATTD of DM, nitrogen, and energy were not affected by amino acid-

chelated zinc. This finding aligns with that of our study, suggesting that the effects of zinc may vary depending on the animal breed, zinc source, and its concentration.

Zinc is an essential trace mineral playing a vital role in regulating the immune systems, including innate and adaptive immunity, and zinc deficiency can negatively affect immune functions, development, and overall health [40]. Disruption of the intestinal barrier and penetration of pathogenic bacteria, because of weaning stress, increase the permeability of intestinal tissues [41,42]. This process activates mucosal immune cells resulting in the upregulation including pro-inflammatory cytokines such as TNF- α , which is a potential marker of inflammatory reactions [43]. In the present study, the serum concentration of TNF- α was decreased in the dietary CZO supplementation compared with the CON diet, but did not differ from the HZO group. This finding suggests that the supplementation of zinc in weaner diets modulated systemic immunity by inhibiting the pathway of nuclear factor- κ B, which is a key transcription factor for the expression of TNF- α [44]. Additionally, a tendency of cortisol concentration was observed on day 7 in the HZO group compared to the CON group. Cortisol, a biomarker of stress, reflects activation of the hypothalamic–pituitary–adrenal (HPA) axis, which is triggered during early weaning stress and leads to the overproduction of cortisol and pro-inflammatory cytokines in response to inflammation [45]. In addition, zinc supplementation in weaned pigs not only regulating immune functions through cell-mediated immunity but also enhancing humoral immunity. IgA, IgM, and IgG are the main antibodies activated by B lymphocytes and are essential for humoral immunity [46]. Our current study indicated that supplemented with CZO group tended to have increased serum IgM levels on day 14. A previous study revealed that ZnO supplementation enhances IgM levels [47], which is consistent with our results. IgM is the first antibody produced in response to antigens and has major functions in the host defense system against infections. These results suggest that CZO supplementation is positively affected during early weaning, which is the most critical period for infections. Our findings indicate that supplementation with pharmacological and low doses of coated ZnO alleviates the stress response and modulates immune responses, supporting the potential effect of enhancing the systemic immunity of weaned pigs.

The gut ecosystem is affected by the interactions between the immune systems and intestinal microorganisms, leading to the utilization of nutrients and contributing to the maintenance of host homeostasis [48]. The intestinal microbiota showed increased diversity and richness after weaning which depends on the species, age, and diet of the animals [49]. In the current study, we observed that dietary HZO reduced the species richness indices (Observed OTUs and Chao1) on day 42 compared with dietary CZO. Previous studies have reported lower richness values in groups treated with high levels of ZnO [15,50], which is in agreements with the results observed in our study. Regarding beta diversity, communities of fecal microbiota were distinctly identified among the dietary treatments on day 42. Based on the results of microbial diversities, the dietary CZO group potentially contributed to the improvement in microbial richness and stable microbial communities among the dietary treatments which may enhance intestinal health [51]. In addition to diversity analysis, the relative taxonomic abundance of fecal microbial communities was investigated to further evaluate the intestinal health of the pigs. Firmicutes and Bacteroidetes phyla were more abundant in the HZO and CZO groups than in the CON group at the phylum level on day 42. These are the most common phyla in the GI tract of humans and pigs and can maintain the balance of energy and promote the production of short chain fatty acids (SCFAs) [52,53]. In the present study, the CZO group showed higher abundance of the genera *Prevotella* and *Eubacterium* compared with the CON group on day 42. *Prevotella* is known as one of the most dominant genera in the GI tract during the weaning period and plays a pivotal role in producing SCFAs that serve as an energy source and help protect against gut inflammation [54,55]. Furthermore, the *Prevotella*-

driven enterotype was positively correlated with weight gain and feed intake [56], and the higher abundance of *Prevotella* observed in the present study indicates the beneficial effects of growth performance on weaned pigs. The genus *Eubacterium* has been reported to be a butyrate-producing bacterium as well as a core component of the gut microbiome [57]. Previous studies have shown that *Eubacterium* spp. contributes to maintaining gut integrity by modulating the gut microbiota and are considered to have similar effects to probiotics, such as *Lactobacillus* and *Bacillus* strain in promoting intestinal health [58]. In addition, the higher *Lactobacillus* abundance was observed in the HZO group than in the CZO and CON groups on day 42. Our current results are consistent with those of previous studies reported that only pigs fed pharmacological levels of ZnO showed a higher abundance of *Lactobacillus* than those fed low doses of modified ZnO [12]. This result may vary depending on the form of ZnO and it may be considered that the proportion of *Lactobacillus* is relatively reduced as the proportion of other beneficial bacteria increased in the CZO group.

CONCLUSION

In summary, our study showed that supplementation with dietary coated ZnO improved growth performance and alleviated the frequency of diarrhea in weaned pigs. Additionally, dietary coated ZnO at low dosages modulated systemic immunity and enhanced the host gut health by altering the microbial communities and shifting the relative microbial compositions of the fecal microbiota. As systemic immune modulation and improved gut health are positively correlated with growth performance, these compounds could potentially replace high levels of ZnO, which mainly contributes to environmental pollution. Further studies should be considered to analyze the metagenomics indicators of zinc to elucidate the potential effects of coated ZnO.

REFERENCES

1. Kinara E, Moturi J, Hosseindoust A, Mun JY, Tajudeen H, Ha SH, et al. Dietary supplementation of *Lactobacillus salivarius* in suckling and weanling piglets modulates intestinal microbiota, morphology and improves growth performance. *J Anim Sci Technol.* 2025;67:826-38. <https://doi.org/10.5187/jast.2024.e58>
2. Song D, Lee J, Yoo Y, Oh H, Chang S, An J, et al. Effects of probiotics on growth performance, intestinal morphology, intestinal microbiota weaning pig challenged with *Escherichia coli* and *Salmonella enterica*. *J Anim Sci Technol.* 2025;67:106-36. <https://doi.org/10.5187/jast.2023.e119>
3. Kyoung H, Kim S, Cho JH, Choe J, Ahn J, Nam J, et al. Dietary flavor supplementation enhanced growth performance and alleviated diarrhea of weaned pigs by modulating gut microbiota and systemic immunity. *J Anim Sci Technol.* 2026;68:168-80. <https://doi.org/10.5187/jast.2025.e3>
4. Oh HJ, Kim MH, Lee JH, Kim YJ, An JW, Chang SY, et al. Effects of different inorganic: organic zinc ratios or combination of low crude protein diet and mixed feed additive in weaned piglet diets. *J Anim Sci Technol.* 2022;64:23-37. <https://doi.org/10.5187/jast.2021.e120>
5. Nam J, Cho JH, Kim HB, Ahn J, Kang Y, Kim S, et al. Dietary coated copper and zinc improved growth performance by modulating immune responses and fecal microbiota of weaned pigs. *J Anim Sci Technol.* 2025;67:1050-66. <https://doi.org/10.5187/jast.2024.e70>
6. Hansen SV, Nørgaard JV, Woyengo T, Nielsen TS. The relationship between zinc intake, dietary content, and fecal excretion in pigs. *Livest Sci.* 2023;271:105228. <https://doi.org/10.1016/j.livsci.2023.105228>
7. Tang X, Xiong K, Zeng Y, Fang R. The mechanism of zinc oxide in alleviating diarrhea in

- piglets after weaning: a review from the perspective of intestinal barrier function. *Int J Mol Sci.* 2024;25:10040. <https://doi.org/10.3390/ijms251810040>
8. Bonetti A, Tugnoli B, Piva A, Grilli E. Towards zero zinc oxide: feeding strategies to manage post-weaning diarrhea in piglets. *Animals.* 2021;11:642. <https://doi.org/10.3390/ani11030642>
 9. Satessa GD, Kjeldsen NJ, Mansouryar M, Hansen HH, Bache JK, Nielsen MO. Effects of alternative feed additives to medicinal zinc oxide on productivity, diarrhoea incidence and gut development in weaned piglets. *Animal.* 2020;14:1638-46. <https://doi.org/10.1017/S1751731120000154>
 10. Burrough ER, De Mille C, Gabler NK. Zinc overload in weaned pigs: tissue accumulation, pathology, and growth impacts. *J Vet Diagn Invest.* 2019;31:537-45. <https://doi.org/10.1177/1040638719852144>
 11. Wei X, Tsai T, Howe S, Zhao J. Weaning induced gut dysfunction and nutritional interventions in nursery pigs: a partial review. *Animals.* 2021;11:1279. <https://doi.org/10.3390/ani11051279>
 12. Oh HJ, Park YJ, Cho JH, Song MH, Gu BH, Yun W, et al. Changes in diarrhea score, nutrient digestibility, zinc utilization, intestinal immune profiles, and fecal microbiome in weaned piglets by different forms of zinc. *Animals.* 2021;11:1356. <https://doi.org/10.3390/ani11051356>
 13. Sun Y, Ma N, Qi Z, Han M, Ma X. Coated zinc oxide improves growth performance of weaned piglets via gut microbiota. *Front Nutr.* 2022;9:819722. <https://doi.org/10.3389/fnut.2022.819722>
 14. Peng P, Deng D, Chen S, Li C, Luo J, Romeo A, et al. The effects of dietary porous zinc oxide supplementation on growth performance, inflammatory cytokines and tight junction's gene expression in early-weaned piglets. *J Nutr Sci Vitaminol.* 2020;66:311-8. <https://doi.org/10.3177/jnsv.66.311>
 15. Shen J, Chen Y, Wang Z, Zhou A, He M, Mao L, et al. Coated zinc oxide improves intestinal immunity function and regulates microbiota composition in weaned piglets. *Br J Nutr.* 2014;111:2123-34. <https://doi.org/10.1017/S0007114514000300>
 16. Lei XJ, Kim IH. Low dose of coated zinc oxide is as effective as pharmacological zinc oxide in promoting growth performance, reducing fecal scores, and improving nutrient digestibility and intestinal morphology in weaned pigs. *Anim Feed Sci Technol.* 2018;245:117-25. <https://doi.org/10.1016/j.anifeedsci.2018.06.011>
 17. Zhang L, Guo Q, Duan Y, Lin X, Ni H, Zhou C, et al. Comparison of the effects of inorganic or amino acid-chelated zinc on mouse myoblast growth in vitro and growth performance and carcass traits in growing-finishing pigs. *Front Nutr.* 2022;9:857393. <https://doi.org/10.3389/fnut.2022.857393>
 18. Kwon CH, Lee CY, Han SJ, Kim SJ, Park BC, Jang I, et al. Effects of dietary supplementation of lipid-encapsulated zinc oxide on colibacillosis, growth and intestinal morphology in weaned piglets challenged with enterotoxigenic *Escherichia coli*. *Anim Sci J.* 2014;85:805-13. <https://doi.org/10.1111/asj.12215>
 19. Dong X, Xu Q, Wang C, Zou X, Lu J. Supplemental-coated zinc oxide relieves diarrhoea by decreasing intestinal permeability in weanling pigs. *J Appl Anim Res.* 2019;47:362-8. <https://doi.org/10.1080/09712119.2019.1645673>
 20. NRC (National Research Council). Nutrient requirements of swine. 11th rev. ed. The National Academies Press; 2012.
 21. Park S, Choe J, Cho JH, Jang KB, Kyoung H, Park KI, et al. Determination of optimal energy system and level for growing pigs. *J Anim Sci Technol.* 2024;66:514-22. <https://doi.org/10.5187/jast.2023.e63>
 22. AOAC (Association of Official Analytical Chemists) International. Official Methods of Analysis. 18th ed. Method 962.09. AOAC International; 2010.

23. Stein HH, Sève B, Fuller MF, Moughan PJ, de Lange CFM. Invited review: amino acid bioavailability and digestibility in pig feed ingredients: terminology and application. *J Anim Sci.* 2007;85:172-80. <https://doi.org/10.2527/jas.2005-742>
24. Kyoung H, Park KI, Shin S, Ahn J, Nam J, Kang Y, et al. Probiotic *Lactiplantibacillus plantarum* improved growth performance of weaned pigs by enhancing intestinal health and modulating immune responses. *J Anim Sci Technol.* 2026;68:541-61. <https://doi.org/10.5187/jast.2025.e22>
25. Lee JJ, Kyoung H, Cho JH, Park KI, Kim Y, Ahn J, et al. Change in the gut microbiota of lactating sows and their piglets by inclusion of dietary spray-dried plasma in sow diets. *J Microbiol Biotechnol.* 2024;34:516-24. <https://doi.org/10.4014/jmb.2311.11001>
26. Parks DH, Tyson GW, Hugenholtz P, Beiko RG. STAMP: statistical analysis of taxonomic and functional profiles. *Bioinformatics.* 2014;30:3123-4. <https://doi.org/10.1093/bioinformatics/btu494>
27. Pluske JR, Hampson DJ, Williams IH. Factors influencing the structure and function of the small intestine in the weaned pig: a review. *Livest Prod Sci.* 1997;51:215-36. [https://doi.org/10.1016/S0301-6226\(97\)00057-2](https://doi.org/10.1016/S0301-6226(97)00057-2)
28. Campbell JM, Crenshaw JD, Polo J. The biological stress of early weaned piglets. *J Anim Sci Biotechnol.* 2013;4:19. <https://doi.org/10.1186/2049-1891-4-19>
29. Højberg O, Canibe N, Poulsen HD, Hedemann MS, Jensen BB. Influence of dietary zinc oxide and copper sulfate on the gastrointestinal ecosystem in newly weaned piglets. *Appl Environ Microbiol.* 2005;71:2267-77. <https://doi.org/10.1128/AEM.71.5.2267-2277.2005>
30. Zhu C, Lv H, Chen Z, Wang L, Wu X, Chen Z, et al. Dietary zinc oxide modulates antioxidant capacity, small intestine development, and jejunal gene expression in weaned piglets. *Biol Trace Elem Res.* 2017;175:331-8. <https://doi.org/10.1007/s12011-016-0767-3>
31. Dahmer PL, Jones CK, Ferreyra FM. Evaluation of a microencapsulated form of zinc oxide on weanling pig growth performance, fecal zinc excretion, and small intestinal morphology. *Transl Anim Sci.* 2024;8:txad146. <https://doi.org/10.1093/tas/txad146>
32. Ouyang Z, Ren P, Zheng D, Huang L, Wei T, Yang C, et al. Hydrothermal synthesis of a new porous zinc oxide and its antimicrobial evaluation in weanling piglets. *Livest Sci.* 2021;248:104499. <https://doi.org/10.1016/j.livsci.2021.104499>
33. Liu Y, Ma YL, Zhao JM, Vazquez-Añón M, Stein HH. Digestibility and retention of zinc, copper, manganese, iron, calcium, and phosphorus in pigs fed diets containing inorganic or organic minerals. *J Anim Sci.* 2014;92:3407-15. <https://doi.org/10.2527/jas.2013-7080>
34. Xiong Y, Cui B, He Z, Liu S, Wu Q, Yi H, et al. Dietary replacement of inorganic trace minerals with lower levels of organic trace minerals leads to enhanced antioxidant capacity, nutrient digestibility, and reduced fecal mineral excretion in growing-finishing pigs. *Front Vet Sci.* 2023;10:1142054. <https://doi.org/10.3389/fvets.2023.1142054>
35. Xiao X, Guo K, Liu J, Liu Y, Yang C, Xu Y, et al. The effect of sodium alginate-coated nano-zinc oxide on the growth performance, serum indexes and fecal microbial structure of weaned piglets. *Animals.* 2024;14:146. <https://doi.org/10.3390/ani14010146>
36. Johanns VC, Ghazisaeedi F, Epping L, Semmler T, Lübke-Becker A, Pfeifer Y, et al. Effects of a four-week high-dosage zinc oxide supplemented diet on commensal *Escherichia coli* of weaned pigs. *Front Microbiol.* 2019;10:2734. <https://doi.org/10.3389/fmicb.2019.02734>
37. Hu C, You ZT, Zhu K, Luan ZS. Effects of nano zinc oxide on growth performance and intestinal mucosal barrier in weaner piglets. *Chin J Anim Nutr.* 2012;24:285-90.
38. Furbeyre H, van Milgen J, Mener T, Gloaguen M, Labussière E. Effects of dietary supplementation with freshwater microalgae on growth performance, nutrient digestibility and gut health in weaned

- piglets. *Animal*. 2017;11:183-92. <https://doi.org/10.1017/S1751731116001543>
39. Biswas S, Dang DX, Kim IH. Comparison of the effects of zinc oxide and zinc aspartic acid chelate on the performance of weaning pigs. *J Anim Sci Technol*. 2024;66:125-34. <https://doi.org/10.5187/jast.2023.e39>
 40. Haase H, Rink L. The immune system and the impact of zinc during aging. *Immun Ageing*. 2009;6:9. <https://doi.org/10.1186/1742-4933-6-9>
 41. Moeser AJ, Pohl CS, Rajput M. Weaning stress and gastrointestinal barrier development: implications for lifelong gut health in pigs. *Anim Nutr*. 2017;3: 313-21. <https://doi.org/10.1016/j.aninu.2017.06.003>
 42. Goto Y, Kiyono H. Epithelial barrier: an interface for the cross-communication between gut flora and immune system. *Immunol Rev*. 2011;245:147-63. <https://doi.org/10.1111/j.1600-065X.2011.01078.x>
 43. Zhang L, Xu YQ, Liu HY, Lai T, Ma JL, Wang JF, et al. Evaluation of *Lactobacillus rhamnosus* GG using an *Escherichia coli* K88 model of piglet diarrhoea: effects on diarrhoea incidence, faecal microflora and immune responses. *Vet Microbiol*. 2010;141:142-8. <https://doi.org/10.1016/j.vetmic.2009.09.003>
 44. Xia T, Lai W, Han M, Han M, Ma X, Zhang L. Dietary ZnO nanoparticles alters intestinal microbiota and inflammation response in weaned piglets. *Oncotarget*. 2017;8:64878-91. <https://doi.org/10.18632/oncotarget.17612>
 45. Sánchez J, Matas M, Ibáñez-López FJ, Hernández I, Sotillo J, Gutiérrez AM. The connection between stress and immune status in pigs: a first salivary analytical panel for disease differentiation. *Front Vet Sci*. 2022;9:881435. <https://doi.org/10.3389/fvets.2022.881435>
 46. Zhang T, Zhang N, Peng S, Zhang Y, Wang H, Huang S, et al. Effects of dietary valine chelated zinc supplementation on growth performance, antioxidant capacity, immunity, and intestine health in weaned piglets. *Biol Trace Elem Res*. 2024;202:2577-87. <https://doi.org/10.1007/s12011-023-03870-2>
 47. Wang Q, Ying J, Zou P, Zhou Y, Wang B, Yu D, et al. Effects of dietary supplementation of humic acid sodium and zinc oxide on growth performance, immune status and antioxidant capacity of weaned piglets. *Animals*. 2020;10:2104. <https://doi.org/10.3390/ani10112104>
 48. Rist VTS, Weiss E, Eklund M, Mosenthin R. Impact of dietary protein on microbiota composition and activity in the gastrointestinal tract of piglets in relation to gut health: a review. *Animal*. 2013;7:1067-78. <https://doi.org/10.1017/S1751731113000062>
 49. Frese SA, Parker K, Calvert CC, Mills DA. Diet shapes the gut microbiome of pigs during nursing and weaning. *Microbiome*. 2015;3:28. <https://doi.org/10.1186/s40168-015-0091-8>
 50. Ortiz Sanjuán JM, Manzanilla EG, Cabrera-Rubio R, Crispie F, Cotter PD, Garrido JJ, et al. Using shotgun sequencing to describe the changes induced by in-feed zinc oxide and apramycin in the microbiomes of pigs one week postweaning. *Microbiol Spectr*. 2022;10:e01597-22. <https://doi.org/10.1128/spectrum.01597-22>
 51. Schokker D, Kar SK, Willems E, Bossers A, Dekker RA, Jansman AJM. Dietary supplementation of zinc oxide modulates intestinal functionality during the post-weaning period in clinically healthy piglets. *J Anim Sci Biotechnol*. 2023;14:122. <https://doi.org/10.1186/s40104-023-00925-1>
 52. Zhang W, Ma C, Xie P, Zhu Q, Wang X, Yin Y, et al. Gut microbiota of newborn piglets with intrauterine growth restriction have lower diversity and different taxonomic abundances. *J Appl Microbiol*. 2019;127:354-69. <https://doi.org/10.1111/jam.14304>
 53. Yang Q, Huang X, Zhao S, Sun W, Yan Z, Wang P, et al. Structure and function of the fecal microbiota in diarrheic neonatal piglets. *Front Microbiol*. 2017;8:502. <https://doi.org/10.3389/>

fmicb.2017.00502

54. Umu ÖCO, Frank JA, Fangel JU, Oostindjer M, da Silva CS, Bolhuis EJ, et al. Resistant starch diet induces change in the swine microbiome and a predominance of beneficial bacterial populations. *Microbiome*. 2015;3:16. <https://doi.org/10.1186/s40168-015-0078-5>
55. Guevarra RB, Hong SH, Cho JH, Kim BR, Shin J, Lee JH, et al. The dynamics of the piglet gut microbiome during the weaning transition in association with health and nutrition. *J Anim Sci Biotechnol*. 2018;9:54. <https://doi.org/10.1186/s40104-018-0269-6>
56. Amat S, Lantz H, Munyaka PM, Willing BP. Prevotella in pigs: the positive and negative associations with production and health. *Microorganisms*. 2020;8:1584. <https://doi.org/10.3390/microorganisms8101584>
57. Mukherjee A, Lordan C, Ross RP, Cotter PD. Gut microbes from the phylogenetically diverse genus Eubacterium and their various contributions to gut health. *Gut Microbes*. 2020;12:1802866. <https://doi.org/10.1080/19490976.2020.1802866>
58. Gibson GR, Hutkins R, Sanders ME, Prescott SL, Reimer RA, Salminen SJ, et al. Expert consensus document: The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nat Rev Gastroenterol Hepatol*. 2017;14:491-502. <https://doi.org/10.1038/nrgastro.2017.75>