

Effects of *Bacillus licheniformis* and *Bacillus subtilis* on growth performance, gut health, and immunity in pigs


Min-Gi Han^{1#}, Ran Lee^{2#†}, Hyeon-Woo Sim², Won-Young Lee³,
Young-Bin Park⁴, Sung-Ho Lee^{4*}, Hyun-Jung Park^{2†*}

¹Department of Stem Cell and Regenerative Biotechnology, KU Institute of Technology (KIT), Konkuk University, Seoul, Korea

²Department of Animal Biotechnology, College of Life Science, Sangji University, Wonju, Korea

³Department of Livestock, Korea National University of Agriculture and Fisheries, Jeonju, Korea

⁴Research center, Woogene B&G Co., LTD, Hwaseong, Korea



Received: Jan 7, 2025

Revised: Jun 30, 2025

Accepted: Jul 11, 2025

#These authors contributed equally to this work.

†Current affiliation: Department of Biotechnology, College of Biomedical and Health Science, Konkuk University, Chungju, Korea

*Corresponding author

Sung-Ho Lee

E-mail: dasal235@gmail.com

Hyun-Jung Park

E-mail: sunrisehj02@kku.ac.kr

Copyright © 2026 Korean Society of Animal Science and Technology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Min-Gi Han

<https://orcid.org/0000-0002-4567-6632>

Ran Lee

<https://orcid.org/0000-0002-1988-5481>

Hyeon-Woo Sim

<https://orcid.org/0000-0002-4288-7450>

Won-Young Lee

<https://orcid.org/0000-0003-0291-5581>

Abstract

This study evaluated the effects of dietary supplementation with *Bacillus licheniformis* and *Bacillus subtilis* on the gut health and growth performance of weaning pigs. A total of 28-day-old piglets (n = 50 per group) were randomly assigned to four groups: CTRL (basal diet), BS (basal diet + *Bacillus subtilis*), BL (basal diet + *Bacillus licheniformis*), or BSL (basal diet + both *Bacillus* strains). After 28 days, pigs fed BSL had the highest body weight, although the difference was not statistically significant. *Bacillus*-treated pigs showed no differences in inflammatory cytokines or hematological characteristics, except for a reduction in cortisol levels in the supplemented groups. The genes encoding inflammatory cytokines were not significantly expressed among the groups. Intestinal morphology was significantly improved with villus height, villus/crypt ratio in BL ($p < 0.05$) and BSL ($p < 0.01$), and smooth circular muscle thickness showing notable increases in BS ($p < 0.05$), BL ($p < 0.001$) and BSL ($p < 0.001$). The tight junction proteins were significantly expressed higher in all *Bacillus*-treated groups, with BSL ($p < 0.001$) exhibiting the greatest improvement. Gene expression levels of *ZO-1* and *occludin* were also significantly elevated in the ileum, with the most pronounced increases observed in BSL ($p < 0.01$). These results suggest that combining *Bacillus subtilis* and *Bacillus licheniformis* synergistically enhances intestinal morphology and barrier function, highlighting their potential as functional feed additives.

Keywords: *Bacillus licheniformis* and *Bacillus subtilis*, Weaning pig, Growth performance, Gut health, Intestinal barrier

INTRODUCTION

Increasing livestock productivity significantly affects the nutrition and environment of animals. Dietary probiotics are widely used as alternatives to antibiotics to promote pig health and productivity [1].

Young-Bin Park
<https://orcid.org/0000-0002-2342-7475>
 Sung-Ho Lee
<https://orcid.org/0009-0001-5753-3516>
 Hyun-Jung Park
<https://orcid.org/0000-0002-4589-2052>

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This work was supported by Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry (IPET) through Agri-Food Export Enhancement Technology Development Program, funded by Ministry of Agriculture, Food and Rural Affairs (MAFRA) (No. RS-2023-00234143). This study was supported by the Korea Institute of Planning and Evaluation of Technology in Food, Agriculture and Forestry (IPET) through the Agriculture and Food Convergence Technologies Program for Research Manpower Development Program funded by the Ministry of Agriculture, Food, and Rural Affairs (MAFRA). Grant number: RS-2024-00400922.

Acknowledgements

All pig feeds and additives used in this study were provided by Woogene B&G Co., LTD.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Lee SH, Park HJ.
 Data curation: Lee WY, Park HJ.
 Formal analysis: Lee WY, Park HJ.
 Methodology: Park YB.
 Software: Lee R, Sim HW.
 Validation: Sim HW, Lee SH.
 Investigation: Han MG, Lee R, Sim HW, Park YB.
 Writing - original draft: Han MG, Lee R.
 Writing - review & editing: Han MG, Lee R, Sim HW, Lee WY, Park YB, Lee SH, Park HJ.

Ethics approval and consent to participate

The animal study protocol was approved by the Institutional Animal Care and Use Committees (IACUC, WG-IACUC-2024-003). The procedure was performed by veterinary professionals according to the clinical standard.

Declaration of generative AI

No AI tools were used in this article.

The gut flora of pigs profoundly affects host health [2]. Supplementing pig feed with probiotics improves growth performance, gut health, and immunity [3]. Intestinal microorganisms influence numerous immune tissues and cells in the gut, which in turn affect systemic host homeostasis [4,5]. Specifically, expression of immune-related genes in the gut and associated lymphoid tissues has been studied, highlighting the relations between the gut microbiota and immune function [6]. Additionally, pro-biotic administration alleviates colitis and increases mucosal expression through the action of tight junction proteins, such as Zonula occludens-1 (ZO-1) and Occludin [7].

Among probiotics, *Bacillus* strains are known to improve the intestinal health of various hosts, including pigs. These strains exhibit antibacterial and antioxidant properties and positively influence immune function [8,9]. They are also effective in preventing and alleviating intestinal inflammation [10,11]. Originating in the soil, *Bacillus* strains have been found in the intestines of humans and pigs, where they act as probiotics [12–14]. Specifically, *Bacillus subtilis* and *Bacillus licheniformis* are noted for their ability to form biofilms, enabling them to persist in the intestine by producing anaerobic spores that can withstand gastrointestinal stress [15]. These unique characteristics indicate that both species are valuable feed additives.

Feeding *Bacillus subtilis* to pigs increases average daily gain and feed efficiency while reducing the incidence of diarrhea [16]. Moreover, *Bacillus subtilis* enhances pig growth and disease resistance by promoting digestive enzyme activity and increasing villus height in the ileum [17]. *Bacillus subtilis* also suppresses harmful bacteria in the large intestine, strengthens the probiotic microbial community, and modulates metabolic activity, collectively improving growth efficiency in pigs [18]. *Bacillus subtilis* and *Lactobacillus* can alleviate colitis and enhance mucosal expression by regulating the expression of tight junction proteins, including ZO-1 and Occludin [19]. Fermented *Bacillus subtilis* products have similarly been shown to boost immunity in the small intestine, contributing to growth efficiency [20].

Bacillus licheniformis also supports digestion by producing proteases and amylases [21,22]. It detoxifies zearalenone, a mycotoxin found in feed [23], and produces surfactin, which exhibits potent antibacterial activity against *Clostridium perfringens*, a common cause of diarrhea in pigs [24]. Fermented products containing *Bacillus licheniformis* positively affect pig gut microbiota and reduce the frequency of diarrhea [25]. Additionally, the combined administration of *Bacillus licheniformis* and other beneficial probiotics has been associated with improved immunity and reduced diarrheal symptoms [26].

Recent studies have confirmed that feeding both *Bacillus licheniformis* and *Bacillus subtilis* directly to pigs increases digestibility, enhances *Lactobacillus* populations in feces, and supports growth performance [27,28]. Despite these reports, the effects of *Bacillus licheniformis* and *Bacillus subtilis* on intestinal tissue and villi in relation to intestinal immunity and the function of weaning pigs have not been clearly identified. Therefore, this study aimed to determine the contribution of dietary supplementation with these two bacterial species to villus structure, particularly with respect to tight junctions in the intestines of weaning pigs, in the context of serological immune indicators.

MATERIALS AND METHODS

Experimental design

In this experiment, 28-day-old piglets (n = 50 per group; total = 200) were randomly selected, excluding underweight individuals and assigned to four groups: CTRL (basal diet), BS (basal diet + *Bacillus subtilis*), BL (basal diet + *Bacillus licheniformis*), or BSL (basal diet + both *Bacillus* strains). All piglets were provided with the same starter diet, which did not contain antibiotics, prebiotics, or probiotics. The handling procedures followed the ethical guidelines established by the Institutional Animal Care and

Use Committee, and the study was conducted with approval under protocol WG-IACUC-2024-003. The experimental feeding regimen consisted of two phases: Phase 1 and Phase 2 diets were supplied as the basal diet for 14 days each, while feed additives were continuously administered for 28 days. Resistant maltodextrin (2%, Daesang, Korea) was included in BS, BL and BSL groups as a prebiotic to function synergistically with probiotics, thus constituting a synbiotic formulation. Throughout the experimental period, piglets had unrestricted access to both feed and water. The detailed composition of the basal diet and the additional feed additives are presented in Tables 1 and 2, respectively.

Material collection and preparation

At both the beginning and the end of the experiment, body weight changes were measured in 40 piglets, with 10 animals assigned to the control (CTRL) group and 10 to each treatment group

Table 1. Composition and nutrient levels of the basal diet (air-dry basis)

Item	Phase 1	Phase 2
Ingredients (%)		
Corn	33.39	48.23
Soybean meal	15.20	18.10
Whey	15.00	10.00
Lactose	12.00	6.00
Fishmeal	5.00	3.00
Spray dry plasma protein	5.00	2.00
Animal fat	4.59	3.99
Sugar	6.00	4.00
L-Lysine (78%)	0.35	0.49
DL-Methionine (99%)	0.13	0.18
L-Threonine (99%)	0.11	0.17
L-Tryptophan (100%)	0.12	0.31
Limestone	1.10	1.10
Monocalcium phosphate	0.88	1.31
Salt	0.50	0.50
ZnO	0.30	0.30
Vitamin premix ¹⁾	0.11	0.11
Mineral premix ²⁾	0.22	0.22
Total	100.00	100.00
Chemical composition (%)		
ME	3,400	3,350
Crude protein	20.00	18.00
Crude fat	6.38	5.94
Ash	5.76	5.59
Ca	0.85	0.85
P	0.62	0.69
Lysine	1.53	1.40
Met + Cys	0.87	0.79
Phenylalanine	0.90	0.82

¹⁾Supplied per kg of diet: 16,000 IU vitamin A (palmitate), 2.00 mg vitamin B₁ (thiamin), 5.00 mg vitamin B₂ (riboflavin), 2.00 mg vitamin B₆ (pyridoxine), 0.03 mg vitamin B₁₂ (cyanocobalamin), 25.00 mg niacin, 0.40 mg folic acid, 0.05 mg biotin, 5.00 mg ethoxyquin, 2,000 IU vitamin D₃ (cholecalciferol), 75.00 mg vitamin E (dl- α -tocopheryl acetate), and 2.00 mg vitamin K₃ (menadione).

²⁾Supplied per kg of diet: 100 mg Fe, 6 mg Cu, 4 mg Mn, 0.3 mg Se, 0.14 mg I, and 0.25 mg Co.

Table 2. Dietary supplementation of three different treatments based on the basal diet

Item	CTRL	Treat 1 (BS)	Treat 2 (BL)	Treat 3 (BSL)
<i>Bacillus subtilis</i> (CFU/g)	-	1×10^8	-	1×10^8
<i>Bacillus licheniformis</i> (CFU/g)	-	-	1×10^8	1×10^8
Dietary fiber (resistant maltodextrin)	-	2%	2%	2%

CTRL, control group (basal diet); BS, basal diet + *Bacillus subtilis*; BL, basal diet + *Bacillus licheniformis*; BSL, basal diet + *Bacillus subtilis* + *Bacillus licheniformis*.

(BS, BL, and BSL). Additionally, a total of 24 piglets (six per group: CTRL, BS, BL, and BSL) were selected for molecular biological analysis. Blood samples were collected in the morning under fasting conditions. The blood for complete blood count (CBC) analysis was collected from the anterior inferior vena cava, placed in EDTA K3 tubes (AB medical), and stirred at room temperature for 2 h. Blood for serum analysis was centrifuged at $3,000 \times g$ for 10 min at 4°C to separate serum, and stored at 4°C . The collected whole blood (1 mL) and serum (600 μL) were analyzed for alanine aminotransferase (ALT), aspartate aminotransferase (AST), creatinine, cortisol, and CBC at Samkwang Medical Research Institute. All piglets were humanely euthanized using electrical stunning (250 V, 0.5 A) for 5–6 seconds. Ileum tissue samples were immediately excised and sectioned into 2.0-cm-thick slices using phosphate-buffered saline (PBS; Welgene).

Determination of inflammatory cytokines

Serum was analyzed using enzyme-linked immunosorbent assay (ELISA) kits to quantify interferon- γ (IFN- γ , MyBioSource), interleukin 6 (IL-6, MyBioSource), and tumor necrosis factor- α (TNF- α , MyBioSource), following the manufacturer's guidelines. Absorbance readings were taken at 450 nm, with a reference wavelength of 570 nm, using an Epoch spectrophotometer (BioTek). Six serum samples from each group (total: 24 samples) were tested, with measurements conducted at least three times. Until analysis, serum samples were preserved at -80°C .

Intestinal histomorphology

Ileum tissue samples were fixed in 4% paraformaldehyde at 4°C for 48 hours. The samples were then cut into 1.0 cm sections, rinsed with distilled water for 30 minutes, and sequentially treated with ethanol at concentrations of 70%, 80%, 90%, 95%, and 100% (each for 90 minutes) to ensure dehydration while being gently agitated at 24°C . The dehydrated tissues were embedded in paraffin wax for 2 hours and subsequently formed into blocks at room temperature. Using a Leica Biosystems microtome (Nussloch, Germany), the paraffin-embedded ileum tissues were sectioned into 4- μm -thick slices, dried in a slide dryer at 40°C for 12 hours, and mounted onto slides.

For hematoxylin and eosin (H&E) staining and immunohistochemical analysis, the ileum sections were deparaffinized by immersing them in xylene three times for 5 minutes each, followed by rehydration through a graded ethanol series (100%, 90%, 80%, and 70%) and distilled water for 10 minutes at each step. H&E staining was performed at room temperature using hematoxylin for 1 minute and eosin for 30 seconds. After staining, the sections were dehydrated with ethanol (95% and 100%, each for 5 minutes) and then cleared in xylene for 5 minutes. Stained samples were visualized using a microscope (Olympus IX73; Olympus). Five ileum samples from each group (total 20 slides) were selected, and at least three samples were included per replicate. The paraffin blocks were stored at -80°C until further analysis.

Immunostaining

Rehydrated slides underwent heat-induced epitope retrieval by treatment with 10 mM sodium

citrate buffer (pH 6.0) at 100°C for 15 minutes. Primary antibodies (detailed in Table 3) were diluted according to the manufacturer's instructions and incubated with ileum tissue sections 16 hours at 4°C. After washing with PBS, the sections were treated with Alexa Fluor 488- and 568-conjugated secondary fluorescent antibodies for 1 hour at room temperature. Following additional PBS washes, nuclei were counterstained with 4',6-diamidino-2-phenylindole (DAPI; 1 µg/mL). Immunostained ileum tissues were imaged using a fluorescence microscope, and fluorescence intensity was quantified with ImageJ software. Three ileum samples from each group (total 12 slides) were selected, with at least three samples included per replicate.

Quantitative polymerase chain reaction analysis

For qPCR analysis, total RNA was extracted using the AccuPrep® Universal RNA Extraction Kit (Bioneer) with on-column DNase treatment (Qiagen), following the manufacturer's instructions. The extracted RNA was then reverse transcribed into complementary DNA using the RevertAid First Strand cDNA Synthesis Kit (Thermo Fisher Scientific) at a concentration of 1 µg/20 µL.

QPCR was performed using a QuantStudio 1 system (Applied Biosystems) with SYBR Green Master Mix (Thermo Fisher Scientific). The reaction conditions were set as follows: initial denaturation at 94°C for 1 minute, followed by 40 cycles of 94°C for 10 seconds, 57°C for 10 seconds, and 72°C for 20 seconds. Gene expression levels were normalized to the endogenous *β-actin*. Primers were designed using Primer3 software (<http://frodo.wi.mit.edu>) and are detailed in Table 4. Three RNA samples from each group (total 12 samples) were analyzed per group, with each sample tested in triplicate.

Western blotting

Ileal tissue was lysed using radioimmunoprecipitation assay buffer (Thermo Fisher Scientific) supplemented with a protease inhibitor mixture (Roche) to extract proteins. A total of 30 µg of ileal protein was separated on a 4%–20% sodium dodecyl sulfate-polyacrylamide gel (SDS-PAGE; Bio-

Table 3. List of primary antibodies used

Antibody	Manufacturer	Catalog number	Dilution (usage)
β-Actin	Santa Cruz	SC-47778	1:2000 (WB)
Occludin	Invitrogen	71-1500	1:1000 (WB), 1:300 (IHC)
Zonula occludens-1	Invitrogen	40-2200	1:300 (IHC)

WB; western blot; IHC, immunohistochemistry.

Table 4. List of primers for quantitative PCR

Gene	Forward primer	Reverse primer
<i>β-Actin</i>	5'-CTGTCCCTGTACGCCTCTG-3'	5'-GTGGTGGTGAAGCTGTAGCC-3'
<i>IL-1β</i>	5'-GTTCTGCATGAGCTTTGTG-3	5'-TTCTCCATGTCCCTCTTTGG-3
<i>IL-6</i>	5'-TTCAGTCCAGTCGCCTTCT-3'	5'-TGCCAGTACCTCCTTGCTG-3
<i>IL-10</i>	5'-TGCTCTATTGCCTGATCTT-3	5'-CCATCTGGTCCCTTCGTTTG-3
<i>INF-γ</i>	5'-TTCAGCTTTGCGTGACTTT-3'	5'-TGAATGGCCTGGTTATCTTTG-3'
<i>TNF-α</i>	5'-TCCTCACTCACACCATCA-3'	5'-TAGTCGGGCAGGTTGATCTC-3'
<i>Occludin</i>	5'-CAGCAAAGGGATTCTTCC-3'	5'-GCAATGAACACCATCACACC-3'
<i>Zonula occludens-1</i>	5'-GAGTTTGATAGTGGCG-3'	5'-TGGGAGGATGCTGTTG-3'

PCR, polymerase chain reaction; IL-1β, interleukin-1 beta; IL-6, interleukin-6; IL-10, interleukin-10; INF-γ, interferon-gamma; TNF-α, tumor necrosis factor-alpha.

Rad) and subsequently transferred onto a 0.45- μ m polyvinylidene difluoride membrane.

Primary antibodies were diluted in 1% bovine serum albumin buffer containing 0.05% Tween-20, following the manufacturer's instructions, and incubated with the membranes at 4 °C for 16 hours. After washing with TBS-0.05% Tween, the membranes were incubated with HRP-conjugated anti-mouse and anti-rabbit secondary antibodies for 1 hour at 24 °C. Protein detection was performed using Pierce ECL substrate (Thermo Fisher Scientific), and images were captured using the iBright™ Imaging System (Thermo Fisher Scientific). β -Actin was used as quantitative CTRL for protein normalization. The primary antibodies used for western blotting are listed in Table 3. Three protein samples from each group (total 12 samples) were analyzed, with all assays conducted in triplicate.

Statistical analysis

All data are presented as the mean $\pm$ standard error of the mean from independent experiments conducted at least three times in triplicate. Statistical analysis was performed using one-way ANOVA followed by Tukey's post hoc test. All analyses were conducted using SPSS version 29 for Windows (IBM). Statistical significance was defined as follows: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

RESULTS

Physiological effects of *Bacillus* supplementation in weaning pigs

A schematic design of the 28-d supplementation period is presented (Fig. 1A). The body weight of pigs showed no significant difference between the CTRL and test groups (Table 5). To examine the effects of dietary supplementation on organ stress, serum ALT, AST, creatinine, and cortisol levels were evaluated after 28 d (Fig. 1B). ALT and AST levels showed no significant differences. Similarly, creatinine levels were not significantly different across all groups. However, the cortisol levels in BS, BL, and BSL groups ($p < 0.05$) were significantly lower than those in the CTRL group.

Effects of *Bacillus* supplementation on hematological characteristics

To evaluate the effect of *Bacillus* supplementation on hematological characteristics, a complete blood count was conducted for the CTRL, BS, BL, and BSL groups (Fig. 2). There were no significant differences observed in white blood cell (WBC) counts. Similarly, red blood cell (RBC), hematocrit, and hemoglobin levels were consistent across all groups. The platelet counts also showed no notable differences among the groups. Regarding WBC composition, the lymphocyte and monocyte percentage showed no significant differences among all groups. The percentages of neutrophils, basophils and eosinophils were comparable across all groups with no significant. Additionally, RBC quality indicators, including mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), and mean corpuscular hemoglobin concentration (MCHC), showed no significant differences among groups.

Effect of *Bacillus* supplementation on the small intestine of weaning pigs

To examine the effects of *Bacillus* supplementation on the small intestine, the crypt depth, villus height, smooth circular muscle thickness, and villus/crypt ratio were measured using H&E staining (Fig. 3A). H&E staining revealed differences in villus height and smooth muscle thickness between the CTRL and supplemented groups. Crypt depth showed no significant differences among the groups (Fig. 3B). BS did not show significant differences in villus height and villus/crypt ratios compared to CTRL. However, the villus height showed increased in the BL and BSL supplemented groups. Notably, BL ($p < 0.05$) and BSL ($p < 0.01$) groups showed significantly




Fig. 1. Effects of dietary supplementation on liver and kidney function and stress. (A) A schematic descriptive design of the 28-day dietary supplementation with Phase 1 (14 days) and Phase 2 (14 days) of weanling pigs in this study; (B) Evaluation of liver function (AST and ALT), renal function (creatinine), and stress level reduction (cortisol) in the serum of weanling pigs after dietary supplementation. Data represent the mean $\pm$ SE ($n = 6$, $*p < 0.05$ compared to the CTRLs). ALT, alanine transaminase; AST, aspartate transaminase; CTRL, basal diet; BS, basal diet + *Bacillus subtilis*; BL, basal diet + *Bacillus licheniformis*; BSL, basal diet + *Bacillus subtilis* + *Bacillus licheniformis*.

Table 5. Effects of dietary probiotics and symbiotics supplementation on growth performance in weanling pigs

Item	CTRL		BS		BL		BSL	
	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM
Initial BW (kg)	6.36	0.03	6.40	0.04	6.39	0.03	6.37	0.02
Final BW (kg)	15.7	0.26	15.93	0.09	16.15	0.15	16.20	0.17

BW, body weight; SEM, standard error of the mean; CTRL, control group (basal diet); BS, basal diet + *Bacillus subtilis*; BL, basal diet + *Bacillus licheniformis*; BSL, basal diet + *Bacillus subtilis* + *Bacillus licheniformis*.

increased villus height and villus/crypt ratios, with BSL showing the highest values (Fig. 3B). The thickness of the smooth circular muscle was significantly greater in the BS ($p < 0.05$), BL ($p < 0.001$), and BSL ($p < 0.001$) groups compared with that in the CTRL group (Fig. 3B).




Fig. 2. Effects of dietary supplementation on blood characteristics in weaning pigs. The analysis includes the following parameters ($n = 6$): white blood cell (WBC, $10^9/\mu\text{L}$), red blood cell (RBC, $10^9/\mu\text{L}$), hematocrit (Hct, %), hemoglobin (Hb, g/dL), platelet count ($10^9/\mu\text{L}$), lymphocyte (%), monocyte (%), mean corpuscular volume (MCV, fL), mean corpuscular hemoglobin (MCH, pg), mean corpuscular hemoglobin concentration (MCHC, g/dL), neutrophils (%), basophils (%), and eosinophil (%). CTRL, basal diet; BS, basal diet + *Bacillus subtilis*; BL, basal diet + *Bacillus licheniformis*; BSL, basal diet + *Bacillus subtilis* + *Bacillus licheniformis*. Data represent the mean $\pm$ SE.

Effect of *Bacillus* supplementation on tight junctions in the small intestine

To further investigate the improvement of scaffold protein in intestinal epithelial cells to maintain barrier integrity and regulate paracellular permeability, immunostaining for ZO-1 and Occludin was conducted, respectively (Fig. 4). Immunofluorescence images showed that ZO-1 expression was enhanced in all *Bacillus*-supplemented groups. Similarly, Occludin expression increased in the supplemented groups. Furthermore, intensity levels showed that ZO-1 expression was significantly




Fig. 3. Effects of dietary supplementation on ileum characteristics in weaning pigs. (A) Hematoxylin & eosin-stained image of the ileum. Black double-headed arrows indicate crypt depth, villus height, and smooth circular muscle thickness. Scale bar = 500 μm; (B) Graphical representation of villus height, crypt depth, villi/crypt ratio, and smooth circular muscle measurements. Data represent the mean ± SE (n = 3, *p < 0.05, **p < 0.01, ***p < 0.001 compared to the CTRLs). CTRL, basal diet; BS, basal diet + *Bacillus subtilis*; BL, basal diet + *Bacillus licheniformis*; BSL, basal diet + *Bacillus subtilis* + *Bacillus licheniformis*.

increased in the BS ($p < 0.01$), BL ($p < 0.001$), and BSL groups ($p < 0.001$) compared with that in the CTRL. The intensity of Occludin also significantly increased at BS ($p < 0.05$) and was more pronounced at BL ($p < 0.001$) and BSL ($p < 0.001$).

Effect of *Bacillus* supplementation on inflammatory cytokines

Inflammatory cytokine analysis was conducted to evaluate immune responses in CTRL and *Bacillus*-supplemented groups. The concentration of IFN- γ showed no significant difference among the CTRL, BS, BL, and BSL groups. Similarly, no notable changes were observed in TNF- α levels across all groups. IL-6 concentrations were also not significantly different (Fig. 5A).




Fig. 4. Effect of dietary supplementation on tight junctions of ileal villi in weaning pigs. Immunofluorescence staining of ileum tissues. Paraffin-embedded ileum sections were subjected to heat-induced antigen retrieval, followed by incubation with primary antibodies and Alexa Fluor 488/568-conjugated secondary antibodies. Nuclei were counterstained with DAPI, and fluorescence signals were quantified using ImageJ. Zonula occludens-1 (ZO-1, red fluorescence) and Occludin (green fluorescence) protein expression in villi. Scale bar = 50 μm. Graphical representation of quantitative fluorescence expression. Data represent the mean ± SE (n = 3, *p < 0.05, **p < 0.01 and ***p < 0.001 compared to the CTRLs). CTRL, basal diet; BS, basal diet + *Bacillus subtilis*; BL, basal diet + *Bacillus licheniformis*; BSL, basal diet + *Bacillus subtilis* + *Bacillus licheniformis*.




Fig. 5. Effects of dietary supplementation on inflammatory cytokines in weaning pigs. (A) Changes in inflammatory cytokines interferon-gamma (IFN-γ), tumor necrosis factor-alpha (TNF-α), and interleukin-6 (IL-6) in the serum of weaned pigs. (B) mRNA expression levels of inflammatory cytokines, including *IL-1β*, *IL-6*, *IL-10*, *IFN-γ*, and *TNF-α* in the ileum. Data represent the mean ± SE (n=3). CTRL, basal diet; BS, basal diet + *Bacillus subtilis*; BL, basal diet + *Bacillus licheniformis*; BSL, basal diet + *Bacillus subtilis* + *Bacillus licheniformis*.

Subsequently, the relative mRNA expression of inflammatory cytokines was analyzed in the ileum, *interleukin-1 beta* (*IL-1β*), *IL-6*, *interleukin 10* (*IL-10*), *IFN-γ*, and *TNF-α* (Fig. 5B). The proinflammatory cytokines *IL-1β* and *IL-6* showed no significant changes in expression among the CTRL, BS, BL, and BSL groups. The anti-inflammatory cytokine *IL-10* was also consistently expressed, with no notable differences between the groups. Similarly, *IFN-γ* and *TNF-α* did not exhibit significant differences.

Tight junction-related protein and gene expression in the ileum through *Bacillus* supplementation

Tight junction-related protein and gene expression in the ileum were examined to determine the influence of *Bacillus* supplementation. Occludin expression was significantly increased in all *Bacillus*-supplemented groups (Fig. 6A). Notably, the quantified protein data showed a significant increase at BS ($p < 0.05$), with more pronounced increases observed at BL ($p < 0.01$) and BSL ($p < 0.01$). Correspondingly, the mRNA expression of *occludin* was significantly increased in all *Bacillus*-supplemented groups ($p < 0.01$, Fig. 6B). Moreover, the expression of *ZO-1* was also elevated in all *Bacillus*-supplemented groups, particularly in BL ($p < 0.05$) and BSL ($p < 0.05$), which showed significant upregulation (Fig. 6B). Uncropped full western blot images were displayed in Fig. 7.

DISCUSSION

The weaning period is a critical phase in piglet development, significantly influencing growth




Fig. 6. Effect of dietary supplementation on tight junction proteins expressed in the ileum of weaning pigs. (A) Occludin protein expression levels in the ileum of weaning pigs; (B) mRNA expression levels of *occludin* and *zonula occludens-1* (ZO-1) in the ileum of weaning pigs. Data represent the mean $\pm$ SE ($n = 3$, * $p < 0.05$ and ** $p < 0.01$ compared to the CTRLs). CTRL, basal diet; BS, basal diet + *Bacillus subtilis*; BL, basal diet + *Bacillus licheniformis*; BSL, basal diet + *Bacillus subtilis* + *Bacillus licheniformis*.

performance and overall livestock productivity. During this stage, disruptions in gut microbiota balance can lead to intestinal infections, resulting in impaired growth and, in severe cases, increased mortality [3]. Traditionally, large-scale antibiotic administration has been employed to mitigate such infections. However, recent research efforts have focused on replacing antibiotics with probiotics to promote a healthier gut environment [1]. Thus, the application of *Bacillus*-based probiotics to maintain intestinal microbial homeostasis and intestinal development during the weaning period is of considerable interest.

The supplementation of probiotics in weaning piglets has been associated with enhanced disease resistance and improved growth performance [16]. Furthermore, studies have reported increased feed efficiency and a reduction in diarrhea incidence following probiotic administration [29]. Although no significant differences in body weight or feed intake were observed, probiotic-treated piglets exhibited beneficial effects on intestinal immune responses and gut microbiota composition [30]. Although studies on feeding *Bacillus* to pigs have shown inconsistent results regarding growth performance, many have consistently reported that nutrient digestibility is enhanced through improved intestinal microflora and gut health [16,18,29,30]. This suggests that growth performance is influenced by multiple factors and cannot be determined solely by improvements in gut health. In this study, although no statistically significant differences were observed between the CTRL and all the supplemented groups, body weight was the highest in the BSL group compared to the CTRL. The average body weight levels in this study were similar to findings from other studies comparing




Fig. 7. Western full membrane image.

CTRL and probiotic-treated groups with significantly difference [31].

Studies involving long-term probiotic intake have reported decreased ALT, AST, and creatinine levels [32]. However, this study found no signs of toxicity or functional decline in the liver or kidneys of the BS, BL, and BSL groups. This absence of toxicity may be attributed to the focus of this study on the weaning period, which was limited to 28 d. Interestingly, cortisol, a key indicator of stress, was significantly reduced in all *Bacillus*-supplemented groups. *Bacillus*-supplementation during the weaning period may help alleviate stress from both internal and external environmental factors. This is consistent with findings that probiotic administration lowers cortisol levels and reduces mortality in weaned pigs [33]. While no significant differences in WBC composition were observed between the negative CTRL and pigs supplemented with *Bacillus subtilis*, pigs fed *E. coli* exhibited increased monocyte and neutrophil counts [29]. These results suggest that, in the absence of a toxin or pathogen challenge, blood characteristics may not display marked changes. Among the hematological characteristics, no significant differences were observed in blood components. The numerical reduction in BSL monocytes was likely the result of improved immunity owing to the synergistic effects of *Bacillus licheniformis* and *Bacillus subtilis*.

Transcriptome analysis has demonstrated that probiotics regulate cytokine expression [34]. Additionally, a study involving *Bacillus* supplementation in pigs reported an increase in immune indicators because of the antibacterial and antioxidant properties of probiotics [8,9]. However, in a study where probiotics were fed to weaning pigs during the same period, no significant differences in IL-6 and IL-10 levels in the blood were observed [31]. Similarly, our study did not observe significant changes in blood inflammatory cytokine levels, and the expression of both proinflammatory and anti-inflammatory cytokine genes in the ileum showed no differences compared to the CTRL. This minimal immune response across all groups, including the CTRL group, likely reflects a well-managed breeding environment. The reduction of immune responses in weaning pigs is typically more pronounced when probiotics are administered following intestinal diseases caused by enterotoxigenic *E. coli* [35].

Maintaining gastrointestinal health and promoting development during weaning is crucial for pigs [36]. During the neonatal period, gut microbiota influences the stem cell niche of intestinal cells and plays a critical role in villus formation [37]. In the small intestine, villus height and

villus/crypt ratio are key indicators of nutrient absorption capacity [38]. Consistent with studies demonstrating that *Bacillus* supplementation increases villus height in the ileum [17], our results showed increased villus height in all supplemented groups. Notably, BSL exhibited a statistically significant increase, suggesting that the combination of *Bacillus licheniformis* and *Bacillus subtilis* was more effective than individual treatments. Concerning smooth circular muscle thickness, studies have reported that probiotics are associated with intestinal motility [39] and that they relieve intestinal muscle-related diseases, particularly in the large intestine [40]. Similarly, our findings indicate that the thickness of the smooth circular muscle increased in all supplemented groups, with the combined supplementation of *Bacillus licheniformis* and *Bacillus subtilis* showing the greatest effect. These results indicate that the combination of *Bacillus licheniformis* and *Bacillus subtilis* was more effective than individual treatments in improving intestinal morphology. *Bacillus subtilis* plays a crucial role in suppressing pathogenic bacteria in the intestine while promoting the growth of probiotic microbial communities, thereby contributing to a balanced gut environment [18]. Similarly, *Bacillus licheniformis* exhibits detoxification and antibacterial properties that help neutralize harmful compounds and inhibit pathogen growth [23,24].

Due to these characteristics, many studies on *Bacillus* supplementation have consistently demonstrated beneficial changes in gut microbiota composition during the weaning period in pigs [16,29,30]. These findings strongly suggest that in this study, the combined supplementation of *Bacillus licheniformis* and *Bacillus subtilis* contributed to the formation of a beneficial gut microbiota in weaning piglets. Our results indicate that *Bacillus* supplementation promotes intestinal development caused by the increase in beneficial microbiota. The improvements in body weight gain and feeding efficiency appear to be a consequence of this well-developed intestinal environment in the weaning period. This is supported by previous research that intestinal development during the weaning period is influenced by a favorable gut microbiota composition [37]. Additionally, these complementary roles suggest a synergistic interaction between the two strains, leading to an enhanced gut microbiota composition and improved overall intestinal health.

This morphological improvement in the intestine was further confirmed by the increased expression of ZO-1 and Occludin protein. They are essential tight junction proteins in the intestinal epithelium that regulate the proliferation and survival of epithelial cells while supporting nutrient absorption and waste secretion in the intestinal mucosa [41]. Probiotics can improve tight junction proteins, such as ZO-1 and Occludin, thereby supporting a healthy intestinal environment [7]. In line with previous findings of increased tight junctions following *Bacillus subtilis* supplementation [19], we observed enhanced protein expression of ZO-1 and Occludin in BS. Notably, the highest levels of these proteins were observed in the groups supplemented with *Bacillus licheniformis* and *Bacillus subtilis*. These results highlight that *Bacillus* strain supplementation enhanced intestinal health, as evidenced by the increased villus height and villus/crypt ratio in the small intestine during the weaning period. This suggests a synergistic effect between the two strains in enhancing intestinal barrier function. While the mRNA expression of *occludin* and *ZO-1* increased compared with that in the CTRL, the greatest improvement was evident in western blot analyses, which revealed the highest Occludin protein levels in the ileum of pigs supplemented with both *Bacillus licheniformis* and *Bacillus subtilis*. Taken together, *Bacillus* strain supplementation can lower cortisol levels in weaning piglets without inducing significant inflammatory or immune responses. Additionally, by potentially promoting beneficial gut microbiota during the weaning period, that contributes to the development of intestinal structure and the enhancement of intestinal barrier integrity. These results further confirm the synergistic interaction between these two probiotic strains in supporting intestinal health.

CONCLUSIONS

This study demonstrates that the combined supplementation of *Bacillus licheniformis* and *Bacillus subtilis* synergistically enhances intestinal morphology, including villus height, villus/crypt ratio, and smooth circular muscle thickness, while significantly improving tight junction integrity through the increased expression of ZO-1 and Occludin. Although the body weight differences were not statistically significant, the combined supplementation group showed the highest increase, suggesting the potential benefit of combined probiotics. These findings highlight the potential of probiotics as effective feed additives for promoting intestinal health and barrier function in weaning pigs.

REFERENCES

1. Upadhaya SD, Kim IH. Maintenance of gut microbiome stability for optimum intestinal health in pigs – a review. *J Anim Sci Biotechnol*. 2022;13:140. <https://doi.org/10.1186/s40104-022-00790-4>
2. Nowland TL, Plush KJ, Barton M, Kirkwood RN. Development and function of the intestinal microbiome and potential implications for pig production. *Animals*. 2019;9:76. <https://doi.org/10.3390/ani9030076>
3. Gresse R, Chaucheyras-Durand F, Fleury MA, Van de Wiele T, Forano E, Blanquet-Diot S. Gut microbiota dysbiosis in postweaning piglets: understanding the keys to health. *Trends Microbiol*. 2017;25:851-73. <https://doi.org/10.1016/j.tim.2017.05.004>
4. Thaïss CA, Zmora N, Levy M, Elinav E. The microbiome and innate immunity. *Nature*. 2016;535:65-74. <https://doi.org/10.1038/nature18847>
5. Kim YS, Ho SB. Intestinal goblet cells and mucins in health and disease: recent insights and progress. *Curr Gastroenterol Rep*. 2010;12:319-30. <https://doi.org/10.1007/s11894-010-0131-2>
6. Plaza-Diaz J, Gomez-Llorente C, Fontana L, Gil A. Modulation of immunity and inflammatory gene expression in the gut, in inflammatory diseases of the gut and in the liver by probiotics. *World J Gastroenterol*. 2014;20:15632-49. <https://doi.org/10.3748/wjg.v20.i42.15632>
7. Yuan L, Zhu C, Gu F, Zhu M, Yao J, Zhu C, et al. Lactobacillus johnsonii N5 from heat stress-resistant pigs improves gut mucosal immunity and barrier in dextran sodium sulfate-induced colitis. *Anim Nutr*. 2023;15:210-24. <https://doi.org/10.1016/j.aninu.2023.04.012>
8. Ripert G, Racedo SM, Elie AM, Jacquot C, Bressollier P, Urdaci MC. Secreted compounds of the probiotic *Bacillus clausii* strain O/C inhibit the cytotoxic effects induced by *Clostridium difficile* and *Bacillus cereus* toxins. *Antimicrob Agents Chemother*. 2016;60:3445-54. <https://doi.org/10.1128/AAC.02815-15>
9. Khochamit N, Siripornadulsil S, Sukon P, Siripornadulsil W. Antibacterial activity and genotypic-phenotypic characteristics of bacteriocin-producing *Bacillus subtilis* KKU213: potential as a probiotic strain. *Microbiol Res*. 2015;170:36-50. <https://doi.org/10.1016/j.micres.2014.09.004>
10. Zhang HL, Li WS, Xu DN, Zheng WW, Liu Y, Chen J, et al. Mucosa-repairing and microbiota-balancing therapeutic effect of *Bacillus subtilis* alleviates dextrate sulfate sodium-induced ulcerative colitis in mice. *Exp Ther Med*. 2016;12:2554-62. <https://doi.org/10.3892/etm.2016.3686>
11. Lopetuso LR, Scaldaferri F, Franceschi F, Gasbarrini A. *Bacillus clausii* and gut homeostasis: state of the art and future perspectives. *Expert Rev Gastroenterol Hepatol*. 2016;10:943-8. <https://doi.org/10.1080/17474124.2016.1200465>
12. Secaira-Morochó H, Castillo JA, Driks A. Diversity and evolutionary dynamics of spore-coat

- proteins in spore-forming species of Bacillales. *Microb Genomics*. 2020;6:1-32. <https://doi.org/10.1099/mgen.0.000451>
13. Elshaghabee FMF, Rokana N, Gulhane RD, Sharma C, Panwar H. *Bacillus* as potential probiotics: status, concerns, and future perspectives. *Front Microbiol*. 2017;8:1490. <https://doi.org/10.3389/fmicb.2017.01490>
 14. Payne J, Bellmer D, Jadeja R, Muriana P. The potential of *Bacillus* species as probiotics in the food industry: a review. *Foods*. 2024;13:2444. <https://doi.org/10.3390/foods13152444>
 15. Larsen N, Thorsen L, Kpikpi EN, Stuer-Lauridsen B, Cantor MD, Nielsen B, et al. Characterization of *Bacillus* spp. strains for use as probiotic additives in pig feed. *Appl Microbiol Biotechnol*. 2014;98(3):1105-18. <https://doi.org/10.1007/s00253-013-5343-6>
 16. Hu Y, Dun Y, Li S, Zhao S, Peng N, Liang Y. Effects of *Bacillus subtilis* KN-42 on Growth Performance, Diarrhea and Faecal Bacterial Flora of Weaned Piglets. *Asian-Australas J Anim Sci*. 2014;27:1131-40. <https://doi.org/10.5713/ajas.2013.13737>
 17. Deng B, Wu J, Li X, Zhang C, Men X, Xu Z. Effects of *Bacillus subtilis* on growth performance, serum parameters, digestive enzyme, intestinal morphology, and colonic microbiota in piglets. *AMB Express*. 2020;10:212. <https://doi.org/10.1186/s13568-020-01150-z>
 18. Tian Z, Wang X, Duan Y, Zhao Y, Zhang W, Azad MAK, et al. Dietary supplementation with *Bacillus subtilis* promotes growth and gut health of weaned piglets. *Front Vet Sci*. 2020;7:600772. <https://doi.org/10.3389/fvets.2020.600772>
 19. Xie Z, Li M, Qian M, Yang Z, Han X. Co-cultures of *Lactobacillus acidophilus* and *Bacillus subtilis* enhance mucosal barrier by modulating gut microbiota-derived short-chain fatty acids. *Nutrients*. 2022;14:4475. <https://doi.org/10.3390/nu14214475>
 20. Zhang Y, Shi C, Wang C, Lu Z, Wang F, Feng J, et al. Effect of soybean meal fermented with *Bacillus subtilis* BS12 on growth performance and small intestinal immune status of piglets. *Food Agric Immunol*. 2017;29:133-46. <https://doi.org/10.1080/09540105.2017.1360258>
 21. Huang Y, Łężyk M, Herbst FA, Busk PK, Lange L. Novel keratinolytic enzymes, discovered from a talented and efficient bacterial keratin degrader. *Sci Rep*. 2020;10:10033. <https://doi.org/10.1038/s41598-020-66792-2>
 22. Wang C, Niu D, Mchunu NP, Zhang M, Singh S, Wang Z. Secretory expression of amylosucrase in *Bacillus licheniformis* through twin-arginine translocation pathway. *J Ind Microbiol Biotechnol*. 2024;51:kuae004. <https://doi.org/10.1093/jimb/kuae004>
 23. Fu G, Ma J, Wang L, Yang X, Liu J, Zhao X. Effect of degradation of zearalenone-contaminated feed by *Bacillus licheniformis* CK1 on postweaning female piglets. *Toxins*. 2016;8:300. <https://doi.org/10.3390/toxins8100300>
 24. Horng YB, Yu YH, Dybus A, Hsiao FSH, Cheng YH. Antibacterial activity of *Bacillus* species-derived surfactin on *Brachyspira hyodysenteriae* and *Clostridium perfringens*. *AMB Express*. 2019;9:188. <https://doi.org/10.1186/s13568-019-0914-2>
 25. Hung DY, Cheng YH, Chen WJ, Hua KF, Pietruszka A, Dybus A, et al. *Bacillus licheniformis*-fermented products reduce diarrhea incidence and alter the fecal microbiota community in weaning piglets. *Animals*. 2019;9:1145. <https://doi.org/10.3390/ani9121145>
 26. Zong X, Wang TH, Lu ZQ, Song DG, Zhao J, Wang YZ. Effects of *Clostridium butyricum* or in combination with *Bacillus licheniformis* on the growth performance, blood indexes, and intestinal barrier function of weanling piglets. *Livest Sci*. 2019;220:137-42. <https://doi.org/10.1016/j.livsci.2018.12.024>
 27. Lan R, Kim IH. Effects of *Bacillus licheniformis* and *Bacillus subtilis* complex on growth performance and faecal noxious gas emissions in growing-finishing pigs. *J Sci Food Agric*. 2019;99:1554-60. <https://doi.org/10.1002/jsfa.9333>

28. Kim YJ, Cho SB, Song MH, Lee SI, Hong SM, Yun W, et al. Effects of different *Bacillus licheniformis* and *Bacillus subtilis* ratios on nutrient digestibility, fecal microflora, and gas emissions of growing pigs. *J Anim Sci Technol.* 2022;64:291-301. <https://doi.org/10.5187/jast.2022.e12>
29. He Y, Kim K, Kovanda L, Jinno C, Song M, Chase J, et al. *Bacillus subtilis*: a potential growth promoter in weaned pigs in comparison to carbadox. *J Anim Sci.* 2020;98:skaa290. <https://doi.org/10.1093/jas/skaa290>
30. Luise D, Bertocchi M, Motta V, Salvarani C, Bosi P, Luppi A, et al. *Bacillus* sp. probiotic supplementation diminish the *Escherichia coli* F4ac infection in susceptible weaned pigs by influencing the intestinal immune response, intestinal microbiota and blood metabolomics. *J Anim Sci Biotechnol.* 2019;10:74. <https://doi.org/10.1186/s40104-019-0380-3>
31. Pang J, Liu Y, Kang L, Ye H, Zang J, Wang J, et al. *Bifidobacterium animalis* promotes the growth of weaning piglets by improving intestinal development, enhancing antioxidant capacity, and modulating gut microbiota. *Appl Environ Microbiol.* 2022;88:e01296-22. <https://doi.org/10.1128/aem.01296-22>
32. Majlesi M, Shekarfroush SS, Ghaisari HR, Nazifi S, Sajedianfard J, Eskandari MH. Effect of probiotic *Bacillus coagulans* and *Lactobacillus plantarum* on alleviation of mercury toxicity in rat. *Probiotics Antimicrob Proteins.* 2017;9:300-9. <https://doi.org/10.1007/s12602-016-9250-x>
33. Wu J, Wang J, Lin Z, Liu C, Zhang Y, Zhang S, et al. *Clostridium butyricum* alleviates weaned stress of piglets by improving intestinal immune function and gut microbiota. *Food Chem.* 2023;405:135014. <https://doi.org/10.1016/j.foodchem.2022.135014>
34. Shin D, Chang SY, Bogere P, Won K, Choi JY, Choi YJ, et al. Beneficial roles of probiotics on the modulation of gut microbiota and immune response in pigs. *PLoS ONE.* 2019;14:e0220843. <https://doi.org/10.1371/journal.pone.0220843>
35. Castro J, Barros MM, Araujo D, Campos AM, Oliveira R, Silva S, et al. Swine enteric colibacillosis: current treatment avenues and future directions. *Front Vet Sci.* 2022;9:981207. <https://doi.org/10.3389/fvets.2022.981207>
36. Tang X, Xiong K, Fang R, Li M. Weaning stress and intestinal health of piglets: a review. *Front Immunol.* 2022;13:1042778. <https://doi.org/10.3389/fimmu.2022.1042778>
37. Kim JE, Li B, Fei L, Horne R, Lee D, Loe AKH, et al. Gut microbiota promotes stem cell differentiation through macrophage and mesenchymal niches in early postnatal development. *Immunity.* 2023;56:2175. <https://doi.org/10.1016/j.immuni.2023.08.005>
38. Yang H, Xiong X, Wang X, Tan B, Li T, Yin Y. Effects of weaning on intestinal upper villus epithelial cells of piglets. *PLoS ONE.* 2016;11:e0150216. <https://doi.org/10.1371/journal.pone.0150216>
39. Dalziel JE, Spencer NJ, Young W. Microbial signalling in colonic motility. *Int J Biochem Cell Biol.* 2021;134:105963. <https://doi.org/10.1016/j.biocel.2021.105963>
40. Zhang Y, Xia Y, Sun J. Probiotics and microbial metabolites maintain barrier and neuromuscular functions and clean protein aggregation to delay disease progression in TDP43 mutation mice. *Gut Microbes.* 2024;16:2363880. <https://doi.org/10.1080/19490976.2024.2363880>
41. Kuo WT, Odenwald MA, Turner JR, Zuo L. Tight junction proteins occludin and ZO-1 as regulators of epithelial proliferation and survival. *Ann N Y Acad Sci.* 2022;1514:21-33. <https://doi.org/10.1111/nyas.14798>